

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
24 February 2005 (24.02.2005)

PCT

(10) International Publication Number
WO 2005/016894 A1

(51) International Patent Classification⁷: **C07D 239/48**,
405/12, 403/12, 401/12, 401/14, A61K 31/506, 35/00,
A61P 37/00, 29/00, C07D 401/14, 405/14

(21) International Application Number:
PCT/EP2004/009099

(22) International Filing Date: 13 August 2004 (13.08.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0319227.5 15 August 2003 (15.08.2003) GB
0322370.8 24 September 2003 (24.09.2003) GB

(71) Applicant (for all designated States except AT, US): **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(71) Applicant (for AT only): **NOVARTIS PHARMA GMBH** [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).

(71) Applicant (for all designated States except US): **IRM LLC**; Sofia House, 48 Church Street, Hamilton HM XL (BM).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GARCIA-ECHEVERRIA, Carlos** [ES/CH]; Engalgasse 126, CH-4052 Basel (CH). **KANAZAWA, Takanori** [JP/JP]; 27-8-203, Higashi-arai, Tsukuba-shi, Ibaraki Pref. 305-0033 (JP). **KAWAHARA, Eiji** [JP/JP]; 4-20, Inarimae, Tsukuba-shi, Ibaraki Pref. 305-0061 (JP). **MASUYA, Keiichi** [JP/JP]; 1-2-1 Tsukuho, Tsukuba-shi, Ibaraki Pref. 300-3257 (JP). **MATSUURA, Naoko** [JP/JP]; 2-4-1-507, Amakubo, Tsukuba-shi, Ibaraki Pref. 305-0005 (JP). **MIYAKE, Takahiro** [JP/JP]; 2-4-1-102, Amakubo, Tsukuba-shi, Ibaraki Pref. 305-0005 (JP). **OHMORI, Osamu** [JP/JP]; 2-4-6-406, Sengen, Tsukuba-shi, Ibaraki Pref. 305-0047 (JP). **UMEMURA, Ichiro** [JP/JP]; 2-3-7-406, Ninomiya,

Tsukuba-shi, Ibaraki Pref. 305-0051 (JP). **STEENSMA, Ruo** [CN/US]; 6509 Avenida Manana, La Jolla, CA 92037 (US). **CHOPIUK, Greg** [CA/US]; 10994 West Ocean Air Drive No.387, San Diego, CA 92130 (US). **JIANG, Jiqing** [CN/US]; 5225 Fiore Terrace No.D104, San Diego, CA 92122 (US). **WAN, Yongqin** [CN/US]; 7552 Charmant Drive, No.1622, San Diego, CA 92122 (US). **DING, Qiang** [CN/US]; 8465 Regents Road No 264, San Diego, CA 92122 (US). **ZHANG, Qiong** [CN/US]; 8148 Genesee Ave., No.106, San Diego, CA 92122 (US). **GRAY, Nathanael, Schiander** [US/US]; 5652 Lamas Street, San Diego, CA 92122 (US). **KARANEWSKY, Donald** [US/US]; 1797 Continental Lane, Escondido, CA 92029 (US).

(74) Agent: **GRUBB, Philip**; Novartis AG, Corporate Intellectual Property, CH-4002 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 2, 4-PYRIMIDINEDIAMINES USEFUL IN THE TREATMENT OF NEOPLASTIC DISEASES, INFLAMMATORY AND IMMUNE SYSTEM DISORDERS



(I)


(57) Abstract: Novel pyrimidine derivatives of formula (I) Wherein R is selected from C₁₆₋₁₀ aryl, C₅₋₁₀heteroaryl, C₃₋₁₂cycloalkyl and C₃₋₁₀heterocycloalkyl; R⁰-R⁶ as described herein; and their use for the manufacture of a medicament for the treatment or prevention of a disease which responds to inhibition of FAK and/or ALK and/or ZAP-70 and/or IGF-IR.

WO 2005/016894 A1

2,4-PYRIMIDINEDIAMINES USEFUL IN THE TREATMENT OF NEOPLASTIC DISEASES,
INFLAMMATORY AND IMMUNE SYSTEM DISORDERS.

The present invention relates to the use novel pyrimidine derivatives, the certain novel pyrimidine derivatives, to processes for their production, their use as pharmaceuticals and to pharmaceutical compositions comprising them.

More particularly the present invention provides in a first aspect, the use of a compound of formula I



(I)

wherein

R is selected from C₆₋₁₀aryl, C₅₋₁₀heteroaryl, C₃₋₁₂cycloalkyl and C₃₋₁₀heterocycloalkyl;

each of R⁰, R¹, R², and R³ independently is hydrogen, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkinyl, C₃₋₈cycloalkyl, C₃₋₈cycloalkylC₁₋₈alkyl, C₅₋₁₀arylC₁₋₈alkyl, hydroxyC₁₋₈alkyl, C₁₋₈alkoxyC₁₋₈alkyl, aminoC₁₋₈alkyl, haloC₁₋₈alkyl, unsubstituted or substituted C₅₋₁₀aryl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1, 2 or 3 hetero atoms selected from N, O and S, hydroxy, C₁₋₈alkoxy, hydroxyC₁₋₈alkoxy, C₁₋₈alkoxyC₁₋₈alkoxy, haloC₁₋₈alkoxy, unsubstituted or substituted C₅₋₁₀arylC₁₋₈alkoxy, unsubstituted or substituted heterocyclyloxy, or unsubstituted or substituted heterocyclylC₁₋₈alkoxy, unsubstituted or substituted amino, C₁₋₈alkylthio, C₁₋₈alkylsulfinyl, C₁₋₈alkylsulfonyl, C₅₋₁₀arylsulfonyl, halogen, carboxy, C₁₋₈alkoxycarbonyl, unsubstituted or substituted carbamoyl, unsubstituted or substituted sulfamoyl, cyano, nitro, -S(O)₀₋₂NR₁₂R₁₃, -S(O)₀₋₂R₁₃, -NR₁₂S(O)₀₋₂R₁₃, -C(O)NR₁₂R₁₃, -C(O)R₁₃ and -C(O)OR₁₃; wherein R₁₂ is selected from hydrogen and C₁₋₆alkyl; and R₁₃ is selected from hydrogen, C₁₋₆alkyl and C₃₋₁₂cycloalkyl;

or R⁰ and R¹, R¹ and R², and/or R² and R³ form, together with the carbon atoms to which they are attached, a 5 or 6 membered carbocyclic or heterocyclic ring comprising 0, 1, 2 or 3 heteroatoms selected from N, O and S;

R⁴ is hydrogen or C₁₋₈alkyl;

each of R^5 and R^6 independently is hydrogen, C_1 - C_8 alkyl, C_1 - C_8 alkoxy C_1 - C_8 alkyl, halo C_1 - C_8 alkyl, C_1 - C_8 alkoxy, halogen, carboxy, C_1 - C_8 alkoxycarbonyl, unsubstituted or substituted carbamoyl, cyano, or nitro;

R is unsubstituted or substituted by R_7 , R_8 , R_9 , R_{10} , and R'_{10} ;

R_7 , R_8 , R_9 , R_{10} , or R'_{10} is a substituent independently selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkinyl, C_3 - C_8 cycloalkyl, C_3 - C_8 cycloalkyl C_1 - C_8 alkyl, C_5 - C_{10} aryl C_1 - C_8 alkyl, hydroxy C_1 - C_8 alkyl, C_1 - C_8 alkoxy C_1 - C_8 alkyl, amino C_1 - C_8 alkyl, halo C_1 - C_8 alkyl, unsubstituted or substituted C_5 - C_{10} aryl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1, 2 or 3 hetero atoms selected from N, O and S, hydroxy, C_1 - C_8 alkoxy, hydroxy C_1 - C_8 alkoxy, C_1 - C_8 alkoxy C_1 - C_8 alkoxy, halo C_1 - C_8 alkoxy, unsubstituted or substituted amino C_1 - C_8 alkoxy, unsubstituted or substituted C_5 - C_{10} aryl C_1 - C_8 alkoxy, unsubstituted or substituted heterocyclyloxy, or unsubstituted or substituted heterocyclyl C_1 - C_8 alkyl, unsubstituted or substituted heterocyclyl C_1 - C_8 alkoxy, unsubstituted or substituted amino, C_1 - C_8 alkylthio, C_1 - C_8 alkylsulfinyl, C_1 - C_8 alkylsulfonyl, C_5 - C_{10} arylsulfonyl, heterocyclosulfonyl, halogen, carboxy, C_1 - C_8 alkylcarbonyl, C_1 - C_8 alkoxycarbonyl, unsubstituted or substituted carbamoyl, unsubstituted or substituted sulfamoyl, cyano, nitro, $-S(O)_{0-2}NR_{12}R_{13}$, $-S(O)_{0-2}R_{12}$, $-C(O)R_{11}$, $-OXR_{11}$, $-NR_{12}XR_{11}$, $-NR_{12}XNR_{12}R_{13}$, $-OXNR_{12}R_{13}$, $-OXOR_{12}$ and $-XR_{11}$;

or two adjacent substituents on R may form together with the carbon atoms to which they are attached, a unsubstituted or substituted 5 or 6 membered carbocyclic or heterocyclic ring comprising 0, 1, 2 or 3 heteroatoms selected from N, O and S;

X is a bond or C_{1-6} alkylene; and

R_{11} is independently selected from C_{6-10} aryl, C_{5-10} heteroaryl, C_{3-12} cycloalkyl and C_{3-10} heterocycloalkyl;

and any aryl, heteroaryl, cycloalkyl or heterocycloalkyl of R_{11} is optionally substituted by 1 to 3 radicals independently selected from C_{1-6} alkyl, C_{3-10} heterocycloalkyl- C_{0-4} alkyl optionally substituted with C_{1-6} alkyl, $-C(O)R_{12}$, $-C(O)NR_{12}R_{13}$, $-XNR_{12}R_{13}$, $-NR_{12}XNR_{12}R_{13}$ and $-NR_{12}C(O)R_{13}$; wherein X is a bond or C_{1-6} alkylene; R_{12} and R_{13} are independently selected from hydrogen and C_{1-6} alkyl;

and salts thereof in the treatment of of a disease associated to tyrosine kinase activity of anaplastic lymphoma kinase (ALK) or for the manufacture of pharmaceutical compositions for use in the treatment of said diseases, as well as use in the treatment of said diseases, to methods of use of such pyrimidine derivatives in the treatment of said diseases, and to pharmaceutical compositions comprising such pyrimidine derivatives for the treatment of said diseases.

The general terms used hereinbefore and hereinafter preferably have within the context of this disclosure the following meanings, unless otherwise indicated:

Where the plural form is used for compounds, salts, and the like, this is taken to mean also a single compound, salt, or the like.

Any asymmetric carbon atoms may be present in the (R)-, (S)- or (R,S)-configuration, preferably in the (R)- or (S)-configuration. The compounds may thus be present as mixtures of isomers or as pure isomers, preferably as enantiomer-pure diastereomers.

The invention relates also to possible tautomers of the compounds of formula I.

C₁-C₈alkyl denotes a an alkyl radical having from 1 up to 8, especially up to 4 carbon atoms, the radicals in question being either linear or branched with single or multiple branching; preferably, C₁-C₈alkyl is butyl, such as n-butyl, sec-butyl, isobutyl, tert-butyl, propyl, such as n-propyl or isopropyl, ethyl or methyl; especially methyl, propyl or tert-butyl.

C₂-C₈alkenyl denotes a an alkenyl radical having from 2 up to 8, especially up to 5 carbon atoms, the radicals in question being either linear or branched with single or multiple branching; preferably, C₂-C₈alkenyl is pentenyl, such as 3-methyl-2-buten-2-yl, butenyl, such as 1- or 2-butenyl or 2-buten-2-yl, propenyl, such as 1-propenyl or allyl, or vinyl.

C₂-C₈alkinyl denotes a an alkynyl radical having from 2 up to 8, especially up to 5 carbon atoms, the radicals in question being either linear or branched; preferably, C₂-C₈alkinyl is propinyl, such as 1-propinyl or propargyl, or acetylenyl.

C₃-C₈cycloalkyl denotes a cycloalkyl radical having from 3 up to 8 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl, preferably cyclopropyl, cyclopentyl or cyclohexyl.

C₁-C₈alkoxy is especially methoxy, ethoxy, isopropoxy, or tert-butoxy.

HydroxyC₁-C₈alkyl is especially hydroxymethyl, 2-hydroxyethyl or 2-hydroxy-2-propyl.

HydroxyC₁-C₈alkoxy is especially 2-hydroxyethoxy or 3-hydroxypropoxy.

C₁-C₈alkoxyC₁-C₈alkoxy is especially 2-methoxyethoxy.

C₁-C₈alkoxyC₁-C₈alkyl is especially methoxymethyl, 2-methoxyethyl or 2-ethoxyethyl.

Halogen is preferably fluorine, chlorine, bromine, or iodine, especially fluorine, chlorine, or bromine.

HaloC₁-C₈alkyl is preferably chloroC₁-C₈alkyl or fluoroC₁-C₈alkyl, especially trifluoromethyl or pentafluoroethyl.

HaloC₁-C₈alkoxy is preferably chloroC₁-C₈alkoxy or fluoroC₁-C₈alkoxy, especially trifluoromethoxy.

C₁-C₈alkoxycarbonyl is especially tert-butoxycarbonyl, iso-propoxycarbonyl, methoxycarbonyl or ethoxycarbonyl.

Unsubstituted or substituted carbamoyl is carbamoyl substituted by one or two substituents selected from hydrogen, C₁-C₈alkyl, C₂-C₈alkenyl, C₂-C₈alkinyl, C₃-C₈cycloalkyl, C₃-C₈cycloalkylC₁-C₈alkyl, C₅-C₁₀arylC₁-C₈alkyl, hydroxyC₁-C₈alkyl, C₁-C₈alkoxyC₁-C₈alkyl, haloC₁-C₈alkyl, unsubstituted or substituted C₅-C₁₀aryl, or aminoC₁-C₈alkyl, or carbamoyl wherein the substituents and the nitrogen atom of the carbamoyl group represent a 5 or 6 membered heterocyclyl further comprising 0, 1 or 2 hetero atoms selected from N, O and S; and is preferably carbamoyl, methylcarbamoyl, dimethylcarbamoyl, propylcarbamoyl, hydroxyethyl-methyl-carbamoyl, di(hydroxyethyl)carbamoyl, dimethylaminoethylcarbamoyl, or pyrrolidinocarbonyl, piperidinocarbonyl, N-methylpiperazinocarbonyl or morpholinocarbonyl, especially carbamoyl or dimethylcarbamoyl.

Unsubstituted or substituted sulfamoyl is sulfamoyl substituted by one or two substituents selected from hydrogen, C₁-C₈alkyl, C₂-C₈alkenyl, C₂-C₈alkinyl, C₃-C₈cycloalkyl, C₃-C₈cycloalkylC₁-C₈alkyl, C₅-C₁₀arylC₁-C₈alkyl, hydroxyC₁-C₈alkyl, C₁-C₈alkoxyC₁-C₈alkyl, haloC₁-

C₈alkyl, unsubstituted or substituted C₅-C₁₀aryl, or aminoC₁-C₈alkyl, or sulfamoyl wherein the substituents and the nitrogen atom of the sulfamoyl group represent a 5 or 6 membered heterocyclyl further comprising 0, 1 or 2 hetero atoms selected from N, O and S; and is preferably sulfamoyl, methylsulfamoyl, propylsulfamoyl, cyclopropylmethyl-sulfamoyl, 2,2,2-trifluoroethylsulfamoyl, dimethylaminoethylsulfamoyl, dimethylsulfamoyl, hydroxyethyl-methyl-sulfamoyl, di(hydroxyethyl)sulfamoyl, or pyrrolidinosulfonyl, piperidinosulfonyl, N-methylpiperazinosulfonyl or morpholinosulfonyl, especially sulfamoyl or methylsulfamoyl.

Unsubstituted or substituted amino is amino substituted by one or two substituents selected from hydrogen, C₁-C₈alkyl, C₂-C₈alkenyl, C₂-C₈alkinyl, C₃-C₈cycloalkyl, C₃-C₈cycloalkylC₁-C₈alkyl, C₅-C₁₀arylC₁-C₈alkyl, hydroxyC₁-C₈alkyl, C₁-C₈alkoxyC₁-C₈alkyl, haloC₁-C₈alkyl, unsubstituted or substituted C₅-C₁₀aryl, aminoC₁-C₈alkyl, acyl, e.g. formyl, C₁-C₈alkylcarbonyl, C₅-C₁₀arylcarbonyl, C₁-C₈alkylsulfonyl or C₅-C₁₀arylsulfonyl, and is preferably amino, methylamino, dimethylamino, propylamino, benzylamino, hydroxyethyl-methyl-amino, di(hydroxyethyl)amino, dimethylaminoethylamino, acetylamino, acetyl-methyl-amino, benzoylamino, methylsulfonylamino or phenylsulfonylamino, especially amino or dimethylamino.

AminoC₁-C₈alkyl is especially aminoethyl, methylaminoethyl, dimethylaminoethyl or dimethylaminopropyl.

Unsubstituted or substituted C₅-C₁₀aryl is, for example, phenyl, indenyl, indanyl, naphthyl, or 1,2,3,4-tetrahydronaphthalenyl, optionally substituted by C₁-C₈alkyl, C₁-C₈alkoxyC₁-C₈alkyl, haloC₁-C₈alkyl, hydroxy, C₁-C₈alkoxy, methylenedioxy, amino, substituted amino, halogen, carboxy, C₁-C₈alkoxycarbonyl, carbamoyl, sulfamoyl, cyano or nitro; preferably phenyl, tolyl, trifluoromethylphenyl, methoxyphenyl, dimethoxyphenyl, methylenedioxyphenyl, chlorophenyl or bromophenyl, whereby the substituents may be in ortho, meta or para position, preferably meta or para.

C₅-C₁₀aryloxy is especially phenoxy or methoxyphenoxy, e.g. p-methoxyphenoxy.

C₅-C₁₀arylC₁-C₈alkyl is especially benzyl or 2-phenylethyl.

C₅-C₁₀arylC₁-C₈alkoxy is especially benzyloxy or 2-phenylethoxy.

Unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1, 2 or 3 hetero atoms selected from N, O and S may be unsaturated, partially unsaturated or saturated, and further condensed to a benzo group or a 5 or 6 membered heterocyclyl group, and may be bound through a hetero or a carbon atom, and is, for example, pyrrolyl, indolyl, pyrrolidinyl, imidazolyl, benzimidazolyl, pyrazolyl, triazolyl, benzotriazolyl, tetrazolyl, pyridyl, quinolinyl, isoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, piperidyl, pyrimidinyl, pyrazinyl, piperazinyl, purinyl, tetrazinyl, oxazolyl, isoxalyl, morpholinyl, thiazolyl, benzothiazolyl, oxadiazolyl, and benzoxadiazolyl. Substituents considered are C₁-C₈alkyl, hydroxyC₁-C₈alkyl, C₁-C₈alkoxyC₁-C₈alkyl, C₁-C₈alkoxyC₁-C₈alkoxy, haloC₁-C₈alkyl, hydroxy, amino, substituted amino, C₁-C₈alkoxy, halogen, carboxy, C₁-C₈alkylcarbonyl, C₁-C₈alkoxycarbonyl, carbamoyl, C₁-C₈alkylcarbamoyl, cyano, oxo, or unsubstituted or substituted 5 or 6 membered heterocyclyl as defined in this paragraph. 5 or 6 membered heterocyclyl preferably comprises 1 or 2 hetero atoms selected from N, O and S, and is especially indolyl, pyrrolidinyl, pyrrolidonyl, imidazolyl, N-methylimidazolyl, benzimidazolyl, S,S-dioxisothiazolidinyl, piperidyl, 4-acetylaminopiperidyl, 4-methylcarbamoylpiperidyl, 4-piperidinopiperidyl, 4-cyanopiperidyl, piperazinyl, N-methylpiperazinyl, N-(2-hydroxyethyl)piperazinyl, morpholinyl, 1-aza-2,2-dioxo-2-thiacyclohexyl, or sulfolanyl.

In unsubstituted or substituted heterocyclyloxy, heterocyclyl has the meaning as defined above, and is especially N-methyl-4-piperidyloxy. In unsubstituted or substituted heterocyclylC₁-C₈alkoxy, heterocyclyl has the meaning as defined above, and is especially 2-pyrrolidinoethoxy, 2-morpholinoethoxy, 3-morpholinopropoxy, 1-methyl-piperidin-3-ylmethoxy, 3-(N-methylpiperazino)propoxy or 2-(1-imidazolyl)ethoxy.

In a 5 or 6 membered carbocyclic or heterocyclic ring comprising 0, 1, 2 or 3 heteroatoms selected from N, O and S, and formed by two adjacent substituents together with the benzene ring, the ring may be further substituted, e.g. by C₁-C₈alkyl, C₁-C₈alkoxy, haloC₁-C₈alkyl, hydroxy, amino, substituted amino, C₁-C₈alkoxy, halogen, carboxy, C₁-C₈alkoxycarbonyl, carbamoyl, cyano, or oxo. The two adjacent substituents forming such a ring are preferably propylene, butylene, 1-aza-2-propylidene, 3-aza-1-propylidene, 1,2-diaza-2-propylidene, 2,3-diaza-1-propylidene, 1-oxapropylene, 1-oxapropylidene, methylenedioxy, difluoromethylenedioxy, 2-aza-1-oxopropylene, 2-aza-2-methyl-1-oxopropylene, 1-aza-2-oxopropylene, 2-aza-1,1-dioxo-1-thiapropylene or the corresponding butylene derivatives forming a 6 membered ring.

Salts are especially the pharmaceutically acceptable salts of compounds of formula I.

Such salts are formed, for example, as acid addition salts, preferably with organic or inorganic acids, from compounds of formula I with a basic nitrogen atom, especially the pharmaceutically acceptable salts. Suitable inorganic acids are, for example, halogen acids, such as hydrochloric acid, sulfuric acid, or phosphoric acid. Suitable organic acids are, for example, carboxylic, phosphonic, sulfonic or sulfamic acids, for example acetic acid, propionic acid, octanoic acid, decanoic acid, dodecanoic acid, glycolic acid, lactic acid, fumaric acid, succinic acid, adipic acid, pimelic acid, suberic acid, azelaic acid, malic acid, tartaric acid, citric acid, amino acids, such as glutamic acid or aspartic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, cyclohexanecarboxylic acid, adamantanecarboxylic acid, benzoic acid, salicylic acid, 4-aminosalicylic acid, phthalic acid, phenylacetic acid, mandelic acid, cinnamic acid, methane- or ethane-sulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 2-naphthalenesulfonic acid, 1,5-naphthalene-disulfonic acid, 2-, 3- or 4-methylbenzenesulfonic acid, methylsulfuric acid, ethylsulfuric acid, dodecylsulfuric acid, N-cyclohexylsulfamic acid, N-methyl-, N-ethyl- or N-propyl-sulfamic acid, or other organic protonic acids, such as ascorbic acid.


For isolation or purification purposes it is also possible to use pharmaceutically unacceptable salts, for example picrates or perchlorates. For therapeutic use, only pharmaceutically acceptable salts or free compounds are employed (where applicable in the form of pharmaceutical preparations), and these are therefore preferred.

In view of the close relationship between the novel compounds in free form and those in the form of their salts, including those salts that can be used as intermediates, for example in the purification or identification of the novel compounds, any reference to the free compounds hereinbefore and hereinafter is to be understood as referring also to the corresponding salts, as appropriate and expedient.

The compounds of formula I have valuable pharmacological properties, as described hereinbefore and hereinafter.

In formula I the following significances are preferred independently, collectively or in any combination or sub-combination. R is C₆₋₁₀aryl, C₅₋₁₀heteroaryl, C₃₋₁₂cycloalkyl or C₃₋₁₀heterocycloalkyl, preferably R is

- 8 -



wherein R_7 , R_8 , R_9 , R_{10} , or R'_{10} are as defined above;

In each of the following significances A, D or E is C or N but A, D and E may not all be N, preferably A, D or E is C:

- (a) each of R^0 or R^2 independently is hydrogen, C_1 - C_8 alkyl, e.g. methyl, ethyl or isopropyl, hydroxy C_1 - C_8 alkyl, e.g. hydroxyethyl or hydroxybutyl, halo C_1 - C_8 alkyl, e.g. trifluoromethyl, unsubstituted or substituted C_5 - C_{10} aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C_1 - C_8 alkoxy, e.g. methoxy, ethoxy or isopropoxy, halo C_1 - C_8 alkoxy, e.g. trifluoromethoxy, C_5 - C_{10} aryloxy, e.g. phenoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclyl C_1 - C_8 alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetylamino, C_1 - C_8 alkylsulfonyl, e.g. methylsulfonyl, halogen, e.g. fluoro or chloro, unsubstituted or substituted carbamoyl, e.g. cyclohexylcarbamoyl, piperidinocarbonyl, piperazinocarbonyl, N-methylpiperazinocarbonyl or morpholinocarbonyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl; preferably hydrogen, piperazino, N-methylpiperazino or 1-methyl-4-piperidyloxy, $-S(O)_{0-2}NR_{12}R_{13}$, $-S(O)_{0-2}R_{13}$, $-NR_{12}S(O)_{0-2}R_{13}$, $-C(O)NR_{12}R_{13}$, and $-C(O)OR_{13}$ in particular hydrogen;
- (b) R^1 is hydrogen, C_1 - C_8 alkyl, e.g. methyl, ethyl or isopropyl, hydroxy C_1 - C_8 alkyl, e.g. hydroxyethyl or hydroxybutyl, halo C_1 - C_8 alkyl, e.g. trifluoromethyl, unsubstituted or substituted C_5 - C_{10} aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C_1 - C_8 alkoxy, e.g. methoxy, ethoxy or isopropoxy, halo C_1 - C_8 alkoxy, e.g. trifluoromethoxy, C_5 - C_{10} aryloxy, e.g. phenoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclyl C_1 - C_8 alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetylamino, C_1 - C_8 alkylsulfonyl, e.g. methylsulfonyl, halogen, e.g. fluoro or chloro,

unsubstituted or substituted carbamoyl, e.g. cyclohexylcarbamoyl, piperidinocarbonyl, piperazinocarbonyl, N-methylpiperazinocarbonyl or morpholinocarbonyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl; preferably hydrogen, piperazino, N-methylpiperazino, morpholino, 1-methyl-4-piperidinyloxy, 3-morpholinopropoxy or 2-morpholinoethoxy, in particular hydrogen;

(c) R^3 is hydrogen, C_1 - C_8 alkyl, e.g. methyl or ethyl, hydroxy C_1 - C_8 alkyl, e.g. hydroxyethyl or hydroxybutyl, halo C_1 - C_8 alkyl, e.g. trifluoromethyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 heteroatoms selected from N, O and S, e.g. 2-pyrrolidonyl or S,S-dioxoisothiazolidinyl, C_1 - C_8 alkoxy, e.g. methoxy, substituted amino, e.g. acetylamino, acetyl-methyl-amino, benzoylamino, methylsulfonylamino or phenylsulfonylamino, C_1 - C_8 alkylsulfonyl, e.g. methylsulfonyl, propyl-sulfonyl, cyclohexyl-sulfonyl, isopropyl-sulfonyl, C_5 - C_{10} arylsulfonyl, e.g. phenylsulfonyl, halogen, e.g. fluoro or chloro, carboxy, substituted or unsubstituted carbamoyl, e.g. carbamoyl, methylcarbamoyl, ethyl-amino-carbonyl or dimethylcarbamoyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl, propylsulfamoyl, isopropylsulfamoyl, isobutylsulfamoyl, cyclopropylmethyl-sulfamoyl, 2,2,2-trifluoroethylsulfamoyl, dimethylsulfamoyl or morpholinosulfonyl dimethyl-sulfamoyl, ethyl-sulfamoyl, 1-ethyl-propyl-sulfamoyl, cyclopentyl-sulfamoyl, cyclobutyl-sulfamoyl; preferably sulfamoyl, methylsulfamoyl or propylsulfamoyl;

(d) each pair of adjacent substituents R^0 and R^1 , or R^1 and R^2 , or R^2 and R^3 are $-CH_2-NH-CO-$, $-CH_2-CH_2-NH-CO-$, $-CH_2-CO-NH-$, $-CH_2-CH_2-CO-NH-$, $-CH_2-NH-SO_2-$, $-CH_2-CH_2-NH-SO_2-$, $-CH_2-SO_2-NH-$, $-CH_2-CH_2-SO_2-NH-$, $-CH_2-CH_2-SO_2-$, $-CH_2-CH_2-CH_2-SO_2-$, $-O-CH_2-O-$, or $-O-CF_2-O-$, and such pairs wherein hydrogen in NH is replaced by C_1 - C_8 alkyl; preferably the pair of adjacent substituents R^0 and R^1 , or R^1 and R^2 being $-O-CH_2-O-$, and the pair of adjacent substituents R^2 and R^3 being $-CH_2-NH-CO-$ or $-CH_2-NH-SO_2-$.

(e) R^4 is hydrogen or C_1 - C_8 alkyl, e.g. methyl; preferably hydrogen;

(f) R^5 is hydrogen; C_1 - C_8 alkyl, e.g. methyl or ethyl, halogen, e.g. chloro or bromo, halo C_1 - C_8 alkyl, e.g. trifluoromethyl, cyano or nitro; preferably hydrogen, methyl, ethyl, chloro, bromo, trifluoromethyl or nitro; in particular chloro or bromo;

(g) R^6 is hydrogen;

(h) each of R^7 and R^9 independently is hydrogen, C_1 - C_8 alkyl, e.g. methyl, ethyl or isopropyl, hydroxy C_1 - C_8 alkyl, e.g. hydroxyethyl or hydroxybutyl, C_1 - C_8 alkylcarbonyl, e.g. methyl carbonyl, aminoalkoxy, e.g. diethylaminoethoxy, halo C_1 - C_8 alkyl, e.g. trifluoromethyl,

unsubstituted or substituted C₅-C₁₀aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C₁-C₈alkoxy, e.g. methoxy, ethoxy or isopropoxy, haloC₁-C₈alkoxy, e.g. trifluoromethoxy, C₅-C₁₀aryloxy, e.g. phenoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclylC₁-C₈alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetilamino, C₁-C₈alkylsulfonyl, e.g. methylsulfonyl, heterocyclosulfonyl, e.g. piperazinylsulfonyl, heterocyclocarbonyl, e.g.

methylpiperazinylcarbonyl, cyano, halogen, e.g. fluoro or chloro, unsubstituted or substituted carbamoyl, e.g. cyclohexylcarbamoyl, piperidinocarbonyl, piperazinocarbonyl, N-methylpiperazinocarbonyl or morpholinocarbonyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl; preferably hydrogen, methyl, isopropyl, trifluoromethyl, phenyl, methoxyphenyl, piperidino, piperazino, N-methylpiperazino, morpholino, methoxy, ethoxy, isopropoxy, phenoxy, 3-morpholinopropoxy, 2-morpholinoethoxy, 2-(1-imidazolyl)ethoxy, dimethylamino, fluoro, morpholinocarbonyl, piperidinocarbonyl, piperazinocarbonyl or cyclohexylcarbamoyl;

(i) R⁸ is hydrogen, C₁-C₈alkyl, e.g. methyl, ethyl or isopropyl, hydroxyC₁-C₈alkyl, e.g.

hydroxyethyl or hydroxybutyl, haloC₁-C₈alkyl, e.g. trifluoromethyl, C₅-C₁₀aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, heterocyclylalkyl, e.g. methylpiperazinoethyl, heterocyclylcarbonyl, e.g. piperazinocarbonyl, heterocyclyl C₁-C₈alkylamino, e.g. pyridylethyl(methyl)amino, C₁-C₈alkoxy, e.g. methoxy, ethoxy or isopropoxy, haloC₁-C₈alkoxy, e.g. trifluoromethoxy, C₅-C₁₀aryloxy, e.g. phenoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclylC₁-C₈alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino or dimethylamino, C₁-C₈alkylamino-C₁-C₈alkylamino, e.g. dimethylamino-propylamino, C₁-C₈alkylsulfonyl, e.g. methylsulfonyl, halogen, e.g. fluoro or chloro, unsubstituted or substituted carbamoyl, e.g. cyclohexylcarbamoyl, piperidinocarbonyl, piperazinocarbonyl, N-methylpiperazinocarbonyl or morpholinocarbonyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl, cyano, or nitro; preferably hydrogen, methyl, piperidino, piperazino, N-methylpiperazino, morpholino, methoxy, ethoxy, trifluoromethoxy, phenoxy, 1-methyl-4-

- piperidyloxy, 3-morpholinopropoxy, 2-morpholinoethoxy, 3-(N-methylpiperazino)-propoxy, methylamino, fluoro, chloro, sulfamoyl or nitro;
- (j) R^{10} is hydrogen, C_1 - C_8 alkyl, e.g. methyl, ethyl or butyl, hydroxy, cyano, hydroxy C_1 - C_8 alkyl, e.g. hydroxyethyl or hydroxybutyl, halo C_1 - C_8 alkyl, e.g. trifluoromethyl, C_1 - C_8 alkoxy, e.g. methoxy or ethoxy, cycloalkylalkoxy, aryloxy, halo C_1 - C_8 alkoxy, unsubstituted or substituted heterocyclyl C_1 - C_8 alkoxy, e.g. 2-(1-imidazolyl)ethoxy, unsubstituted or substituted amino, e.g. methylamino or dimethylamino, halogen, e.g. fluoro or chloro; carboxy, carbamoyl, or unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl; preferably methyl, butyl, methoxy, ethoxy, 2-(1-imidazolyl)ethoxy, methylamino, dimethylamino or fluoro; and
- (k) each pair of adjacent substituents R^7 and R^8 , or R^8 and R^9 or R^9 and R^{10} , are $-NH-CH=CH-$, $-CH=CH-NH-$, $-NH-N=CH-$, $-CH=N-NH-$, $-CH_2-CH_2-CH_2-$, $-CH_2-CH_2-CH_2-CH_2-$, $-CH_2-CH_2-O-$, $-CH_2C(CH_3)_2O-$, $-CH=C(CH_3)O-$, $-OCH_2CH_2O-$, $-(Morpholinopropyl)N-CH=CH-$, $-CH=CH-O-$, $-O-CH_2-O-$, or $-O-CF_2-O-$; preferably the pair of adjacent substituents R^7 and R^8 or R^8 and R^9 being $-O-CH_2-O-$ or the pair of adjacent substituents R^9 and R^{10} being $-NH-CH=CH-$, $-CH=N-NH-$, $-CH_2-CH_2-CH_2-$, $-CH_2-CH_2-CH_2-CH_2-$ or $-O-CF_2-O-$.
- (l) or R^7 , R^8 , R^9 , R^{10} and R^{10} are ethoxy, ethyl, propyl, methyl, t-butyl, trifluoromethyl, nitrile, cyclobutyloxy, 2,2,2-trifluoroethoxy, methoxy, isobutyloxy, t-butyloxy, isopropoxy, methyl-amino-carbonyl, cyclopropyl-methoxy, dimethylamino-propyl-amino, methoxy-ethoxy, $-XR_{11}$, $-C(O)R_{11}$ and $-OXR_{11}$; wherein X is a bond, methylene or ethylene; R_{11} is selected from piperaziny, piperidiny, pyrrolidiny, morpholino, azepanyl and 1,4-dioxo-8-aza-spiro[4.5]dec-8-yl; wherein R_{11} is optionally substituted by 1 to 3 radicals independently selected from methyl, isopropyl, acetyl, acetyl-methyl-amino, 3-dimethylamino-2,2-dimethyl-propylamino, ethyl-methyl-amino-ethoxy, diethyl-amino-ethoxy, amino-carbonyl, ethyl, 2-oxo-pyrrolidin-1-yl, pyrrolidiny, pyrrolidiny-methyl, piperidiny optionally substituted with methyl or ethyl, morpholino, dimethylamino, dimethylamino-propyl-amino, methyl-amino and ethyl-amino.

More preferred are the following meanings, independently, collectively or in any combination or sub-combination:

- (a') each of R^0 or R^2 independently is hydrogen, C_1 - C_8 alkyl, e.g. methyl, ethyl or isopropyl, halo C_1 - C_8 alkyl, e.g. trifluoromethyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C_1 - C_8 alkoxy, e.g. methoxy, ethoxy or

isopropoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclylC₁-C₈alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetylamino, halogen, e.g. fluoro or chloro; preferably hydrogen, piperazino, N-methylpiperazino or 1-methyl-4-piperidyloxy, in particular hydrogen;

- (b') R¹ is hydrogen, C₁-C₈alkyl, e.g. methyl, ethyl or isopropyl, haloC₁-C₈alkyl, e.g. trifluoromethyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C₁-C₈alkoxy, e.g. methoxy, ethoxy or isopropoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclylC₁-C₈alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetylamino, halogen, e.g. fluoro or chloro; preferably hydrogen, piperazino, N-methylpiperazino, morpholino, 1-methyl-4-piperidyloxy, 3-morpholinopropoxy or 2-morpholinoethoxy, in particular hydrogen;
- (c') R³ is hydrogen, C₁-C₈alkyl, e.g. methyl or ethyl, haloC₁-C₈alkyl, e.g. trifluoromethyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 heteroatoms selected from N, O and S, e.g. 2-pyrrolidinyl or S,S-dioxoisothiazolidinyl, C₁-C₈alkoxy, e.g. methoxy, substituted amino, e.g. acetylamino, acetyl-methyl-amino, benzoylamino, methylsulfonylamino or phenylsulfonylamino, C₁-C₈alkylsulfonyl, e.g. methylsulfonyl, C₅-C₁₀arylsulfonyl, e.g. phenylsulfonyl, halogen, e.g. fluoro or chloro, carboxy, substituted or unsubstituted carbamoyl, e.g. carbamoyl, methylcarbamoyl or dimethylcarbamoyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl, propylsulfamoyl, isopropylsulfamoyl, isobutylsulfamoyl, cyclopropylmethyl-sulfamoyl, 2,2,2-trifluoroethylsulfamoyl, dimethylsulfamoyl or morpholinosulfonyl; preferably sulfamoyl, methylsulfamoyl or propylsulfamoyl;
- (d') each pair of adjacent substituents R⁰ and R¹, or R¹ and R², or R² and R³ are -CH₂-NH-CO-, -CH₂-NH-SO₂-, -CH₂-CH₂-SO₂-, -O-CH₂-O-, or -O-CF₂-O-, and such pairs wherein hydrogen in NH is replaced by C₁-C₈alkyl; preferably the pair of adjacent substituents R⁰ and R¹, or R¹ and R² being -O-CH₂-O-, and the pair of adjacent substituents R² and R³ being -CH₂-NH-CO- or -CH₂-NH-SO₂-.
- (e') R⁴ is hydrogen;


- (f') R⁵ is hydrogen, halogen, e.g. chloro or bromo, haloC₁-C₈alkyl, e.g. trifluoromethyl, or nitro; preferably hydrogen, chloro, bromo, trifluoromethyl or nitro; in particular chloro or bromo;
- (g') R⁶ is hydrogen;
- (h') each of R⁷ and R⁹ independently is hydrogen, C₁-C₈alkyl, e.g. methyl, ethyl or isopropyl, haloC₁-C₈alkyl, e.g. trifluoromethyl, unsubstituted or substituted C₅-C₁₀aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C₁-C₈alkoxy, e.g. methoxy, ethoxy or isopropoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclylC₁-C₈alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetylamino, halogen, e.g. fluoro or chloro, unsubstituted or substituted carbamoyl, e.g. cyclohexylcarbamoyl, piperidinocarbonyl, piperazinocarbonyl, N-methylpiperazinocarbonyl or morpholinocarbonyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl; preferably hydrogen, methyl, isopropyl, trifluoromethyl, phenyl, o-, m- or p-methoxyphenyl, piperidino, piperazino, N-methylpiperazino, morpholino, methoxy, ethoxy, isopropoxy, phenoxy, 3-morpholinopropoxy, 2-morpholinoethoxy, 2-(1-imidazolyl)ethoxy, dimethylamino, fluoro, morpholinocarbonyl, piperidinocarbonyl, piperazinocarbonyl or cyclohexylcarbamoyl;
- (i') R⁸ is hydrogen, C₁-C₈alkyl, e.g. methyl, ethyl or isopropyl, haloC₁-C₈alkyl, e.g. trifluoromethyl, C₅-C₁₀aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C₁-C₈alkoxy, e.g. methoxy, ethoxy or isopropoxy, haloC₁-C₈alkoxy, e.g. trifluoromethoxy, C₅-C₁₀aryloxy, e.g. phenoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclylC₁-C₈alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino or dimethylamino, halogen, e.g. fluoro or chloro, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl, or nitro; preferably hydrogen, methyl, piperidino, piperazino, N-methylpiperazino, morpholino, methoxy, ethoxy, trifluoromethoxy, phenoxy, 1-methyl-4-piperidyloxy, 3-morpholinopropoxy, 2-morpholinoethoxy, 3-(N-methylpiperazino)-propoxy, methylamino, fluoro, chloro, sulfamoyl or nitro;
- (j') R¹⁰ is C₁-C₈alkyl, e.g. methyl, ethyl or butyl, haloC₁-C₈alkyl, e.g. trifluoromethyl, C₁-C₈alkoxy, e.g. methoxy or ethoxy, unsubstituted or substituted heterocyclylC₁-C₈alkoxy, e.g. 2-(1-

imidazolyl)ethoxy, unsubstituted or substituted amino, e.g. methylamino or dimethylamino, halogen, e.g. fluoro or chloro; preferably methyl, butyl, methoxy, ethoxy, 2-(1-imidazolyl)ethoxy, methylamino, dimethylamino or fluoro; and

(k') each pair of adjacent substituents R^7 and R^8 , or R^8 and R^9 or R^9 and R^{10} , are $-NH-CH=CH-$, $-CH=CH-NH-$, $-NH-N=CH-$, $-CH=N-NH-$, $-CH_2-CH_2-CH_2-$, $-CH_2-CH_2-CH_2-CH_2-$, $-O-CH_2-O-$, or $-O-CF_2-O-$; preferably the pair of adjacent substituents R^7 and R^8 or R^8 and R^9 being $-O-CH_2-O-$ or the pair of adjacent substituents R^9 and R^{10} being $-NH-CH=CH-$, $-CH=N-NH-$, $-CH_2-CH_2-CH_2-$, $-CH_2-CH_2-CH_2-CH_2-$ or $-O-CF_2-O-$.

Most preferred as compounds of the formula I are those wherein the substituents have the meaning given in the Examples.

In another embodiment of the invention the invention provides a compound of formula I' with the proviso that this does not include any of the compounds of examples 1 to 52 inclusive.



in which:

n' is selected from 1, 2 and 3;

R'_1 is selected from C_{6-10} aryl, C_{5-10} heteroaryl, C_{3-12} cycloalkyl and C_{3-10} heterocycloalkyl;

wherein any aryl, heteroaryl, cycloalkyl or heterocycloalkyl of R'_1 is optionally substituted by 1 to 3 radicals independently selected from C_{1-6} alkyl, C_{1-6} alkoxy, alkoxy-substituted- C_{1-6} alkyl, halo-substituted- C_{1-6} alkyl, halo-substituted- C_{1-6} alkoxy, $-C(O)NR'_5R'_6$, $-S(O)_{0-2}NR'_5R'_6$, $-S(O)_{0-2}R'_5$, $-C(O)R'_4$, $-OXR'_4$, $-NR'_5XNR'_5R'_6$, $-OXNR'_5R'_6$, $-OXOR'_5$ and $-XR'_4$;

wherein X' is a bond or C_{1-6} alkylene; R'_5 is selected from hydrogen and C_{1-6} alkyl; R'_6 is selected from hydrogen, C_{1-6} alkyl and C_{3-12} cycloalkyl- C_{1-4} alkyl; and R'_4 is independently selected from C_{6-10} aryl, C_{5-10} heteroaryl, C_{3-12} cycloalkyl and C_{3-10} heterocycloalkyl;

and any aryl, heteroaryl, cycloalkyl or heterocycloalkyl of R'_4 is optionally substituted by 1 to 3 radicals independently selected from C_{1-6} alkyl, C_{3-10} heterocycloalkyl- C_{0-4} alkyl optionally substituted with C_{1-6} alkyl, $-C(O)NR'_5R'_6$, $-XNR'_5R'_6$, $-NR'_5XNR'_5R'_6$ and $-NR'_5C(O)R'_6$; wherein X is a bond or C_{1-6} alkylene; R'_5 and R'_6 are independently selected from hydrogen and C_{1-6} alkyl;

R'_2 is selected from hydrogen and halo, cyano, C_{1-6} alkyl, halo-substituted- C_{1-6} alkyl;

R'_3 is selected from halo, $-S(O)_{0-2}NR'_5R'_6$, $-S(O)_{0-2}R'_6$, $-NR'_5S(O)_{0-2}R'_6$, $-C(O)NR'_5R'_6$, $-C(O)R'_6$ and $-C(O)OR'_6$; wherein R'_5 is selected from hydrogen and C_{1-6} alkyl; and R'_6 is selected from hydrogen, C_{1-6} alkyl and C_{3-12} cycloalkyl;

and the pharmaceutically acceptable salts, hydrates, solvates, isomers and prodrugs thereof.

Preferably a compound of formula I' in which:

n' is selected from 1 and 2;

R'_1 is selected from C_{6-10} aryl and C_{5-10} heteroaryl; wherein any aryl or heteroaryl of R'_1 is optionally substituted by 1 to 3 radicals independently selected from C_{1-6} alkyl, C_{1-6} alkoxy, $-C(O)NR'_5R'_6$, $-OX'R'_4$, $-C(O)R'_4$, $-NR'_5XNR'_5R'_6$, $-OX'NR'_5R'_6$, $-OX'OR'_5$ and $-X'R'_4$; wherein X' is a bond or C_{1-6} alkylene; R'_5 is selected from hydrogen and C_{1-6} alkyl; R'_6 is selected from hydrogen, C_{1-6} alkyl and C_{3-12} cycloalkyl- C_{1-4} alkyl; and R'_4 is C_{3-10} heterocycloalkyl optionally substituted by 1 to 3 radicals independently selected from C_{1-6} alkyl, halo-substituted- C_{1-6} alkyl, C_{3-10} heterocycloalkyl- C_{0-4} alkyl optionally substituted with C_{1-6} alkyl, $-C(O)NR'_5R'_6$, $-X'NR'_5R'_6$, $-NR'_5XNR'_5R'_6$ and $-NR'_5C(O)R'_6$; wherein X' is a bond or C_{1-6} alkylene; R'_5 and R'_6 are independently selected from hydrogen and C_{1-6} alkyl;

R'_2 is selected from hydrogen and halo;

R'_3 is selected from halo, $-S(O)_{0-2}NR'_5R'_6$, $-S(O)_{0-2}R'_6$, $-NR'_5S(O)_{0-2}R'_6$, $-C(O)NR'_5R'_6$ and $-C(O)OR'_6$; wherein R'_5 is selected from hydrogen and C_{1-6} alkyl; and R'_6 is selected from hydrogen, C_{1-6} alkyl and C_{3-12} cycloalkyl.


more preferably a compound of formula I' in which R'_1 is selected from phenyl, pyridinyl, pyrazolyl and pyrimidinyl; wherein any aryl or heteroaryl of R'_1 is optionally substituted by 1 to 3 radicals independently selected from ethoxy, ethyl, propyl, methyl, t-butyl, trifluoromethyl, nitrile,

cyclobutyloxy, 2,2,2-trifluoroethoxy, methoxy, isobutyloxy, t-butyloxy, isopropoxy, methyl-amino-carbonyl, cyclopropyl-methoxy, dimethylamino-propyl-amino, methoxy-ethoxy, $-X'R'_4$, $-C(O)R'_4$ and $-OX'R'_4$; wherein X' is a bond, methylene or ethylene; R'_4 is selected from piperazinyl, piperidinyl, pyrrolidinyl, morpholino, azepanyl and 1,4-dioxo-8-aza-spiro[4.5]dec-8-yl; wherein R'_4 is optionally substituted by 1 to 3 radicals independently selected from methyl, isopropyl, acetyl, acetyl-methyl-amino, 3-dimethylamino-2,2-dimethyl-propylamino, ethyl-methyl-amino-ethoxy, diethyl-amino-ethoxy, amino-carbonyl, ethyl, 2-oxo-pyrrolidin-1-yl, pyrrolidinyl, pyrrolidinyl-methyl, piperidinyl optionally substituted with methyl or ethyl, morpholino, dimethylamino, dimethylamino-propyl-amino, methyl-amino and ethyl-amino.

Even more preferably a compound of formula I' in which R'_2 is selected from hydrogen and halo; and R'_3 is selected from halo, dimethyl-sulfamoyl, isobutyl-sulfamoyl, methyl-sulfamoyl, ethyl-sulfamoyl, propyl-sulfonyl, ethyl-amino-carbonyl, 1-ethyl-propyl-sulfamoyl, cyclopentyl-sulfamoyl, isopropyl-sulfamoyl, cyclohexyl-sulfonyl, cyclopropyl-methyl-sulfamoyl, cyclobutyl-sulfamoyl, isopropyl-sulfonyl,

Most preferably a compound of example 53


In a yet further embodiment of the invention the present invention also provides a process for the production of a compound of formula I, comprising reacting a compound of formula II



(II)

wherein R^0 , R^1 , R^2 , R^3 , R^4 , R^5 , and R^6 are as defined above, and Y is a leaving group, preferably halogen such as bromide, iodine, or in particular chloride;

with a compound of formula III



(III)

wherein R⁷, R⁸, R⁹ and R¹⁰ are as defined above;

and, if desired, converting a compound of formula I, wherein the substituents have the meaning as defined above, into another compound of formula I as defined;

and recovering the resulting compound of formula I in free form or as a salt, and, when required, converting the compound of formula I obtained in free form into the desired salt, or an obtained salt into the free form.

The reaction can be carried out in a manner known per se, the reaction conditions being dependent especially on the reactivity of the leaving group Y and the reactivity of the amino group in the aniline of formula III, usually in the presence of a suitable solvent or diluent or of a mixture thereof and, if necessary, in the presence of an acid or a base, with cooling or, preferably, with heating, for example in a temperature range from approximately -30°C to approximately +150°C, especially approximately from 0°C to +100°C, preferably from room temperature (approx. +20 °C) to +80 °C, in an open or closed reaction vessel and/or in the atmosphere of an inert gas, for example nitrogen. Alternatively, the reaction can proceed in the presence of a suitable catalyst (for example, palladium di-benzyl-acetone), in the presence of a base (for example, caesium carbonate) and in the presence of a suitable reaction facilitator (for example, xanthphos).

If one or more other functional groups, for example carboxy, hydroxy or amino, are or need to be protected in a compound of formula II or III, because they should not take part in the reaction, these are such groups as are usually used in the synthesis of peptide compounds, cephalosporins and penicillins, as well as nucleic acid derivatives and sugars.

The protecting groups may already be present in precursors and should protect the functional groups concerned against unwanted secondary reactions, such as substitution reaction or

solvolysis. It is a characteristic of protecting groups that they lend themselves readily, i.e. without undesired secondary reactions, to removal, typically by solvolysis, reduction, photolysis or also by enzyme activity, for example under conditions analogous to physiological conditions, and that they are not present in the end-products. The specialist knows, or can easily establish, which protecting groups are suitable with the reactions mentioned hereinabove.

Salts of a compound of formula I with a salt-forming group may be prepared in a manner known per se. Acid addition salts of compounds of formula I may thus be obtained by treatment with an acid or with a suitable anion exchange reagent.


Salts can usually be converted to compounds in free form, e.g. by treating with suitable basic agents, for example with alkali metal carbonates, alkali metal hydrogencarbonates, or alkali metal hydroxides, typically potassium carbonate or sodium hydroxide.

Stereoisomeric mixtures, e.g. mixtures of diastereomers, can be separated into their corresponding isomers in a manner known per se by means of suitable separation methods. Diastereomeric mixtures for example may be separated into their individual diastereomers by means of fractionated crystallization, chromatography, solvent distribution, and similar procedures. This separation may take place either at the level of a starting compound or in a compound of formula I itself. Enantiomers may be separated through the formation of diastereomeric salts, for example by salt formation with an enantiomer-pure chiral acid, or by means of chromatography, for example by HPLC, using chromatographic substrates with chiral ligands.

It should be emphasized that reactions analogous to the conversions mentioned in this chapter may also take place at the level of appropriate intermediates.


The compounds of formula I, including their salts, are also obtainable in the form of hydrates, or their crystals can include for example the solvent used for crystallization (present as solvates).

The compound of formula II used as starting materials may be obtained by reacting a compound of formula IV



(IV)

with a compound of formula V



(V)

wherein R¹, R², R³, R⁴, R⁵ and R⁶ are as defined above, and Y¹ and Y² are identical or different leaving groups as defined above for Y. The reaction conditions are those mentioned above for the reaction of a compound of formula II with a compound of formula III.

The compounds of formula IV and V are known or may be produced in accordance with known procedures.

The compounds of formula I and their pharmaceutically acceptable salts exhibit valuable pharmacological properties when tested in vitro in cell-free kinase assays and in cellular assays, and are therefore useful as pharmaceuticals. In particular, the compounds of the invention are inhibitors of Focal Adhesion Kinase, and are useful as pharmaceuticals to treat conditions caused by a malfunction of signal cascades connected with Focal Adhesion Kinase, in particular tumors as described hereinbelow.

Focal Adhesion Kinase (FAK) is a key enzyme in the integrin-mediated outside-in signal cascade (D. Schlaepfer et al., Prog Biophys Mol Biol 1999, 71, 435-478). Interaction between cells and extracellular matrix (ECM) proteins is transduced as intracellular signals important for growth, survival and migration through cell surface receptors, integrins. FAK plays an essential role in these integrin-mediated outside-in signal cascades. The trigger in the signal transduction cascade is the autophosphorylation of Y397. Phosphorylated Y397 is a SH2 docking site for Src family tyrosine kinases. The bound c-Src kinase phosphorylates other tyrosine residues in FAK. Among them, phosphorylated Y925 becomes a binding site for the SH2 site of Grb2 small adaptor protein. This direct binding of Grb2 to FAK is one of the key steps for the activation of down stream targets such as the Ras-ERK2/MAP kinase cascade.

The inhibition of endogenous FAK signalling results in reduced motility and in some cases induces cell death. On the other hand, enhancing FAK signalling by exogenous expression increases cell motility and transmitting a cell survival signal from ECM. In addition FAK is overexpressed in invasive and metastatic epithelial, mesenchymal, thyroid and prostate cancers. Consequently, an inhibitor of FAK is likely to be a drug for anti-tumor growth and metastasis. The compounds of the invention are thus indicated, for example, to prevent and/or treat a vertebrate and more particularly a mammal, affected by a neoplastic disease, in particular breast tumor, cancer of the bowel (colon and rectum), stomach cancer and cancer of the ovary and prostate, non-small cell lung cancer, small cell lung cancer, cancer of liver, melanoma, bladder tumor and cancer of head and neck.

The relation between FAK inhibition and immuno-system is described e.g. in G.A. van Seventer et al., Eur. J. Immunol. 2001, 31, 1417-1427. Therefore, the compounds of the invention are, for example, useful to prevent and/or treat a vertebrate and more particularly a mammal, affected by immune system disorders, diseases or disorders mediated by T lymphocytes, B lymphocytes, mast cells and/or eosinophils e.g. acute or chronic rejection of organ or tissue allo- or xenografts, atherosclerosis, vascular occlusion due to vascular injury such as angioplasty, restenosis, hypertension, heart failure, chronic obstructive pulmonary disease, CNS disease such as Alzheimer disease or amyotrophic lateral sclerosis, cancer, infectious disease such as AIDS, septic shock or adult respiratory distress syndrome, ischemia/reperfusion injury e.g. myocardial infarction, stroke, gut ischemia, renal failure or hemorrhage shock, or traumatic shock. The agent of the invention are also useful in the treatment and/or prevention of acute or chronic inflammatory diseases or disorders or autoimmune diseases e.g. rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, diabetes (type I and II) and the disorders associated with therewith, respiratory diseases such as asthma or inflammatory liver injury, inflammatory glomerular injury, cutaneous manifestations of immunologically-mediated disorders or illnesses, inflammatory and hyperproliferative skin diseases (such as psoriasis, atopic dermatitis, allergic contact dermatitis, irritant contact dermatitis and further eczematous dermatitises, seborrhoeic dermatitis), inflammatory eye diseases, e.g. Sjogren's syndrome, keratoconjunctivitis or uveitis, inflammatory bowel disease, Crohn's disease or ulcerative colitis.

Compounds of the invention are active in a FAK assay system as described in the Examples, and show an inhibition IC_{50} in the range of 1 nM to 100 nM. Particularly active are the compounds Example No. 3-12 and No. 3-17 described hereinbelow showing IC_{50} values in the range of 1 to 5 nM.

Some of the compounds of the invention exhibit also ZAP-70 (zeta chain-associated protein of 70 kD) protein tyrosine kinase inhibiting activity. ZAP-70 protein tyrosine kinase interaction of the agents of the invention may be demonstrated by their ability to prevent phosphorylation of e.g. LAT-11 (linker for activation of T cell) by human ZAP-70 protein tyrosine kinase in aqueous solution, as described in the Examples. The compounds of the invention are thus also indicated for the prevention or treatment of disorders or diseases where ZAP-70 inhibition inhibition play a role.

Compounds of the invention are active in a ZAP-70 assay system as described in the Examples, and show an inhibition IC_{50} in the range of 1 μ M to 10 μ M, e.g. the compounds Example No. 2 and No. 3-2 described hereinbelow.

Compounds of the present invention are also good inhibitors of the IGF-IR (insulin like growth factor receptor 1) and are therefore useful in the treatment of IGF-1R mediated diseases for example such diseases include proliferative diseases, such as tumours, like for example breast, renal, prostate, colorectal, thyroid, ovarian, pancreas, neuronal, lung, uterine and gastro-intestinal tumours as well as osteosarcomas and melanomas. The efficacy of the compounds of the invention as inhibitors of IGF-IR tyrosine kinase activity can be demonstrated using a cellular "Capture ELISA". In this assay the activity of the compounds of the invention against Insulin-like growth factor I (IGF-I) induced autophosphorylation of the IGF-IR is determined.

The compounds of formula I and their pharmaceutically acceptable salts exhibit valuable pharmacological properties when tested in vitro in cell-free kinase assays and in cellular assays, and are therefore useful as pharmaceuticals. In particular, the compounds of the invention are inhibitors of Anaplastic Lymphoma Kinase (ALK), and are useful as pharmaceuticals to treat conditions caused by a malfunction of signal cascades connected with Anaplastic Lymphoma Kinase, in particular tumors as described hereinbelow.

ALK-mediated signaling could play a role in the development and/or progression of a number of common solid tumors (Pulford, K., et al., J. Cell. Physiol. 2004 Jun;199(3):330-58). The

compounds of the present invention also exhibit powerful inhibition of the tyrosine kinase activity of anaplastic lymphoma kinase (ALK) and its fusion proteins, particularly the fusion protein of NPM-ALK. This protein tyrosine kinase results from a gene fusion of nucleophosmin (NPM) and the anaplastic lymphoma kinase (ALK), rendering the protein tyrosine kinase activity of ALK ligand-independent. NPM-ALK plays a key role in signal transmission in a number of hematopoietic and other human cells leading to hematological and neoplastic diseases, for example in anaplastic large-cell lymphoma (ALCL) and non-Hodgkin's lymphomas (NHL), specifically in ALK⁺ NHL or Alkomas, in inflammatory myofibroblastic tumors (IMT) and neuroblastomas. (Duyster J et al. 2001 Oncogene 20, 5623-5637). NPM-ALK has been shown to be a potent oncogene in vitro, being able to transform various cell lines and primary hematopoietic cells. Furthermore, NPM-ALK transduced bone marrow cells are able to induce a lymphoma-like disease after transplantation into irradiated recipient mice. Signaling pathways activated by NPM-ALK include ras, PLC and PI3K pathways and, in addition, STAT5 has been shown to be phosphorylated by NPM-ALK. In addition to NPM-ALK, other gene fusions have been identified in human hematological and neoplastic diseases; mainly TPM3-ALK (a fusion of nonmuscle tropomyosin 3 with ALK). Further, the ALK fusion protein CLTC-ALK, is associated with diseases that include classical T cell or null ALCL, ALK⁺ DLBCL and inflammatory myofibroblastic tumors. CLTC-ALK is also thought to play a role in the pathogenesis of large B-cell lymphomas.

Further, the ALK fusion protein CLTC-ALK is associated with diseases that include classical T cell or null ALCL, ALK⁺ DLBCL and inflammatory myofibroblastic tumors. CLTC-ALK is also thought to play a role in the pathogenesis of large B-cell lymphomas.

Aberrant activity of ALK is involved in the development of brain tumors and overexpression of ALK has been reported in neuroblastomas and several cell lines derived from neural tissue. ALK-mediated signaling could play a role in the development and/or progression of a number of common solid tumors (Pulford, K., et al., J. Cell. Physiol. 2004 Jun;199(3):330-58).

The inhibition of ALK tyrosine kinase activity can be demonstrated using known methods, for example using the recombinant kinase domain of the ALK in analogy to the VEGF-R kinase assay described in J. Wood et al. Cancer Res. 60, 2178-2189 (2000). In vitro enzyme assays using GST-ALK protein tyrosine kinase are performed in 96-well plates as

a filter binding assay in 20 mM Tris-HCl, pH = 7.5, 3 mM MgCl₂, 10 mM MnCl₂, 1 mM DTT, 0.1 µCi/assay (=30 µl) [γ -³³P]-ATP, 2 µM ATP, 3 µg/ml poly (Glu, Tyr 4:1) Poly-EY (Sigma P-0275), 1 % DMSO, 25 ng ALK enzyme. Assays are incubated for 10 min at ambient temperature. Reactions are terminated by adding 50 µl of 125 mM EDTA, and the reaction mixture is transferred onto a MAIP Multiscreen plate (Millipore, Bedford, MA, USA), previously wet with methanol, and rehydrated for 5 min with H₂O. Following washing (0.5 % H₃PO₄), plates are counted in a liquid scintillation counter. IC₅₀ values are calculated by linear regression analysis of the percentage inhibition. Compared with the control without inhibitor, the compounds of formula I inhibit the enzyme activity by 50 % (IC₅₀), for example in a concentration of from 0.001 to 0.5 µM, especially from 0.01 to 0.1 µM.

The compounds of formula I potentially inhibit the growth of human NPM-ALK overexpressing murine BaF3 cells (DSMZ Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany). The expression of NPM-ALK is achieved by transfecting the BaF3 cell line with an expression vector pCIneo™ (Promega Corp., Madison WI, USA) coding for NPM-ALK and subsequent selection of G418 resistant cells. Non-transfected BaF3 cells depend on IL-3 for cell survival. In contrast NPM-ALK expressing BaF3 cells (named BaF3-NPM-ALK hereinafter) can proliferate in the absence of IL-3 because they obtain proliferative signal through NPM-ALK kinase. Putative inhibitors of the NPM-ALK kinase therefore abolish the growth signal and result in antiproliferative activity. The antiproliferative activity of putative inhibitors of the NPM-ALK kinase can however be overcome by addition of IL-3 which provides growth signals through an NPM-ALK independent mechanism. [For an analogous cell system using FLT3 kinase see E Weisberg et al. Cancer Cell; 1, 433-443 (2002)]. The inhibitory activity of the compounds of formula I is determined, briefly, as follows: BaF3-NPM-ALK cells (15,000/microtitre plate well) are transferred to 96-well microtitre plates. The test compounds [dissolved in dimethyl sulfoxide (DMSO)] are added in a series of concentrations (dilution series) in such a manner that the final concentration of DMSO is not greater than 1 % (v/v). After the addition, the plates are incubated for two days during which the control cultures without test compound are able to undergo two cell-division cycles. The growth of the BaF3-NPM-ALK cells is measured by means of Yopro™ staining [T Idziorek et al. J. Immunol. Methods; 185: 249-258 (1995)]: 25 µl of lysis buffer consisting of 20 mM sodium citrate, pH 4.0, 26.8 mM sodium chloride, 0.4 % NP40, 20 mM EDTA and 20 mM is added to each well. Cell lysis is completed within 60 min at room temperature and total amount of Yopro bound to DNA is determined by

measurement using the Cytofluor II 96-well reader (PerSeptive Biosystems) with the following settings: Excitation (nm) 485/20 and Emission (nm) 530/25.

IC₅₀ values are determined by a computer-aided system using the formula:

$$IC_{50} = [(ABS_{test} - ABS_{start}) / (ABS_{control} - ABS_{start})] \times 100. \text{ (ABS = absorption)}$$

The IC₅₀ value in those experiments is given as that concentration of the test compound in question that results in a cell count that is 50 % lower than that obtained using the control without inhibitor. The compounds of formula I exhibit inhibitory activity with an IC₅₀ in the range from approximately 0.01 to 1 µM.

The antiproliferative action of the compounds of formula I can also be determined in the human KARPAS-299 lymphoma cell line (DSMZ Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany) [described in WG Dirks et al. Int. J. Cancer 100, 49-56 (2002)] using the same methodology described above for the BaF3-NPM-ALK cell line. The compounds of formula I exhibit inhibitory activity with an IC₅₀ in the range from approximately 0.01 to 1 µM.

The action of the compounds of formula I on autophosphorylation of the ALK can be determined in the human KARPAS-299 lymphoma cell line by means of an immunoblot as described in WG Dirks et al. Int. J. Cancer 100, 49-56 (2002). In that test the compounds of formula I exhibit an IC₅₀ of approximately from 0.001 to 1 µM.

Among the compounds of formula I, 2-[5-chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzamide is an especially potent ALK inhibitor, in that this compound inhibits the growth of the BaF3-NPM-ALK cells with an IC₅₀ of 97 nM. Further specifically preferred compounds that inhibit the tyrosine kinase activity of anaplastic lymphoma kinase (ALK) are the compounds described hereinafter in the examples 7A and 7B, as well as 7-2, 7-15, 19-5, 21-1, 26-3 and 28-5, respectively, all of which are having an IC₅₀ within the range from <0.5 to 200 nM.

For the above uses in the treatment of neoplastic diseases and immune system disorders the required dosage will of course vary depending on the mode of administration, the particular condition to be treated and the effect desired. In general, satisfactory results are indicated to be

obtained systemically at daily dosages of from about 0.1 to about 100 mg/kg body weight. An indicated daily dosage in the larger mammal, e.g. humans, is in the range from about 0.5 mg to about 2000 mg, conveniently administered, for example, in divided doses up to four times a day or in retard form.

The compounds of the invention may be administered by any conventional route, in particular parenterally, for example in the form of injectable solutions or suspensions, enterally, preferably orally, for example in the form of tablets or capsules, topically, e.g. in the form of lotions, gels, ointments or creams, or in a nasal or a suppository form. Pharmaceutical compositions comprising a compound of the invention in association with at least one pharmaceutical acceptable carrier or diluent may be manufactured in conventional manner by mixing with a pharmaceutically acceptable carrier or diluent. Unit dosage forms for oral administration contain, for example, from about 0.1 mg to about 500 mg of active substance. Topical administration is e.g. to the skin. A further form of topical administration is to the eye.

The pharmaceutical compositions of the present invention are prepared in a manner known per se, for example by means of conventional mixing, granulating, coating, dissolving or lyophilizing processes.

Preference is given to the use of solutions of the active ingredient, and also suspensions or dispersions, especially isotonic aqueous solutions, dispersions or suspensions which, for example in the case of lyophilized compositions comprising the active ingredient alone or together with a carrier, for example mannitol, can be made up before use. The pharmaceutical compositions may be sterilized and/or may comprise excipients, for example preservatives, stabilizers, wetting agents and/or emulsifiers, solubilizers, salts for regulating osmotic pressure and/or buffers and are prepared in a manner known per se, for example by means of conventional dissolving and lyophilizing processes. The said solutions or suspensions may comprise viscosity-increasing agents, typically sodium carboxymethylcellulose, carboxymethylcellulose, dextran, polyvinylpyrrolidone, or gelatins, or also solubilizers, e.g. Tween 80® (polyoxyethylene(20)sorbitan mono-oleate).

Suspensions in oil comprise as the oil component the vegetable, synthetic, or semi-synthetic oils customary for injection purposes. In respect of such, special mention may be made of liquid fatty acid esters that contain as the acid component a long-chained fatty acid having from 8 to

22, especially from 12 to 22, carbon atoms, for example lauric acid, tridecylic acid, myristic acid, pentadecylic acid, palmitic acid, margaric acid, stearic acid, arachidic acid, behenic acid or corresponding unsaturated acids, for example oleic acid, elaidic acid, erucic acid, brassidic acid or linoleic acid, if desired with the addition of antioxidants, for example vitamin E, β -carotene or 3,5-di-tert-butyl-4-hydroxytoluene. The alcohol component of these fatty acid esters has a maximum of 6 carbon atoms and is a monovalent or polyvalent, for example a mono-, di- or trivalent, alcohol, for example methanol, ethanol, propanol, butanol or pentanol or the isomers thereof, but especially glycol and glycerol. As fatty acid esters, therefore, the following are mentioned: ethyl oleate, isopropyl myristate, isopropyl palmitate, "Labrafil M 2375" (polyoxyethylene glycerol), "Labrafil M 1944 CS" (unsaturated polyglycolized glycerides prepared by alcoholysis of apricot kernel oil and consisting of glycerides and polyethylene glycol ester), "Labrasol" (saturated polyglycolized glycerides prepared by alcoholysis of TCM and consisting of glycerides and polyethylene glycol ester; all available from Gattefossé, France), and/or "Miglyol 812" (triglyceride of saturated fatty acids of chain length C_8 to C_{12} from Hüls AG, Germany), but especially vegetable oils such as cottonseed oil, almond oil, olive oil, castor oil, sesame oil, soybean oil and more especially groundnut oil.

The manufacture of injectable preparations is usually carried out under sterile conditions, as is the filling, for example, into ampoules or vials, and the sealing of the containers.

Pharmaceutical compositions for oral administration can be obtained, for example, by combining the active ingredient with one or more solid carriers, if desired granulating a resulting mixture, and processing the mixture or granules, if desired or necessary, by the inclusion of additional excipients, to form tablets or tablet cores.

Suitable carriers are especially fillers, such as sugars, for example lactose, saccharose, mannitol or sorbitol, cellulose preparations, and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, and also binders, such as starches, for example corn, wheat, rice or potato starch, methylcellulose, hydroxypropyl methylcellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone, and/or, if desired, disintegrators, such as the above-mentioned starches, also carboxymethyl starch, crosslinked polyvinylpyrrolidone, alginic acid or a salt thereof, such as sodium alginate. Additional excipients are especially flow conditioners and lubricants, for example silicic acid, talc, stearic acid or salts thereof, such as magnesium or calcium stearate, and/or polyethylene glycol, or derivatives thereof.

Tablet cores can be provided with suitable, optionally enteric, coatings through the use of, inter alia, concentrated sugar solutions which may comprise gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol and/or titanium dioxide, or coating solutions in suitable organic solvents or solvent mixtures, or, for the preparation of enteric coatings, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate. Dyes or pigments may be added to the tablets or tablet coatings, for example for identification purposes or to indicate different doses of active ingredient.

Pharmaceutical compositions for oral administration also include hard capsules consisting of gelatin, and also soft, sealed capsules consisting of gelatin and a plasticizer, such as glycerol or sorbitol. The hard capsules may contain the active ingredient in the form of granules, for example in admixture with fillers, such as corn starch, binders, and/or glidants, such as talc or magnesium stearate, and optionally stabilizers. In soft capsules, the active ingredient is preferably dissolved or suspended in suitable liquid excipients, such as fatty oils, paraffin oil or liquid polyethylene glycols or fatty acid esters of ethylene or propylene glycol, to which stabilizers and detergents, for example of the polyoxyethylene sorbitan fatty acid ester type, may also be added.

Pharmaceutical compositions suitable for rectal administration are, for example, suppositories that consist of a combination of the active ingredient and a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, paraffin hydrocarbons, polyethylene glycols or higher alkanols.

For parenteral administration, aqueous solutions of an active ingredient in water-soluble form, for example of a water-soluble salt, or aqueous injection suspensions that contain viscosity-increasing substances, for example sodium carboxymethylcellulose, sorbitol and/or dextran, and, if desired, stabilizers, are especially suitable. The active ingredient, optionally together with excipients, can also be in the form of a lyophilizate and can be made into a solution before parenteral administration by the addition of suitable solvents.

Solutions such as are used, for example, for parenteral administration can also be employed as infusion solutions.

Preferred preservatives are, for example, antioxidants, such as ascorbic acid, or microbicides, such as sorbic acid or benzoic acid.

The compounds of the invention may be administered as the sole active ingredient or together with other drugs useful against neoplastic diseases or useful in immunomodulating regimens. For example, the agents of the invention may be used in accordance with the invention in combination with pharmaceutical compositions effective in various diseases as described above, e.g. with cyclophosphamide, 5-fluorouracil, fludarabine, gemcitabine, cisplatin, carboplatin, vincristine, vinblastine, etoposide, irinotecan, paclitaxel, docetaxel, rituxan, doxorubicine, gefitinib, or imatinib; or also with cyclosporins, rapamycins, ascomycins or their immunosuppressive analogs, e.g. cyclosporin A, cyclosporin G, FK-506, sirolimus or everolimus, corticosteroids, e.g. prednisone, cyclophosphamide, azathioprene, methotrexate, gold salts, sulfasalazine, antimalarials, brequinar, leflunomide, mizoribine, mycophenolic acid, mycophenolate, mofetil, 15-deoxyspergualine, immuno-suppressive monoclonal antibodies, e.g. monoclonal antibodies to leukocyte receptors, e.g. MHC, CD2, CD3, CD4, CD7, CD25, CD28, CD40, CD45, CD58, CD80, CD86, CD152, CD137, CD154, ICOS, LFA-1, VLA-4 or their ligands, or other immunomodulatory compounds, e.g. CTLA4Ig.

In accordance with the foregoing, the present invention also provides:

- (1) A compound of the invention for use as a pharmaceutical;
- (2) a compound of the invention for use as a 5-Chloro-N²*-{2-methoxy-4-[4-(4-methyl-piperazin-1-yl)-piperidin-1-yl]-phenyl}-N⁴*-[2-(propane-2-sulfonyl)-phenyl]-pyrimidine-2,4-diamine, for example for use in any of the particular indications hereinbefore set forth;
- (3) a pharmaceutical composition, e.g. for use in any of the indications herein before set forth, comprising a compound of the invention as active ingredient together with one or more pharmaceutically acceptable diluents or carriers;
- (4) a method for the treatment of any particular indication set forth hereinbefore in a subject in need thereof which comprises administering an effective amount of a compound of the invention or a pharmaceutical composition comprising same;
- (5) the use of a compound of the invention for the manufacture of a medicament for the treatment or prevention of a disease or condition in which FAK and/or ALK and/or ZAP-70 and/or IGF-I activation plays a role or is implicated, preferably ALK;

(6) the method as defined above under (4) comprising co-administration, e.g. concomitantly or in sequence, of a therapeutically effective amount of a compound of the invention and one or more further drug substances, said further drug substance being useful in any of the particular indications set forth hereinbefore;

(7) a combination comprising a therapeutically effective amount of a compound of the invention and one or more further drug substances, said further drug substance being useful in any of the particular indications set forth hereinbefore;

(8) use of a compound of the invention for the manufacture of a medicament for the treatment or prevention of a disease which responds to inhibition of the anaplastic lymphoma kinase;

(9) the use according to (8), wherein the disease to be treated is selected from lymphoma, anaplastic large-cell lymphoma, non-Hodgkin's lymphomas, inflammatory myofibroblastic tumors and neuroblastomas;

(10) the use according to (8) or (9), wherein the compound is 2-[5-chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzamide or 5-Chloro-N²-(2-methoxy-4-[4-(4-methyl-piperazin-1-yl)-piperidin-1-yl]-phenyl)-N⁴-[2-(propane-2-sulfonyl)-phenyl]-pyrimidine-2,4-diamine or a pharmaceutically acceptable salt thereof, or any of the compounds described hereinafter in the examples or a pharmaceutically acceptable salt of any one of these;

(11) a method for the treatment of a disease which responds to inhibition of the anaplastic lymphoma kinase, especially a disease selected from anaplastic large-cell lymphoma, non-Hodgkin's lymphomas, inflammatory myofibroblastic tumors and neuroblastomas, comprising administering an effective amount of a compound of the invention, especially 2-[5-chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzamide or 5-Chloro-N²-(2-methoxy-4-[4-(4-methyl-piperazin-1-yl)-piperidin-1-yl]-phenyl)-N⁴-[2-(propane-2-sulfonyl)-phenyl]-pyrimidine-2,4-diamine or a pharmaceutically acceptable salt thereof.

Additionally preferred a compound according to the present invention that is useful as herein before described is a compound specifically mentioned in the examples.

Additional specifically preferred compounds according to the present invention that are useful either as FAK inhibitor, as ALK inhibitor or for inhibition of both and which may be prepared essentially according to the methods described hereinbefore are the following:

2-[5-chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzamide,

N²-(4-[1,4']Bipiperidinyl-1'-yl-2-methoxy-phenyl)-5-chloro-N⁴-[2-(propane-1-sulfonyl)-phenyl]-pyrimidine-2,4-diamine,
2-{5-Chloro-2-[2-methoxy-4-(4-methyl-piperazin-1-yl)-phenylamino]-pyrimidin-4-ylamino}-N-isopropyl-benzenesulfonamide,
2-[5-Bromo-2-(2-methoxy-5-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzenesulfonamide
2-[2-[5-(1-Acetyl-piperidin-4-yloxy)-2-methoxy-phenylamino]-5-bromo-pyrimidin-4-ylamino]-N-methyl-benzenesulfonamide,
N-[5-Bromo-2-(2,5-dimethoxy-phenylamino)-pyrimidin-4-yl]-N-(4-morpholin-4-yl-phenyl)-methanesulfonamide,
5-Bromo-N-4-(4-fluoro-phenyl)-N²*(2-methoxy-4-morpholin-4-yl-phenyl)-pyrimidine-2,4-diamine,
2-[5-Chloro-2-(2-methoxy-4-piperazin-1-yl-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzenesulfonamide,
2-[5-Bromo-2-(5-fluoro-2-methoxy-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzenesulfonamide,
2-[5-Chloro-2-(5-fluoro-2-methoxy-phenylamino)-pyrimidin-4-ylamino]-N-isobutyl-benzenesulfonamide, and
2-{5-Chloro-2-[2-methoxy-5-(4-methyl-piperazin-1-ylmethyl)-phenylamino]-pyrimidin-4-ylamino}-N-methyl-benzenesulfonamide,
5-Chloro-N²*-{2-methoxy-4-[4-(4-methyl-piperazin-1-yl)-piperidin-1-yl]-phenyl}-N⁴*-[2-(propane-2-sulfonyl)-phenyl]-pyrimidine-2,4-diamine.

The invention also provides a compound of formula 2-{5-Chloro-2-[4-(3-methylamino-pyrrolidin-1-yl)-phenylamino]-pyrimidin-4-ylamino}-N-isopropyl-benzenesulfonamide


The invention also provides a compound of formula 5-Chloro-N²*-{2-methoxy-4-[4-(4-methyl-piperazin-1-yl)-piperidin-1-yl]-phenyl}-N⁴*-[2-(propane-2-sulfonyl)-phenyl]-pyrimidine-2,4-diamine

The following Examples serve to illustrate the invention without limiting the invention in its scope.

Examples**Abbreviations**

AcOH = acetic acid, ALK = anaplastic lymphoma kinase, ATP = adenosine 5'-triphosphate, brine = saturated sodium chloride solution, BSA = bovine serum albumin, DIAD = diisopropyl azodicarboxylate, DIPCDI = N,N'-diisopropylcarbodiimide, DMAP = 4-dimethylaminopyridine, DMF = N,N-dimethylformamide, DTT = 1,4-dithio-D,L-threitol, EDTA = ethylene diamine tetraacetic acid, Et = ethyl, EtOAc = ethyl acetate, EtOH = ethanol, Eu-PT66 = LANCE™ europium-W1024-labelled anti-phosphotyrosine antibody (Perkin Elmer), FAK = Focal Adhesion Kinase, FRET = fluorescence resonance energy transfer, HEPES = N-2-hydroxyethyl-piperazine-N'-2-ethanesulfonic acid, HOAt = 1-hydroxy-7-azabenzotriazole, Me = methyl, RT-PCR = reverse transcription polymerase chain reaction, SA-(SL)APC = Streptavidin conjugated to SuperLight™ allophycocyanin (Perkin Elmer), subst. = substituted, TBTU = O-(benzotriazol-1-yl)-N,N,N',N'-tetramethylammonium tetrafluoroborate, THF = tetrahydrofuran.

Example 1: 2-[2-(2,5-Dimethoxy-phenylamino)-5-nitro-pyrimidin-4-ylamino]-N-methyl-benzenesulfonamide



To a solution of 2-(2-chloro-5-nitro-pyrimidin-4-ylamino)-N-methyl-benzenesulfonamide (100 mg, 0.29 mmol) in EtOH (3 mL), 2,5-dimethoxyaniline (49 mg, 0.32 mmol) is added at room temperature. The mixture is heated at 78°C for 5 h. The solvent is evaporated, and the mixture is purified by reverse phase HPLC to give the title product in.


R_f = 0.47 (n-hexane : ethyl acetate = 1:1). ¹H-NMR (400 MHz, CDCl₃), δ (ppm): 2.36 (d, 3H), 3.57 (s, 3H), 3.73 (s, 3H), 6.72 (d, 1H), 6.99 (d, 1H), 7.17 (s, 1H), 7.35 (t, 1H), 7.4-7.6 (m, 1H), 7.63 (d, 1H), 7.81 (d, 1H), 8.0-8.2 (m, 1H), 9.13 (s, 1H), 9.41 (br.s, 1H), 11.0 (s, 1H).

Preparation of 2-(2-chloro-5-nitro-pyrimidin-4-ylamino)-N-methyl-benzenesulfonamide:

2,4-Dichloro-5-nitro-pyrimidine (1.94 g, 10 mmol) and 2-amino-N-methyl-benzenesulfonamide (1.86 g, 10 mmol) are dissolved in CHCl₃ (30 mL). The reaction mixture is heated at 61°C for 2 h. The solvent is evaporated and the residue is washed with ether to give the title product.

R_f = 0.5 (n-hexane : ethyl acetate = 1:1). ¹H-NMR (400MHz, CDCl₃), δ (ppm): 2.67 (d, 3H), 4.6-4.7 (m, 2H), 7.41 (dd, 1H), 7.7 (dd, 1H), 8.04 (d, 1H), 8.15 (d, 1H), 9.21 (s, 1H), 11.2 (s, 1H).

Example 2: 2-[5-Bromo-2-(2,4-dimethoxy-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzenesulfonamide



To a solution of 2-(5-bromo-2-chloro-pyrimidin-4-ylamino)-N-methyl-benzenesulfonamide (300 mg, 0.79 mmol), 2,4-dimethoxyaniline (181.5 mg, 1.18 mmol) in ethanol (3 mL), 1 N hydrochloric acid (0.03 mL) is added and stirred under reflux condition for 5 hours. The reaction mixture is cooled to room temperature, poured into water and extracted twice with ethyl acetate. The organic layer is successively washed with water and brine, dried over magnesium sulfate, and evaporated in vacuo. The residue is purified with silica gel column chromatography (n-hexane : ethyl acetate = 5:1 to 1:1) to afford the title compound.

¹H-NMR (CDCl₃), δ (ppm): 8.95 (s, 1H), 8.44 (d, 1H), 8.20 (s, 1H), 7.98 (dd, 1H), 7.58 (ddd, 1H), 7.22-7.32 (m, 1H), 6.51 (d, 1H), 6.40 (d, 1H), 4.56-4.48 (m, 1H), 3.86 (s, 3H), 3.81 (s, 3H), 2.64 (d, 3H). R_f (n-hexane : ethyl acetate = 1:1): 0.31.


Preparation of 2-(5-bromo-2-chloro-pyrimidin-4-ylamino)-N-methyl-benzenesulfonamide

A solution of 5-bromo-2,4-dichloropyrimidine (684 mg, 3.0 mmol) and 2-amino-N-methyl-benzenesulfonamide (559 mg, 3.0 mmol) in N,N-dimethylformamide (10 mL) containing potassium carbonate (830 mg, 6.0 mmol) is stirred at room temperature for 23 hours. Saturated aqueous ammonium chloride is added and the mixture is poured into water and extracted twice with ethyl acetate. The organic layer is washed with brine, dried over sodium sulfate, and evaporated in vacuo. The residue is purified with silica gel column chromatography (n-hexane - ethyl acetate gradient) to afford the title compound as a slightly yellow solid.








¹H-NMR (CDCl₃), δ (ppm): 2.67 (d, 3H), 4.79 (q, 1H), 7.26 (s, 1H), 7.29 (ddd, 1H), 7.66 (ddd, 1H), 7.95 (dd, 1H), 8.37 (s, 1H), 8.48 (d, 1H), 9.52 (s, 1H). R_f (n-hexane : ethyl acetate = 10:3): 0.33.








Example 3:







The following 2-[5-bromo-2-(subst. phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzene-sulfonamides are prepared from 2-(5-bromo-2-chloro-pyrimidin-4-ylamino)-N-methyl-benzenesulfonamide and the corresponding aniline following the procedure of Example 2:

















ExptNo.	Rx	Rf (solvent) or MS	¹ H-NMR (400MHz), δ (ppm)
3-1		0.48 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 2.64(d, 3H), 4.48-4.40(m, 1H), 6.78(d, 1H), 6.87(bs, 1H), 6.99(dd, 1H), 6.82(s, 1H), 7.54(ddd, 1H), 7.79(d, 1H), 7.97(dd, 1H), 8.28(s, 1H), 8.32(dd, 1H), 9.07(s, 1H)
3-2		0.58 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 2.25(s, 3H), 2.33(s, 3H), 2.63(d, 3H), 4.53-4.45(m, 1H), 6.61(bs, 1H), 6.99(dd, 1H), 7.04(s, 1H), 7.18(ddd, 1H), 7.43(ddd, 1H), 7.56(d, 1H), 7.92(dd, 1H), 8.19(s, 1H), 8.41(dd, 1H), 9.08(s, 1H)
3-3		0.36 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 2.23(s, 3H), 2.62(d, 3H), 3.69(s, 3H), 4.53-4.44(m, 1H), 6.62(dd, 1H), 6.69(bs, 1H), 7.10(d, 1H), 7.19(dd, 1H), 7.48(d, 1H), 7.51(dd, 1H), 7.93(dd, 1H), 8.22(s, 1H), 8.44(dd, 1H), 9.09(s, 1H)
3-4		0.41 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 2.32(s, 3H), 2.63(d, 3H), 4.45-4.44(m, 1H), 6.85(d, 1H), 6.91(d, 1H), 7.00(bs, 1H), 7.28-7.24(m, 1H), 7.57(dd, 1H), 7.99(dd, 1H), 8.25(s, 1H), 8.39(d, 1H), 9.00(bs, 1H)






3-5		0.39 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 2.33(s, 3H), 2.63(d, 3H), 3.87(s, 3H), 4.46-4.44(m, 1H), 6.66(d, 1H), 6.71(s, 1H), 7.48(bs, 1H), 7.63-7.59(m, 1H), 7.97(dd, 1H), 8.05(d, 1H), 8.23 (s, 1H), 8.44(d, 1H), 8.92(bs, 1H)
3-6		0.27 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 2.63(d, 3H), 3.90(s, 3H), 4.45-4.40(m, 1H), 6.90-6.86(m, 2H), 7.00-6.96(m, 1H), 7.23-7.17 (m, 3H), 7.45(dd, 1H), 7.50-7.60(m, 2H), 7.97(dd, 1H), 8.22(d, 1H), 8.26 (s, 1H), 8.43(d, 1H), 8.94(bs, 1H)
3-7		0.34 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 2.30(s, 3H), 2.63(d, 3H), 4.44-4.43(m, 1H), 6.68 (bs, 1H), 7.00-6.68(m, 1H), 7.23-7.17(m, 2H), 7.46-7.43(m, 1H), 7.76(d, 1H), 7.93(dd, 1H), 8.22 (s, 1H), 8.40(d, 1H), 9.01(bs, 1H)
3-8		0.12 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 2.62(d, 3H), 2.81(s, 3H), 4.07-3.98(m, 1H), 4.52-4.45(m, 1H), 6.37(bs, 1H), 6.77-6.73 (m, 2H), 7.12(dd, 1H), 7.24-7.20(m, 1H), 7.30-7.27(m, 1H), 7.35(dd, 1H), 7.88(dd, 1H), 8.18 (s, 1H), 8.41(d, 1H), 9.19(bs, 1H)
3-9		0.28 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 2.62(d, 3H), 3.94(s, 3H), 4.49-4.43(m, 1H), 6.99-6.90 (m, 3H), 7.18-7.23(m, 1H), 7.31-7.24(m, 3H), 7.63(bs, 1H), 7.93-7.86(m, 1H), 8.28-8.23(m, 1H), 8.28 (s, 1H), 8.45(bs, 1H), 8.89(bs, 1H)
3-10		0.23 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 0.91(t, 3H), 1.37 (dd, 2H), 1.64-1.55 (m, 2H), 2.64-2.60 (m, 2H), 4.45-4.40 (m, 1H), 6.69 (bs, 1H), 7.23-7.10(m, 1H), 7.46-7.38 (m, 1H), 7.73 (d 1H), 7.92 (d, 1H), 8.21 (s, 1H), 8.38-8.46 (m, 1H), 9.09 (bs, 1H)
3-11		0.12 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 2.63 (d, 3H), 4.15-4.10 (m, 1H), 6.58 (bs, 1H), 7.31-7.10(m, 4H), 7.53-7.49 (m, 1H), 7.71(d 1H), 7.95 (d, 1H), 8.30-8.23 (m, 1H), 8.26 (s, 1H), 8.45 (d, 1H), 9.03 (bs, 1H)






3-12		0.4 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 2.09 (dd, 2H), 2.63 (d, 3H), 2.85(t, 2H), 2.96 (t, 2H), 4.46-4.43 (m, 2H), 6.73 (bs, 1H), 6.99 (d, 1H), 7.09 (dd, 1H), 7.25-7.20(m, 1H), 7.52 (dd, 1H), 7.74 (d 1H), 7.92 (dd, 1H), 8.22 (s, 1H), 8.42 (d, 1H), 9.02 (bs, 1H)
3-13		0.33 (AcOEt)	CDCl ₃ : 2.63 (d, 3H), 4.63-4.64 (m, 1H), 7.11(d, 2H), 7.18(dd, 1H), 7.42-7.34(m, 1H), 7.58-7.55(m, 1H), 7.96(d, 1H), 8.07(s, 1H), 8.19-8.10(m, 1H), 8.24(s, 1H), 9.15(s, 1H), 11.6-11.4(m, 1H)
3-14		0.28 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 2.63(d, 3H), 3.88(s, 3H), 3.89(s, 3H), 4.47-4.41(m, 1H), 6.60(d,1H), 6.92 (dd,1H), 7.64 (dd, 1H),7.66-7.61(m,1H), 7.89(d, 1H), 7.98(dd, 1H), 8.26(s, 1H), 8.43(d, 1H), 8.95(s, 1H)
3-15		0.30 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 2.63(d, 3H), 3.66(s, 3H), 3.85(s, 3H), 4.45-4.44(m, 1H), 6.48(dd,1H), 6.79(d,1H), 7.64(dd, 1H), 7.97(dd, 2H), 8.26(s, 1H), 8.44(d, 1H), 8.96(s, 1H)
3-16		0.22 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 2.17(s, 3H), 2.22(s, 3H), 2.64(s, 3H), 2.63(d, 3H), 4.46-4.44(m, 1H), 6.57(bs, 1H), 7.00(s,1H), 7.17(dd,1H), 7.44-7.40(m,1H), 7.44(s, 1H), 7.93(dd, 1H), 8.19(s, 1H), 8.43(d, 1H), 9.06(s, 1H)
3-17		0.46 (AcOEt)	CDCl ₃ : 2.22(s,3H), 2.63(d, 3H), 3.68(s, 3H), 3.89(s, 3H), 4.52-4.47(m, 1H), 6.51(s,1H), 6.74(s,1H), 7.12(s,1H), 7.16-7.12(m,1H), 7.40(dd, 1H), 7.91(dd, 1H), 8.19(s, 1H), 8.42(d, 1H), 9.12(s, 1H)
3-18		0.35 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 1.16(d, 6H), 2.25 (s, 3H), 2.62(d, 3H), 2.77(t, 1H), 4.49-4.48(m, 1H), 7.00(s,1H), 7.15(d,1H), 7.41-7.37(m,1H), 7.49(d,2H), 7.54(dd, 1H), 7.92(dd, 1H), 8.21(s, 1H), 8.32(d, 1H), 9.02(s, 1H)







3-19		0.23 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 2.63(d, 3H), 3.13-3.10 (m, 4H), 3.87(s, 3H), 3.89-3.86(m, 4H), 4.97-4.93(m, 1H), 6.41(dd, 1H), 6.52(d, 1H), 7.24-7.22(m, 1H), 7.32(s, 1H), 7.57(dd, 1H), 7.96(dd, 1H), 8.01(d, 1H), 8.14(s, 1H), 8.44(d, 1H), 8.98 (s, 1H)
3-20		0.36 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 2.22(s, 3H), 2.64(d, 3H), 3.00-3.2.97 (m, 4H), 3.76-3.74(m, 4H), 4.54-4.50(m, 1H), 6.64(d, 1H), 6.66(dd, 1H), 7.11(d, 1H), 7.18(dd, 1H), 7.37(d, 1H), 7.46(dd, 1H), 7.93(dd, 1H), 8.22(s, 1H), 8.42(d, 1H), 9.09 (s, 1H)
3-21			
3-22		0.27 (AcOEt)	CDCl ₃ : 2.33(s, 3H), 2.65(d, 3H), 3.60-3.45(m, 8H), 4.53-4.49(m, 1H), 6.74(s, 1H), 7.11(d, 1H), 7.22-7.18(m, 1H), 7.58-7.54(m, 1H), 7.94(dd, 1H), 8.00(d, 1H), 8.22(s, 1H), 8.37(d, 1H), 9.13(s, 1H)
3-23		0.38 (AcOEt)	CDCl ₃ : 1.24-1.08(m, 2H), 1.46-1.32(m, 2H), 1.76-1.67(m, 2H), 1.98-1.90(m, 2H), 2.33(s, 3H), 2.64(d, 3H), 3.95-3.90(m, 1H), 4.49-4.47(m, 1H), 5.89-5.80(m, 1H), 6.66(s, 1H), 7.15(dd, 1H), 7.48-7.31(m, 2H), 7.91(dd, 1H), 8.12(s, 1H), 8.23(s, 1H), 8.41(d, 1H), 9.18(s, 1H)
3-24		0.11 (AcOEt)	CDCl ₃ : 2.35(s, 3H), 2.71(s, 3H), 3.07-2.73(m, 2H), 3.86-3.31(m, 6H), 6.85(s, 1H), 7.10(d, 1H), 7.24-7.19(m, 1H), 7.52-7.48(m, 1H), 7.66-7.59(m, 2H), 7.93(d, 1H), 8.06(s, 1H), 8.27-8.21(m, 1H), 8.23(s, 1H), 9.11(s, 1H)








3-25		0.5 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 2.52(d, 3H), 2.62(s, 3H), 4.36-4.32(m, 1H), 6.74(s, 1H), 6.87(d, 2H), 7.00-6.91(m, 2H), 7.00-6.97(m, 2H), 7.38(dd, 2H), 7.86(dd, 1H), 7.98(s, 1H), 8.23(s, 1H), 8.28(d, 1H), 9.04(s, 1H)
3-26		0.45 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 1.62-1.34(m, 6H), 2.13(s, 3H), 2.56(d, 3H), 3.01-2.87(m, 4H), 4.54-4.38(m, 1H), 6.59(s, 1H), 6.69-6.59(m, 1H), 7.02(d, 1H), 7.10-7.07(m, 1H), 7.37(dd, 1H), 7.84(dd, 1H), 8.15(s, 1H), 8.34(d, 1H), 9.01(s, 1H)
3-27		0.45 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 2.32(s, 3H), 2.58(d, 3H), 3.75(s, 3H), 4.37-4.44(m, 1H), 6.77-6.73(m, 1H), 6.89-6.82(m, 1H), 6.97-6.91(m, 2H), 6.96(d, 1H), 7.20(dd, 1H), 7.25-7.24(m, 1H), 7.33-7.29(m, 1H)
3-28		0.35 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 2.34(s, 3H), 2.64(d, 3H), 3.81(s, 3H), 4.57-4.50(m, 1H), 6.76(bs, 1H), 6.91-6.84(m, 4H), 7.04(d, 1H), 7.83(dd, 1H), 8.06(d, 1H), 8.19(dd, 1H), 8.23(s, 1H), 9.00(s, 1H)
3-29		0.45 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 1.50(t, 3H), 2.62(d, 3H), 4.17(dd, 2H), 4.51-4.44(m, 1H), 6.95-6.89(m, 2H), 6.94(d, 1H), 7.16(dd, 1H), 7.31-7.23(m, 5H), 7.67(s, 1H), 7.11(dd, 1H), 7.23(d, 2H), 7.65(s, 1H), 7.88(dd, 1H), 8.28-8.23(m, 1H), 8.28(s, 1H), 8.43(s, 1H), 8.89(s, 1H)
3-30		0.45 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 1.49(t, 3H), 2.63(d, 3H), 3.85(s, 3H), 4.16(dd, 2H), 4.55-4.48(m, 1H), 6.81(dd, 1H), 6.95-6.91(m, 3H), 7.11(dd, 1H), 7.23(d, 2H), 7.65(s, 1H), 7.90-7.88(m, 1H), 8.28-8.26(m, 1H), 8.27(s, 1H), 8.39(s, 1H), 8.90(s, 1H)
3-31		0.29 (<i>n</i> -hexane: AcOEt=1:1)	¹ H-NMR : (CDCl ₃) 1.83-1.72 (4H, m), 2.63 (3H, d), 2.66-2.62 (2H, m), 2.80 (2H, t), 4.41-4.44 (1H, m), 6.64 (1H, br.s), 6.92 (1H, d), 7.09 (1H, dd), 7.18 (1H, dd), 7.45 (1H, dd), 7.59 (1H, dd), 7.92 (1H, d), 8.20 (1H, s), 8.42 (1H, d), 9.08 (1H, br.s).

3-32		0.3 (n-hexane: AcOEt=1:1)	DMSO-d ₆ : 2.43(s, 3H), 2.80-2.82(m, 4H), 3.61-3.64(m, 4H), 3.75(s, 3H), 6.62(dd, 1H), 6.93(d, 1H), 7.46(d, 1H), 7.54(dd, 1H), 7.77(dd, 2H), 8.14(bs, 1H), 8.32(s, 1H), 8.38-8.30(m, 1H), 9.14(bs, 1H)
3-33		0.61 (MeOH: CH ₂ Cl ₂ =1: 1)	DMSO-d ₆ : 1.59-1.68(m, 2H), 1.88-1.98(m, 2H), 2.13-2.25(m, 2H), 2.19(s, 3H), 2.43(s, 3H), 2.60-2.70(m, 2H), 3.75(s, 3H), 4.32-4.40(m, 1H), 6.51(dd, 1H), 6.64(d, 1H), 7.20(dd, 1H), 7.39(d, 1H), 7.75(dd, 1H), 7.70-7.78(s, 1H), 8.22(s, 1H), 8.26(s, 1H), 8.38-8.41(m, 1H), 9.22(s, 1H)
3-34		0.17 (AcOEt)	CDCl ₃ : 2.11(s, 3H), 2.68(d, 3H), 2.76-2.83(m, 2H), 2.89-2.97(m, 2H), 3.47-3.55(m, 2H), 3.58-3.66(m, 2H), 3.86(s, 3H), 4.70-4.78(m, 1H), 6.53(dd, 1H), 6.81(d, 1H), 7.23(dd, 1H), 7.54-7.62(m, 2H), 7.97(dd, 1H), 8.02-8.03(m, 1H), 8.29(s, 1H), 8.40(d, 1H), 8.99(bs, 1H)
3-35		0.22 (AcOEt only)	DMSO-d ₆ : 2.40-2.48(m, 7H), 2.63(t, 2H), 3.50-3.58(m, 4H), 3.77(s, 3H), 3.91(t, 2H), 6.60(dd, 1H), 6.93(d, 1H), 7.28(dd, 1H), 7.56(d, 1H), 7.60(dd, 1H), 7.75-7.80(m, 1H), 7.80(dd, 1H), 8.10(s, 1H), 8.35(s, 1H), 8.40(d, 1H), 9.21(s, 1H)
3-36		0.4 (n-hexane: AcOEt=1:1)	DMSO-d ₆ : 2.43(s, 3H), 7.03-7.08(m, 1H), 7.21-7.23(m, 1H), 7.25-7.36(m, 1H), 7.47-7.57(m, 2H), 7.74-7.77(m, 2H), 8.28(s, 1H), 8.35(d, 1H), 9.09(s, 1H), 9.24(s, 1H)
3-37		0.4 (n-hexane: AcOEt=1:1)	CDCl ₃ : 2.64(d, 3H), 4.53-4.54(m, 1H), 6.88-6.93(m, 1H), 7.14-7.28(m, 3H), 7.54-7.58(m, 1H), 7.95-7.98(m, 1H), 8.16-8.21(m, 1H), 8.24(s, 1H), 8.33-8.36(m, 1H), 9.05(s, 1H)
3-38		0.42 (n-hexane: AcOEt=1:1)	CDCl ₃ : 2.64(d, 3H), 4.46-4.47(m, 1H), 6.63-6.68(m, 1H), 7.30-7.32(m, 2H), 7.55(s, 1H), 7.64-7.68(m, 1H), 7.97-7.99(m, 1H), 8.20-8.39(m, 3H), 9.03(s, 1H)






3-39		562, 564 [M+1] ⁺	CDCl ₃ : 2.37(s, 3H), 2.58-2.64(m, 7H), 3.15-3.18(m, 4H), 3.87(s, 3H), 4.60-4.65(m, 1H), 6.43(dd, 1H), 6.44-6.54(m, 1H), 7.22(d, 1H), 7.30(s, 1H), 7.57(dd, 1H), 7.94-7.99(m, 2H), 8.18(s, 1H), 8.45(d, 1H), 8.95(s, 1H)
3-40		572, 574 [M+1] ⁺	DMSO-d ₆ : 1.79-1.88(m, 2H), 1.98-2.02(m, 2H), 2.43(s, 3H), 3.02-3.08(m, 3H), 3.28-3.39(m, 2H), 3.76(s, 3H), 6.47(dd, 1H), 6.65(d, 1H), 7.22(dd, 1H), 7.39(d, 1H), 7.45-7.50(m, 1H), 7.74-7.77(m, 2H), 8.18(s, 1H), 8.22(s, 1H), 8.41-8.44(m, 1H), 9.21(bs, 1H)
3-41		565, 567 [M+1] ⁺	DMSO-d ₆ : 2.44(d, 3H), 2.69-2.71(m, 4H), 3.49-3.52(m, 4H), 3.76(s, 3H), 6.45(dd, 1H), 6.62(d, 1H), 7.23(ddd, 1H), 7.38(d, 1H), 7.46-7.50(m, 1H), 7.72-7.77(m, 2H), 8.19(s, 1H), 8.22(s, 1H), 8.42-8.45(m, 1H), 9.22(s, 1H)
3-42		595, 597 [M+1] ⁺	DMSO-d ₆ : 2.44(s, 3H), 3.31(s, 6H), 3.48-3.53(m, 8H), 3.72(s, 3H), 6.24(dd, 1H), 6.37(d, 1H), 7.18-7.21(m, 2H), 7.40-7.55(m, 1H), 7.72-7.76(m, 2H), 8.17-8.19(m, 2H), 8.40-8.50(m, 1H), 9.23(s, 1H)
3-43		590, 592 [M+1] ⁺	DMSO-d ₆ : 1.64-1.71(m, 2H), 1.75-1.82(m, 2H), 2.21-2.28(m, 1H), 2.43(d, 3H), 2.62-2.67(m, 2H), 3.68-3.74(m, 2H), 3.76(s, 3H), 6.45(dd, 1H), 6.63(d, 1H), 6.75-6.81(m, 1H), 7.20(ddd, 1H), 7.25-7.30(m, 1H), 7.35(d, 1H), 7.45-7.52(m, 1H), 7.70-7.77(m, 2H), 8.18(s, 1H), 8.21(s, 1H), 8.40-8.47(m, 1H), 9.22(s, 1H)






3-44		597, 599 [M+1] ⁺	DMSO-d ₆ : 2.44(s, 3H), 3.12-3.17(m, 4H), 3.68-3.85(m, 4H), 3.79(s, 3H), 6.55(dd, 1H), 6.71(d, 1H), 7.19-7.25(m, 1H), 7.43(d, 1H), 7.46-7.53(m, 1H), 7.73-7.78(m, 2H), 8.19-8.22(m, 1H), 8.22(s, 1H), 8.38-8.45(m, 1H), 9.20(bs, 1H)
3-45		600, 602 [M+1] ⁺	DMSO-d ₆ : 1.85-1.95(m, 2H), 2.19(t, 2H), 2.25-2.35(m, 4H), 2.43(s, 3H), 3.52-3.64(m, 4H), 4.19(t, 2H), 6.65(d, 1H), 7.05(dd, 1H), 7.20(d, 1H), 7.23(ddd, 1H), 7.27(d, 1H), 7.40-7.46(m, 1H), 7.42(d, 1H), 7.70-7.75(m, 1H), 7.76(dd, 1H), 8.32(s, 1H), 8.45(d, 1H), 9.22(s, 1H), 9.23(s, 1H)
3-46		590, 592 [M+1] ⁺	DMSO-d ₆ : 2.05(s, 3H), 2.44(s, 3H), 3.08-3.17(m, 4H), 3.55-3.63(m, 4H), 3.77(s, 3H), 6.48(dd, 1H), 6.67(d, 1H), 7.23(dd, 1H), 7.41(d, 1H), 7.45-7.52(m, 1H), 7.76(dd, 1H), 7.72-7.78(m, 1H), 8.19(s, 1H), 8.22(s, 1H), 8.40-8.47(m, 1H), 9.22(bs, 1H)
3-47		548, 550 [M+1] ⁺	DMSO-d ₆ : 2.43(s, 3H), 2.82-2.87(m, 4H), 2.99-3.15(m, 4H), 3.76(s, 3H), 6.43(dd, 1H), 6.61(d, 1H), 7.22(dd, 1H), 7.36(d, 1H), 7.43-7.51(m, 1H), 7.75(dd, 1H), 8.17(s, 1H), 8.21(s, 1H), 8.38-8.45(m, 1H), 9.12-9.28(m, 1H)
3-48		MS 530, 532	CDCl ₃ : 2.65 (d, 3H), 3.96 (s, 3H), 4.40-4.48 (m, 1H), 6.85-6.88 (m, 2H), 7.22 (d, 1H), 7.25-7.31 (m, 1H), 7.56-7.65 (m, 3H), 7.79 (s, 1H), 8.00 (dd, 1H), 8.29 (s, 1H), 8.39 (dd, 1H), 9.00 (s, 1H).







3-49		R _f (AcOEt: MeOH=9:1) 0.20	CDCl ₃ : 2.18-2.50 (m, 4H), 2.28 (s, 3H), 2.65 (d, 3H), 3.10-3.75 (m, 4H), 3.93 (s, 3H), 4.50-4.61 (m, 1H), 6.89 (d, 1H), 7.06 (dd, 1H), 7.59-7.67 (m, 2H), 7.93-7.97 (m, 1H), 8.26 (s, 1H), 8.37-8.43 (m, 2H), 9.02 (s, 1H).
3-50		R _f 0.4 (Hexane/Ac OEt=1/1)	CDCl ₃ : 2.63 (d, 3H), 3.90 (s, 3H), 4.00 (s, 3H), 4.39-4.47 (m, 1H), 6.23 (d, 1H), 7.00 (s, 1H), 7.22-7.25 (m, 1H), 7.57 (dd, 1H), 7.96 (dd, 1H), 8.22 (s, 1H), 8.25 (d, 1H), 8.37 (d, 1H), 8.96 (s, 1H)
3-51		MS 535, 537	CDCl ₃ : 1.17 (t, 3H), 1.71-1.79 (m, 1H), 2.28 (s, 3H), 2.62 (d, 3H), 3.41 (q, 2H), 3.46 (t, 2H), 3.79 (q, 2H), 4.41-4.48 (m, 1H), 6.43 (s, 1H), 6.10-6.18 (m, 2H), 7.15 (dd, 1H), 7.33 (d, 1H), 7.35-7.42 (m, 1H), 7.90 (dd, 1H), 8.16 (s, 1H), 8.45 (d, 1H), 9.07 (s, 1H).
3-52		R _f	CDCl ₃ : 2.66 (d, 3H), 3.91 (s, 3H), 4.41-4.47 (m, 1H), 6.80 (d, 1H), 6.92 (dd, 1H), 7.26-7.35 (m, 1H), 7.54 (s, 1H), 7.76 (dd, 1H), 8.00 (dd, 1H), 8.27-8.32 (m, 2H), 8.38 (dd, 1H), 8.97 (s, 1H).
3-53		MS 491, 493	CDCl ₃ : 2.26 (s, 3H), 2.62 (d, 3H), 2.68 (s, 6H), 4.72 (q, 1H), 6.78 (s, 1H), 6.89 (d, 1H), 7.12 (d, 1H), 7.15 (d, 1H), 7.40-7.47 (m, 2H), 7.91 (dd, 1H), 8.40 (s, 1H), 8.41 (dd, 1H), 9.11 (s, 1H).
3-54		MS 525, 527	CDCl ₃ : 2.04 (s, 3H), 2.65 (d, 3H), 4.42-4.48 (m, 1H), 6.79 (s, 1H), 6.96-7.00 (m, 2H), 7.28-7.34 (m, 4H), 7.87-7.91 (m, 1H), 8.18 (s, 1H), 8.23-8.26 (m, 2H), 8.53 (d, 2H), 9.07 (s, 1H).

3-55		Rf (Hexane:AcOEt=3:1) 0.19	CDCl ₃ : 1.34 (t, 3H), 1.44 (t, 3H), 2.63 (d, 3H), 3.81 (q, 2H), 4.06 (q, 2H), 4.46 (q, 1H), 6.43 (dd, 1H), 6.76 (d, 1H), 7.63-7.69 (m, 2H), 7.94 (d, 1H), 7.98 (dd, 1H), 8.42 (d, 1H), 8.93 (s, 1H).
3-56		MS 570, 572	CDCl ₃ : 2.63 (d, 3H), 3.85 (s, 3H), 3.93 (s, 3H), 4.52 (q, 1H), 6.78-6.83 (m, 2H), 6.93 (d, 2H), 6.99-7.02 (m, 1H), 7.11-7.15 (m, 1H), 7.21-7.27 (m, 1H), 7.61 (s, 1H), 7.87-7.92 (m, 1H), 8.26 (s, 1H), 8.20-8.30 (m, 1H), 8.38-8.41 (m, 1H), 8.92 (s, 1H).
3-57		Rf (Hexane:AcOEt=3:1) 0.16	CDCl ₃ : 1.44 (t, 3H), 2.65 (d, 3H), 2.79-2.89 (m, 4H), 3.65-3.74 (m, 4H), 4.07 (q, 2H), 4.52 (q, 4H), 6.48 (dd, 1H), 6.80 (d, 1H), 7.20-7.25 (m, 1H), 7.55-7.67 (m, 2H), 7.92-7.98 (m, 2H), 8.29 (s, 1H), 8.43 (d, 1H), 8.95 (s, 1H).
3-58		Rf 0.17 (Hexane/AcOEt=1/1)	CDCl ₃ : 1.46 (t, 3H), 2.63 (d, 3H), 3.08-3.13 (m, 4H), 3.83-3.90 (m, 4H), 4.09 (q, 2H), 4.46 (q, 1H), 6.39 (dd, 1H), 6.51 (d, 1H), 7.21-7.28 (m, 1H), 7.37 (s, 1H), 7.58 (dd, 1H), 7.97 (dd, 1H), 8.03 (d, 1H), 8.21 (s, 1H), 8.46 (d, 1H), 8.94 (s, 1H).
3-59		MS 538, 540	CDCl ₃ : 2.63 (d, 3H), 3.44 (s, 3H), 3.65 (s, 3H), 3.69-3.73 (m, 2H), 4.10-4.15 (m, 2H), 4.40 (q, 1H), 6.45 (dd, 1H), 6.85 (d, 1H), 7.19-7.25 (m, 1H), 7.61 (dd, 1H), 7.88 (s, 1H), 7.93-7.97 (m, 2H), 8.27 (s, 1H), 8.46 (d, 1H), 8.95 (s, 1H).
3-60		Rf (AcOEt) 0.54	CDCl ₃ : 2.63 (d, 3H), 3.67 (s, 3H), 4.18 (t, 2H), 4.38-4.49 (m, 3H), 6.46 (dd, 1H), 6.81 (d, 1H), 7.60-7.69 (m, 2H), 7.92-7.99 (m, 2H), 8.27 (s, 1H), 8.49 (d, 1H), 9.00 (s, 1H).
3-61		Rf (Hexane:AcOEt=2:1) 0.46	CDCl ₃ : 1.44 (t, 3H), 2.63 (d, 3H), 3.64 (s, 3H), 4.07 (q, 2H), 4.47 (q, 1H), 6.45 (dd, 1H), 6.78 (d, 1H), 7.21-7.28 (m, 1H), 7.40-7.48 (m, 2H), 7.93-7.99 (m, 2H), 8.26 (s, 1H), 8.44 (d, 1H), 8.96 (s, 1H).

3-62		Rf (Hexane:Ac OEt=3:1) 0.31	CDCl ₃ : 1.36 (d, 6H), 2.63 (d, 3H), 3.63 (s, 3H), 4.41-4.52 (m, 2H), 6.45 (dd, 1H), 6.81 (d, 1H), 7.21-7.26 (m, 1H), 7.59-7.68 (m, 2H), 7.91-7.98 (m, 2H), 8.26 (s, 1H), 8.45 (d, 1H), 8.96 (s, 1H).
3-63		Rf (Hexane: AcOEt=3:1) 0.40	CDCl ₃ : 1.07 (t, 3H), 1.84 (m, 2H), 6.63 (d, 3H), 3.64 (s, 3H), 3.96 (t, 2H), 4.40-4.49 (m, 1H), 6.46 (dd, 1H), 6.79 (d, 1H), 7.20-7.27 (m, 1H), 7.58-7.66 (m, 2H), 7.94-7.97 (m, 2H), 8.26 (s, 1H), 8.45 (d, 1H), 8.97 (s, 1H).
3-64		Rf (Hexane:Ac OEt=3:1) 0.19	CDCl ₃ : 2.62 (d, 3H), 6.68 (s, 6H), 3.84 (s, 3H), 4.41-4.48 (m, 1H), 6.36 (dd, 1H), 6.80 (d, 1H), 7.17-7.24 (m, 1H), 7.51-7.62 (m, 2H), 7.83 (s, 1H), 7.95 (dd, 1H), 8.27 (s, 1H), 8.3*9-8.45 (m, 1H), 8.91 (s, 1H).
3-65		Rf (Hexane:Ac OEt=1:1) 0.12	CDCl ₃ : 2.66 (d, 3H), 3.97 (s, 3H), 4.47-4.55 (m, 1H), 6.96-7.10 (m, 3H), 7.21-7.24 (m, 1H), 7.66 (s, 1H), 7.93 (dd, 1H), 8.25 (d, 1H), 8.31 (s, 1H), 8.47 (d, 2H), 8.59 (s, 1H), 8.96 (s, 1H).
3-66		MS 541, 543	CDCl ₃ : 2.65 (d, 3H), 3.96 (s, 3H), 4.61-4.71 (m, 1H), 6.89-7.05 (m, 3H), 7.16 (dd, 1H), 7.15-7.23 (m, 1H), 7.60 (d, 1H), 7.65 (s, 1H), 7.89 (d, 1H), 8.21 (d, 1H), 8.28 (d, 1H), 8.51 (br. s, 2H), 8.57 (s, 1H), 8.93 (s, 1H).
3-67		MS 541, 543	CDCl ₃ : 2.65 (d, 3H), 3.96 (s, 3H), 4.51 (q, 1H), 6.90-7.06 (m, 3H), 7.11-7.16 (m, 1H), 7.38 (d, 1H), 7.50-7.61 (m, 2H), 7.62-7.67 (m, 1H), 7.89 (dd, 1H), 8.29 (s, 1H), 8.34 (d, 1H), 8.53 (d, 1H), 8.79 (br.s, 1H), 8.94 (s, 1H).
3-68		LC-MS 590	CDCl ₃ : 1.45-1.59 (m, 2H), 1.70-1.78 (m, 1H), 1.82-1.90 (m, 1H), 2.38-2.50 (m, 1H), 2.43 (s, 3H), 2.62-2.77 (m, 2H), 3.56-3.70 (m, 2H), 3.76 (s, 3H), 6.46 (dd, 1H), 6.63 (d, 1H), 6.82-6.88 (br, 1H), 7.22 (dd, 1H), 7.31-7.40 (m, 2H), 7.43-7.51 (m, 1H), 7.50-7.80 (m, 2H), 8.14-8.20 (br, 1H), 8.21 (s, 1H), 8.39-8.48

			(m, 1H), 9.16-9.26 (br, 1H)
3-69		0.34 (CH ₂ Cl ₂ :MeOH=9:1)	CDCl ₃ : 1.58-1.82 (br, 7H), 1.88-2.03 (br, 3H), 2.44-2.45 (m, 5H), 3.42-3.52 (m, 3H), 3.75 (s, 3H), 6.66 (dd, 1H), 6.92 (d, 1H), 7.28 (dd, 1H), 7.44 (br, 1H), 7.51 (dd, 1H), 7.79-7.81 (m, 2H), 8.18 (s, 1H), 8.32 (s, 1H), 8.35-8.37 (m, 1H), 9.17 (s, 1H)
3-70		Ms:607, 609	DMSO-d ₆ : 1.84-1.92(m, 2H), 2.34-2.41(m, 4H), 2.41-2.45(m, 3H), 2.44(t, 2H), 3.58(t, 4H), 3.75(s, 3H), 4.02(t, 2H), 6.48(dd, 1H), 6.63(d, 1H), 7.21(dd, 1H), 7.41(d, 1H), 7.46(dd, 1H), 7.72-7.78(m, 1H), 7.76(dd, 1H), 8.22(s, 1H), 8.25(s, 1H), 8.40(d, 1H), 9.22(s, 1H)
3-71		Ms:591, 593	DMSO-d ₆ : 1.84-1.92(m, 2H), 2.14(s, 3H), 2.35-2.4 (m, 4H), 2.43(t, 2H), 2.44(d, 3H), 3.58(t, 4H), 4.01(t, 2H), 6.77(dd, 1H), 6.82(d, 1H), 7.17(dd, 1H), 7.20(d, 1H), 7.3-7.39(m, 1H), 7.71-7.77(m, 2H), 8.2(s, 1H), 8.35-8.44(m, 1H), 8.71(s, 1H), 9.27(s, 1H)
3-72		Ms:620, 622	DMSO-d ₆ : 1.82-1.9 (m, 2H), 2.13-2.17 (m, 3H), 2.25-2.47(m, 13H), 3.75 (s, 3H), 4.01 (t, 2H), 6.47 (dd, 1H), 6.63(d, 1H), 7.19-7.24 (m, 1H), 7.41 (d, 1H), 7.43-7.5(m, 1H), 7.70-7.79(m, 2H), 8.22(s, 1H), 8.25(brs, 1H), 8.37-8.44(m, 1H), 9.22(s, 1H)
3-73		Ms:607, 609	DMSO-d ₆ : 1.78 (t, 2H), 2.32-2.36 (m, 4H), 2.35-2.38 (m, 3H), 3.54-3.59 (m, 4H), 3.74 (t, 3H), 3.78 (s, 3H), 6.38-6.42 (m, 1H), 6.85 (d, 1H), 6.86-6.95 (m, 1H), 7.33-7.43(m, 2H), 7.63-7.68 (m, 1H), 7.85-8.15 (m, 3H), 8.64-8.8 (m, 1H).

3-74		Ms:605, 607	DMSO-d6: 1.47-1.67(m, 2H), 1.84-2.01(m, 2H), 2.03 (s, 3H), 2.41-2.46 (m, 3H), 3.23-3.39 (m, 2H), 3.65-3.73 (m, 1H), 3.81(s, 3H), 3.8-3.88 (m, 1H), 4.58-4.65 (m, 1H), 6.55 (dd, 1H), 6.68 (d, 1H), 7.2-7.26(m, 1H), 7.43(d, 1H), 7.42-7.51 (m, 1H), 7.7-7.8(m, 2H), 8.23 (s, 1H), 8.26 (brs, 1H), 8.37-8.44(m, 1H), 9.22(brs, 1H)
3-75		Ms:605, 607	DMSO-d6: 1.38-1.6(m, 2H), 1.74-1.9(m, 2H), 2.0 (s, 3H), 2.42-2.47 (m, 3H), 3.12-3.3 (m, 2H), 3.55-3.65 (m, 1H), 3.7-3.8 (m, 1H), 3.78 (s, 3H), 4.27-4.34 (m, 1H), 6.65 (dd, 1H), 6.94 (d, 1H), 7.24-7.3 (m, 1H), 7.53-7.63(m, 2H), 7.74-7.83 (m, 2H), 8.09 (brs, 1H), 8.35 (s, 1H), 8.38(d, 1H), 9.19(brs, 1H)
3-76		Ms:577, 579	DMSO-d6: 1.51-1.61 (m, 2H), 1.79-1.87 (m, 2H), 2.03-2.11 (m, 2H), 2.14 (s, 3H), 2.42-2.47 (m, 3H), 2.52-2.6 (m, 2H), 3.77 (s, 3H), 4.02-4.09 (m, 1H), 6.6 (dd, 1H), 6.92 (d, 1H), 7.24-7.3 (m, 1H), 7.52-7.6(m, 2H), 7.74-7.82 (m, 2H), 8.08 (brs, 1H), 8.34 (s, 1H), 8.4 (d, 1H), 9.2 (brs, 1H)
3-77		Rf : 0.4 (n-hexane: AcOEt=7:3)	DMSO-d6: 2.41-2.45 (m, 3H), 6.89-6.96 (m, 1H), 6.69(bs, 1H), 7.24-7.33 (m, 2H), 7.51-7.57 (m, 1H), 7.63-7.7 (m, 1H), 7.73-7.78 (m, 1H), 7.79 (dd, 1H), 8.37(s, 1H), 8.41(d, 1H), 9.21 (brs, 1H), 9.24 (brs, 1H)
3-78		Ms:563, 565	DMSO-d6: 1.33-1.43 (m, 2H), 1.79-1.86 (m, 2H), 2.43-2.46 (m, 3H), 2.46-2.53 (m, 2H), 2.87-2.94 (m, 2H), 3.77 (s, 3H), 4.07-4.14 (m, 1H), 6.59 (dd, 1H), 6.91 (d, 1H), 7.23-7.28 (m, 1H), 7.53-7.59 (m, 2H), 7.79 (dd, 1H), 8.03 (brs, 1H), 8.32 (s, 1H), 8.38 (d, 1H), 8.7-9.5 (brs, 1H)

3-79		Ms:563, 565	DMSO-d6: 1.41-1.51 (m, 2H), 1.88-1.95 (m, 2H), 2.41-2.45 (m, 3H), 2.54-2.63 (m, 2H), 2.92-3.0 (m, 2H), 3.75 (s, 3H), 4.35-4.43 (m, 1H), 6.50 (dd, 1H), 6.63 (d, 1H), 7.18-7.23 (m, 1H), 7.40 (d, 1H), 7.42-7.48 (m, 1H), 7.75 (dd, 1H), 8.21 (s, 1H), 8.22-8.25 (m, 1H), 8.37-8.42 (m, 1H), 8.9-9.5 (brs, 1H)
3-80		Ms:482, 484	DMSO-d6: 2.4-2.46 (m, 3H), 3.79 (s, 3H), 6.72 (ddd, 1H), 6.99 (dd, 1H), 7.21-7.26 (m, 1H), 7.47-7.53 (m, 1H), 7.59-7.64 (m, 1H), 7.76 (dd, 1H), 8.25 (s, 1H), 8.29-8.37 (m, 2H), 8.8-9.6 (m, 1H)
3-81		Ms:482, 484	DMSO-d6: 2.41-2.49 (m, 3H), 3.82 (s, 3H), 6.80 (ddd, 1H), 7.01 (dd, 1H), 7.3-7.35 (m, 1H), 7.56-7.63 (m, 1H), 7.7-7.8 (m, 1H), 7.82 (dd, 1H), 7.85 (dd, 1H), 8.16 (s, 1H), 8.35 (dd, 1H), 9.18 (brs, 1H)
3-82		Ms:563, 565	DMSO-d6: 1.73-1.82 (m, 1H), 2.23-2.34 (m, 4H), 2.34-2.42(m, 3H), 2.42-2.46 (m, 3H), 2.59 (dd, 1H), 2.62-2.68 (m, 1H), 2.80 (dd, 1H), 3.75 (s, 1H), 4.85-4.91(m, 1H), 6.42 (dd, 1H), 6.57(d, 1H), 7.19-7.24(m, 1H), 7.41 (d, 1H), 7.43-7.51(m, 1H), 7.68-7.79 (m, 2H), 8.22(s, 1H), 8.23(s, 1H), 8.37-8.43 (m, 1H), 9.21 (brs, 1H).
3-83		MS 544, 546	2.36 (s, 3H), 2.65 (d, 3H), 3.93 (s, 3H), 4.46-4.51 (m, 1H), 6.75-6.80 (m, 2H), 6.97-7.04 (m, 2H), 7.25-7.30 (m, 1H), 7.56-7.66 (m, 2H), 7.98 (dd, 1H), 8.29 (s, 1H), 8.36-8.44 (m, 2H), 9.01 (s, 1H).
3-84		MS 562, 564	CDCl3: 2.32 (s, 3H), 2.39-2.47 (m, 4H), 2.64 (d, 3H), 2.89-2.97 (m, 4H), 3.85 (s, 3H), 4.54-4.52 (m, 1H), 6.52 (dd, 1H), 6.79 (d, 1H), 7.22 (m, 1H), 7.52-7.64 (m, 2H), 7.94-7.99 (m, 2H), 8.28 (s, 1H), 8.42 (d, 1H), 8.93 (s, 1H).

Example 4: 2-[5-Bromo-2-(subst. phenylamino)-pyrimidin-4-ylamino]-N-propyl-benzene-sulfonamides

These compounds are prepared in analogy to Example 2 using 2-(5-bromo-2-chloro-pyrimidin-4-ylamino)-N-propyl-benzenesulfonamide and the corresponding aniline to give compounds No. 4-1 to 4-31 having the substituent Rx as listed under Example 3 for compounds No. 3-1 to 3-31.

Preparation of 2-(5-bromo-2-chloro-pyrimidin-4-ylamino)-N-propyl-benzenesulfonamide

To a solution of 5-bromo-2,4-dichloropyrimidine (90 μ L, 0.70 mmol) and 2-amino-N-propyl-benzenesulfonamide (100 mg, 0.47 mmol), sodium hydride (54.2 mg, 0.56 mmol) in DMSO (1.0 mL) is added and the resulting solution is stirred at 80°C for 3.0 h. The mixture is poured into water and extracted with ethyl acetate three times. The organic layer is washed with water and then brine, dried over sodium sulfate, and evaporated in vacuo. The residue is purified with silica gel column chromatography (n-hexane : ethyl acetate = 5 : 1) to afford the title compound as a slightly yellow solid.

$^1\text{H-NMR}$ (δ , ppm) : 0.89 (t, 3H), 1.41 (q, 2H), 3.56 (t, 2H), 4.92 (br.s, 2H), 6.71 (dd, 1H), 6.77 (dd, 1H), 7.33 (dd, 1H), 7.54 (dd, 1H), 8.79 (s, 1H)

R_f (hexane : ethyl acetate = 1:1): 0.64.

Example 5: 2-[5-Trifluoromethyl-2-(subst. phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzenesulfonamides


These compounds are prepared in analogy to Example 2 using 2-(2-chloro-5-trifluoromethyl-pyrimidin-4-ylamino)-N-methyl-benzenesulfonamide and the corresponding aniline to give compounds No. 5-1 to 5-31 having the substituent Rx as listed under Example 3 for compounds No. 3-1 to 3-31.

Preparation of 2-(2-chloro-5-trifluoromethyl-pyrimidin-4-ylamino)-N-methyl-benzenesulfonamide

To a solution of 2,4-dichloro-5-trifluoromethyl-pyrimidine (386 mg, 1.79 mmol) in acetonitrile (10 mL), 2-amino-N-methyl-benzenesulfonamide (333 mg, 1.79 mmol) and 1,8-diaza[5.4.0]-bicyclo-7-undecene (280 μ L, 1.88 mmol) are added successively at ambient temperature. After stirring for 15 h at room temperature, dichloromethane (30 mL) is added to the mixture, and the solution is washed with saturated aqueous sodium hydrogen carbonate and saturated aqueous sodium chloride, dried over magnesium sulfate, and evaporated in vacuo. The resulting solid is purified by flash chromatography.

$^1\text{H NMR}$ (CDCl_3) δ : 3.73(s, 3H), 6.67-6.69(m, 1H), 6.72-6.73(m, 1H), 7.27-7.31(m, 1H), 7.78 (dd, 1H), 8.60(s, 1H). R_f (hexane : ethyl acetate = 1:1): 0.28.

Example 6: 2-[5-Bromo-2-(2,3-[difluoromethylenedioxy]phenylamino)-pyrimidin-4-ylamino]-benzenesulfonamide



This compound is obtained as a side product formed by N-demethylation on reaction of 2-(5-bromo-2-chloropyrimidin-4-ylamino)-N-methyl-benzenesulfonamide with 2,3-(difluoromethylenedioxy)aniline following the procedure of Example 2. It may also be prepared by reaction of 2-(5-bromo-2-chloropyrimidin-4-ylamino)benzenesulfonamide with 2,3-(difluoromethylenedioxy)-aniline.


R_f (*n*-hexane: ethyl acetate = 1:1): 0.46.

¹H-NMR : (CDCl₃) 4.83 (bs, 2H), 6.77 (dd, 1H), 6.86 (s, 1H), 6.97 (dd, 1H), 7.31-7.24 (m, 1H), 7.57 (dd, 1H), 7.81 (d, 1H), 8.02 (dd, 1H), 8.28 (d, 1H), 8.29 (s, 1H), 8.88 (s, 1H).

Preparation of 2-(5-bromo-2-chloropyrimidin-4-ylamino)benzenesulfonamide: To a solution of 5-bromo-2,4-dichloropyrimidine (300 mg, 1.32mmol) and 2-amino-benzenesulfonamide (340 mg, 1.97 mmol) in 2-propanol (3 mL), concentrated hydrochloric acid (0.06 mL) is added and the mixture is stirred at 90°C for 4.5 hours. The mixture is poured into aqueous sodium hydrogen carbonate and extracted with ethyl acetate three times. The organic layer is washed with water, dried over sodium sulfate, and evaporated in vacuo. The residue is purified by column chromatography (hexane : ethyl acetate = 2:1) to afford the title compound.


R_f (hexane : ethyl acetate = 1:1): 0.55. ¹H-NMR (400MHz, CDCl₃) δ : 4.78 (br.s, 2H), 7.22 (dd, 1H), 7.61 (ddd, 1H), 7.95 (dd, 1H), 8.35 (s, 1H), 8.35 (d, 1H), 9.18 (s, 1H).

Example 7A: 2-[5-Chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzamide













To a suspension of 2-(2,5-dichloro-pyrimidin-4-yl-amino)-*N*-methyl-benzamide (5.05 g, 17.0 mmol) in 90 mL of 2-methoxyethanol are added 2-methoxy-4-morpholinoaniline dihydrochloride (4.56 g, 16.2 mmol) and 17.0 mL of 1N ethanolic solution of hydrogen chloride (17.0 mmol). After the reaction mixture is stirred at 110°C for 4 hours and cooled to room temperature, the mixture is neutralized with 1N aqueous NaOH solution and extracted with EtOAc (100 mL × 3). The organic layer is washed with brine, dried over Na₂SO₄ and concentrated under reduced pressure. The resulting black solid is washed with EtOH (90 mL), then purified with silica gel column chromatography (CH₂Cl₂ to CH₂Cl₂: AcOEt=1:2) to give 2-[5-chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-*N*-methyl-benzamide as a pale yellow solid. ¹H-NMR (400MHz, DMSO-d₆, δ): 2.80 (d, 3H, J = 4.52 Hz), 3.10-3.20 (m, 4H), 3.78 (s, 3H), 3.70-3.80 (m, 4H), 6.49 (dd, 1H, J = 8.56, 2.52 Hz), 6.66 (d, 1H, J = 2.52 Hz), 7.08 (dd, 1H, J = 8.04, 8.04 Hz), 7.44 (d, 1H, J = 8.56 Hz), 7.71 (dd, 1H, J = 8.04, 1.48 Hz), 8.10 (s, 1H), 8.13 (s, 1H), 8.59 (d, 1H, J = 8.04 Hz) 8.68-8.75 (m, 1H), 11.59 (s, 1H). MS *m/z* 469, 471 (M+1)⁺.





The following 2-[5-Chloro-2-(substituted phenylamino)-pyrimidin-4-ylamino]-*N*-methyl-benzamide are prepared from 2-(2,5-Dichloro-pyrimidin-4-ylamino)-*N*-methyl-benzamide and the corresponding aniline following the procedure of Example 7A.












Expt No.	Rx	Rf (solvent) or MS	NMR (400MHz), δ (ppm)






7-1		MS: m/z 550, 552 ($M+1$)	DMSO- d_6 : 1.44-1.33 (m, 2H), 1.64-1.45 (m, 6H), 1.73-1.89 (m, 2H), 2.34-2.44 (m, 1H), 2.43-2.55 (m, 4H), 2.65 (t, 2H), 2.80 (d, 3H), 3.75 (s, 3H), 3.72-3.75 (m, 2H), 6.48 (dd, 1H), 6.62 (d, 1H), 7.06 (dd, 1H), 7.32 (dd, 1H), 7.39 (d, 1H), 7.71 (dd, 1H), 8.09(s, 1H), 8.60 (d, 1H), 8.70 (d, 1H), 11.58 (s, 1H)
7-2		0.3 (MeOH: AcOEt =5:95)	$CDCl_3$: 1.70-1.97(m, 4H) ,2.62-2.79(m, 1H), 3.04(d, 3H), 3.02-3.18(m, 2H), 3.23-3.33(m, 2H), 3.88 (s, 3H), 5.39-5.47(m, 1H), 6.15-6.24(m, 1H), 6.55-6.62(m, 2H), 6.74-6.82(m, 1H), 7.09 (dd, 1H),7.23-7.32 (m, 1H), 7.46-7.52(m, 2H), 8.09(s, 1H), 8.15(d, 1H), 8.68(d, 1H) 11.0(bs, 1H)
7-3		MS (ESI) m/z 482, 484 ($M+1$) ⁺	DMSO- d_6 : 2.24 (s, 3H), 2.45-2.55 (m, 4H), 2.80 (d, 3H, $J = 4.52$ Hz), 3.12-3.17 (m, 4H), 3.76 (s, 3H), 6.48 (dd, 1H, $J = 8.56, 2.52$ Hz), 6.63 (d, 1H, $J = 2.52$ Hz), 7.05-7.10 (m, 1H), 7.27-7.35 (m, 1H), 7.40 (d, 1H, $J = 8.56$ Hz), 7.69-7.72 (m, 1H), 8.09 (s, 1H), 8.12 (s, 1H), 8.55-8.65 (m, 1H), 8.67-8.75 (m, 1H), 11.59 (s, 1H)
7-4		0.46 (MeOH: CH_2Cl_2 =1:4)	DMSO- d_6 : 2.48-2.55(m, 4H), 2.71(t, 2H), 2.80(d, 3H), 3.58-3.61(m, 4H), 3.76(s, 3H), 4.11(t, 2H), 6.52(dd, 1H), 6.66(d, 1H), 7.06(dd, 1H), 7.32(dd, 1H), 7.46(d, 1H), 7.71(dd, 1H), 8.11(s, 1H), 8.19 (s, 1H), 8.54-8.60(m, 1H), 8.60-8.75(m, 1H), 11.6(s, 1H)
7-5		m/z 497, 499 ($M+1$) ⁺	DMSO- d_6 : 1.60-1.70 (m, 2H), 1.90-1.98 (m, 2H), 2.13-2.25 (m, 2H), 2.19 (s, 3H), 2.60-2.67 (m, 2H), 2.80 (d, 3H, $J = 4.52$ Hz), 3.75 (s, 3H), 4.30-4.40 (m, 1H), 6.54 (dd, 1H, $J = 8.56, 2.0$ Hz), 6.65 (d, 1H, $J = 2.0$ Hz), 7.04-7.09 (m, 1H), 7.25-7.35 (m, 1H), 7.43 (d, 1H, $J = 8.56$ Hz), 7.68-7.73 (m, 1H), 8.10 (s, 1H), 8.18 (s, 1H) 8.52-8.59 (m, 1H), 8.68-8.75 (m, 1H), 11.57 (s, 1H)



7-6		0.25 (<i>n</i> -hexane: AcOEt=1:2)	CDCl ₃ : 2.95 (m, 4H), 3.03 (d, 3H), 3.75 (m, 4H), 3.86 (s, 3H), 6.21-6.19 (br, 1H), 6.49 (dd, 1H), 6.80 (d, 1H), 7.09-7.05 (m, 1H), 7.50 (dd, 1H), 8.08 (d, 1H), 8.13 (s, 1H), 8.68 (d, 1H), 11.07 (s, 1H)
7-7		MS m/z 510, 512 (M+1)	DMSO-d ₆ : 2.06 (s, 3H), 2.80 (d, 3H), 3.11 (t, 2H), 3.16 (t, 2H), 3.60 (dd, 4H), 3.77 (s, 3H), 6.51 (dd, 1H), 6.68 (d, 1H), 7.08 (dd, 1H), 7.33 (dd, 1H), 7.46 (d, 1H), 7.71 (d, 1H), 8.10(s, 1H), 8.12(s, 1H), 8.59-8.61 (m, 1H), 8.70-8.71(m, 1H), 11.59 (s, 1H)
7-8		0.48 (MeOH: AcOEt =5:95)	CDCl ₃ : 1.46(d, 1H), 1.68-1.82(m, 2H), 2.02-2.09(m, 2H), 2.83-2.96 (m, 2H), 3.03(d, 3H), 3.44-3.53(m, 2H), 3.82-3.92(m, 1H), 3.87(s, 3H), 6.15-6.23(m, 1H), 6.51 (d, 1H), 6.56(bs, 1H), 7.07(dd, 1H), 7.48(d, 2H), 8.08(s, 1H), 8.08-8.10(m, 1H), 8.69(d, 1H), 11.0(bs, 1H)
7-9		0.4 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 1.22 (t, 3H), 1.73-1.85 (m, 2H), 2.00-2.09 (m, 2H), 2.81-2.90 (m, 2H), 3.03 (d, 3H), 3.41-3.56 (m, 3H), 3.56 (dd, 2H), 3.58-3.62 (m, 2H), 3.64-3.68 (m, 2H), 3.86 (s, 3H), 6.15-6.24 (m, 1H), 6.50 (dd, 1H), 6.56 (d, 1H), 7.07(dd, 1H), 7.24-7.30 (m, 1H), 7.45-7.52(m, 2H), 8.08(s, 1H), 8.06-8.08 (m, 1H), 8.69 (d, 1H), 11.0 (bs, 1H)
7-10		0.4 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 1.73-1.85(m, 2H), 2.01-2.10(m, 2H), 2.82-2.90(m, 2H), 3.03(d, 3H), 3.41(s, 3H), 3.45-3.51(m, 2H), 3.56-3.58(m, 2H), 3.65-3.68(m, 2H), 3.86(s, 3H), 6.14-6.22(m, 1H), 6.50 (dd, 1H), 6.56 (d, 1H), 7.07(dd, 1H), 7.23-7.30(m, 1H), 7.44-7.52(m, 2H), 8.08(s, 1H), 8.06-8.08(m, 1H), 8.69(d, 1H), 11.0(bs, 1H)

7-11		0.54 (MeOH: CH ₂ Cl ₂ =1:4)	DMSO-d ₆ : 1.78-1.89(m, 1H), 2.13-2.22(m, 1H), 2.22(s, 6H), 2.77-2.87(m, 1H), 2.79(d, 3H), 3.04-3.10(m, 1H), 3.23-3.50(m, 3H), 3.75(s, 3H), 6.11(dd, 1H), 6.22(d, 1H), 7.05(dd, 1H), 7.21-7.32(m, 1H), 7.26(d, 1H), 7.70(d, 1H), 8.06(s, 1H), 8.08(s, 1H), 8.57-8.66(m, 1H), 8.66-8.73 (m, 1H), 11.6(s, 1H)
7-12		0.27 (MeOH: CH ₂ Cl ₂ =1:1)	DMSO-d ₆ : 1.77-1.87(m, 1H), 2.09-2.18(m, 1H), 2.35(s, 3H), 2.79(d, 1H), 3.02-3.07(m, 1H), 3.23-3.50(m, 4H), 3.74(s, 3H), 6.09(dd, 1H), 6.20(d, 1H), 7.04(dd, 1H), 7.22-7.32(m, 1H), 7.26(d, 1H), 7.70(d, 1H), 8.05(s, 1H), 8.08(s, 1H), 8.57-8.67(m, 1H), 8.67-8.73 (m, 1H), 11.6(s, 1H)
7-13		0.23 (MeOH: AcOEt =5:95)	CDCl ₃ : 1.62-1.74(m, 3H), 1.76-1.85(m, 2H), 2.00-2.09(m, 2H), 2.20-2.31(m, 1H), 2.64-2.69 (m, 2H), 2.79 (d, 3H), 3.56-4.04(m, 2H), 4.04(s, 3H), 6.49(dd, 1H), 6.63(d, 1H), 6.78(bs, 1H), 7.07 (dd, 1H), 7.28-7.38 (m, 1H), 7.39(d, 1H), 7.71(d, 1H), 8.09-8.11(m, 2H), 8.09(s, 1H), 8.60(d, 1H), 8.71(d, 1H), 11.6(bs, 1H)
7-14		0.30 (MeOH: CH ₂ Cl ₂ =4:1)	DMSO-d ₆ : 1.61-1.46(m, 2H), 1.92-1.82 (m, 2H), 2.14 (s, 3H), 2.41-2.23 (m, 5H), 2.60-2.45 (m, 4H), 2.67 (t, 2H), 2.79 (d, 3H), 3.75 (s, 3H), 3.71-3.75 (m, 2H), 6.48 (dd, 1H), 6.63 (d, 1H), 7.10-7.03 (m, 1H), 7.34-7.27 (m, 1H), 7.43-7.35 (m, 1H), 7.71(dd, 1H), 8.09 (s, 1H), 8.11 (bs, 1H), 8.65-8.56 (m, 1H), 8.75-8.67 (m, 1H), 11.6 (s, 1H)


7-15		MS (ESI) m/z 524, 526 (M+1) ⁺	DMSO-d ₆ : 2.19-2.37 (m, 4H), 2.65-2.85 (m, 3H), 2.80 (d, 3H, J = 4.5 Hz), 3.15-3.21 (m, 1H), 3.48-3.59 (m, 2H), 3.61-3.67 (m, 1H), 3.72-3.81 (m, 1H), 3.76 (s, 3H), 6.47 (dd, 1H, J = 8.6, 2.5 Hz), 6.65 (d, 1H, J = 2.5 Hz), 7.04-7.10 (m, 1H), 7.28-7.35 (m, 1H), 7.42 (d, 1H, J = 8.6 Hz), 7.69-7.74 (m, 1H), 8.09 (s, 1H), 8.12 (s, 1H), 8.55-8.63 (m, 1H), 8.68-8.73 (m, 1H), 11.60 (s, 1H)
7-16		MS (ESI) m/z 524, 526 (M+1) ⁺	DMSO-d ₆ : 2.19-2.37 (m, 4H), 2.65-2.85 (m, 3H), 2.80 (d, 3H, J = 4.5 Hz), 3.15-3.21 (m, 1H), 3.48-3.59 (m, 2H), 3.61-3.67 (m, 1H), 3.72-3.81 (m, 1H), 3.76 (s, 3H), 6.47 (dd, 1H, J = 8.6, 2.5 Hz), 6.65 (d, 1H, J = 2.5 Hz), 7.04-7.10 (m, 1H), 7.28-7.35 (m, 1H), 7.42 (d, 1H, J = 8.6 Hz), 7.69-7.74 (m, 1H), 8.09 (s, 1H), 8.12 (s, 1H), 8.55-8.63 (m, 1H), 8.68-8.73 (m, 1H), 11.60 (s, 1H)
7-17		MS 510	DMSO-d ₆ : 0.98 (t, 3H), 1.81-1.71 (m, 3H), 1.95-1.84 (m, 3H), 2.68-2.63 (m, 1H), 2.80 (d, 3H), 3.12-3.08 (m, 4H), 3.28 (d, 2H), 3.76 (s, 3H), 6.50 (dd, 1H), 6.64 (d, 1H), 6.86 (bs, 1H), 7.07 (dd, 1H), 7.46-7.19 (m, 3H), 7.71 (d, 1H), 8.09 (s, 1H), 8.15-8.10 (m, 1H), 8.66-8.58 (m, 1H), 8.77-8.70 (m, 1H), 11.6 (s, 1H)
7-18		MS 510	DMSO-d ₆ : 0.98 (t, 3H), 1.81-1.71 (m, 3H), 1.95-1.84 (m, 3H), 2.68-2.63 (m, 1H), 2.80 (d, 3H), 3.12-3.08 (m, 4H), 3.28 (d, 2H), 3.76 (s, 3H), 6.50 (dd, 1H), 6.64 (d, 1H), 6.86 (bs, 1H), 7.07 (dd, 1H), 7.46-7.19 (m, 3H), 7.71 (d, 1H), 8.09 (s, 1H), 8.15-8.10 (m, 1H), 8.66-8.58 (m, 1H), 8.77-8.70 (m, 1H), 11.6 (s, 1H)


7-19		0.16 (CH ₂ Cl ₂ :MeOH=9:1)	1.40-1.53 (m, 2H), 1.72-1.80 (m, 2H), 2.18 (s, 3H), 2.19-2.44 (m, 5H), 2.80 (d, 3H), 3.46 (m, 2H), 3.74 (s, 3H), 6.65 (dd, 1H), 6.91 (d, 1H), 7.07-7.10 (m, 1H), 7.36-7.40 (m, 1H), 7.45-7.49 (m, 1H), 7.73 (dd, 1H), 8.12 (s, 1H), 8.18 (s, 1H), 8.61 (d, 1H), 8.72-8.77 (m, 1H), 11.68 (s, 1H)
7-20		Ms : 511	1.25-1.37 (m, 2H), 1.62-1.79 (m, 3H), 1.81-1.9 (m, 2H), 2.16 (s, 3H), 2.75-2.85 (m, 5H), 3.76 (s, 3H), 3.8-3.88 (m, 2H), 6.45-6.55 (m, 1H), 6.6-6.67 (m, 1H), 7.02-7.12 (m, 1H), 7.25-7.35 (m, 1H), 7.4-7.5 (m, 1H), 7.67-7.78 (m, 1H), 8.1 (s, 1H), 8.19 (brs, 1H) 8.5-8.62 (m, 1H), 8.66-8.8 (m, 1H), 11.6 (s, 1H)
7-21		Ms : 526	2.17 (s, 3H), 2.29-2.39 (m, 3H), 2.45-2.56 (m, 4H), 2.7 (t, 2H), 3.76 (s, 3H), 4.09 (t, 2H), 6.52 (dd, 1H), 6.66 (d, 1H), 7.06 (dd, 1H), 7.31 (dd, 1H), 7.45 (d, 1H), 7.71 (dd, 1H), 8.1 (s, 1H), 8.19 (s, 1H), 8.5-8.6 (m, 1H), 8.67-8.75 (m, 1H), 11.6 (s, 1H)
7-22		Ms : 482	2.24 (s, 3H), 2.42-2.5 (m, 4H), 2.8 (d, 3H), 2.94-3.0 (m, 4H), 3.74 (s, 3H), 6.65 (dd, 1H), 6.93 (d, 1H), 7.07-7.14 (m, 1H), 7.34-7.4 (m, 1H), 7.45 (d, 1H), 7.73 (dd, 1H), 8.14 (s, 1H), 8.18 (s, 1H), 8.61 (dd, 1H), 8.7-8.77 (m, 1H), 11.7 (s, 1H)
7-23		Ms : 482	1.67-1.76 (m, 1H), 2.0-2.1 (m, 1H), 2.25-2.31 (m, 3H), 2.8 (d, 3H), 2.85-2.91 (m, 1H), 3.04-3.12 (m, 1H), 3.14-3.3 (m, 3H), 3.7 (s, 3H), 6.26 (dd, 1H), 6.91 (d, 1H), 7.01-7.04 (m, 1H), 7.07 (dd, 1H), 7.32 (dd, 1H), 7.72 (d, 1H), 8.14 (s, 1H), 8.17 (s, 1H), 8.63 (d, 1H), 8.7-8.78 (m, 1H), 11.6 (s, 1H)

7-24		Ms : 550	1.35-1.57 (m, 8H), 1.7-1.78 (m, 2H), 2.81 (d, 3H), 3.46-3.52 (m, 2H), 3.74 (s, 3H), 6.65 (dd, 1H), 6.91 (d, 1H), 7.05-7.12 (m, 1H), 7.34-7.42 (m, 1H), 7.46 (d, 1H), 7.73 (dd, 1H), 8.11 (s, 1H), 8.18 (s, 1H), 8.62 (dd, 1H), 8.71-8.78 (m, 1H), 11.7 (s, 1H)
7-25		536 [M+1] ⁺	DMSO-d ₆ : 1.48-1.58(m, 2H), 1.65-1.72(m, 4H), 1.90-1.97(m, 2H), 2.07-2.14(m, 1H), 2.49-2.55(m, 4H), 2.70-2.77(m, 2H), 2.79(d, 3H), 3.60-3.65(m, 2H), 3.75(s, 3H), 6.48(dd, 1H), 6.63(d, 1H), 7.03-7.09(m, 1H), 7.28-7.34(m, 1H), 7.39(d, 1H), 7.71(dd, 1H), 8.09(s, 1H), 8.11(s, 1H), 8.55-8.65(m, 1H), 8.69-8.73(m, 1H), 11.59(s, 1H)
7-26		468 [M+1] ⁺	DMSO-d ₆ : 2.80(d, 3H), 2.84-2.89(m, 4H), 3.04-3.08(m, 4H), 3.76(s, 3H), 6.47(dd, 1H), 6.62(dd, 1H), 7.04-7.10(m, 1H), 7.28-7.35(m, 1H), 7.40(d, 1H), 7.69-7.73(m, 1H), 8.09(s, 1H), 8.12(s, 1H), 8.55-8.63(m, 1H), 8.68-8.73(m, 1H), 11.59(s, 1H) (an aliphatic NH is hidden)
7-27		393 [M+1] ⁺	DMSO-d ₆ : 2.80(d, 3H), 6.64-6.67(m, 1H), 7.01-7.08(m, 2H), 7.15(d, 1H), 7.24-7.29(m, 2H), 7.44(d, 1H), 7.69-7.73(m, 1H), 8.20(s, 1H), 8.65-8.73(m, 2H), 9.15(s, 1H), 11.06(s, 1H), 11.63(s, 1H)
7-28		407 [M+1] ⁺	DMSO-d ₆ : 2.81(d, 3H), 3.79(s, 3H), 6.67(d, 1H), 7.05-7.10(m, 1H), 7.12(d, 1H), 7.17(d, 1H), 7.23(d, 1H), 7.25-7.30(m, 1H), 7.50(d, 1H), 7.70-7.73(m, 1H), 8.20(s, 1H), 8.67(d, 1H), 8.70-8.75(m, 1H), 9.17(s, 1H), 11.64(s, 1H)

7-29		492 [M+1] ⁺	DMSO-d ₆ : 2.80(d, 3H), 2.91-2.99(m, 4H), 3.65-3.81(m, 2H), 3.82-3.95(m, 2H), 4.12(s, 3H), 6.58(d, 1H), 6.90(d, 1H), 7.05-7.09(m, 1H), 7.14(d, 1H), 7.22-7.28(m, 1H), 7.30(d, 1H), 7.70(dd, 1H), 8.16(s, 1H), 8.63-8.67(m, 1H), 8.68-8.72(m, 1H), 9.06(s, 1H), 11.64(s, 1H)
7-30		MS m/z 510	DMSO-d ₆ : 2.02 (s, 3H), 2.80 (d, 3H), 2.82-2.92 (m, 2H), 2.92-3.01 (m, 2H), 3.44-3.53 (m, 4H), 3.76 (s, 3H), 6.68 (dd, 1H), 6.95 (d, 1H), 7.09 (dd, 1H), 7.35-7.40 (m, 1H), 7.50 (brs, 1H), 7.73 (d, 1H), 8.15 (s, 1H), 8.19 (s, 1H), 8.59 (d, 1H), 8.69-8.76 (m, 1H), 11.66 (s, 1H).


The following 2-[5-Bromo-2-(substituted phenylamino)-pyrimidin-4-ylamino]-N-ethyl-benzamide are prepared from 2-(5-bromo-2-chloro-pyrimidin-4-ylamino)-N-ethyl-benzamide and the corresponding aniline following the procedure of Example 7A







Exptl No.	Rx	R _f (solvent) or MS	NMR
8-1		0.27 (<i>n</i> -hexane: AcOEt=1:2)	DMSO-d ₆ : 2.80(d, 3H), 2.88(t, 4H), 3.65 (m, 4H), 3.75 (s, 3H), 6.64 (dd, 1H), 6.94 (d, 1H), 7.11-7.08 (m, 1H), 7.38-7.34 (m, 1H), 7.47-7.46 (m, 1H), 7.70 (dd, 1H), 8.11 (s, 1H), 8.26 (s, 1H), 8.51-8.49 (m, 1H), 8.72-8.71 (m, 1H), 11.41 (s, 1H)



8-2		m/z 513, 515 (M+1)	DMSO-d6: 2.79 (d, 3H, J = 4.04 Hz), 3.10-3.20 (m, 4H), 3.77 (s, 3H), 3.70-3.80 (m, 4H), 6.45-6.55 (m, 1H), 6.63-6.69 (m, 1H), 7.05-7.10 (m, 1H), 7.28-7.34 (m, 1H), 7.40-7.45 (m, 1H), 7.65-7.70 (m, 1H), 8.13 (s, 1H), 8.16 (s, 1H), 8.50-8.56 (m, 1H) 8.65-8.72 (m, 1H), 11.40 (s, 1H)
8-3		0.48 (n-Hexane: AcOEt=4:1)	DMSO-d6: 2.80(d, 3H), 3.83(s, 3H), 4.11(t, 2H), 6.82(ddd, 1H), 7.03(dd, 1H), 7.15(dd, 1H), 7.44(dd, 1H), 7.73(d, 1H), 7.93(dd, 1H), 8.13(s, 1H), 8.33 (s, 1H), 8.50(d, 1H), 8.70-8.77(m, 1H), 11.3(s, 1H).
8-4		MS 446, 448	2.79 (d, 3H), 3.79 (s, 3H), 6.75 (ddd, 1H), 7.0 (dd, 1H), 7.05-7.12 (m, 1H), 7.3-7.36 (m, 1H), 7.62 (dd, 1H), 7.69 (dd, 1H), 8.2 (s, 1H), 8.29 (s, 1H), 8.45 (d, 1H), 8.66-8.73 (m, 1H), 11.4 (brs, 1H).

The following 2-[5-Chloro-2-(substituted phenylamino)-pyrimidin-4-ylamino]-N-ethyl-benzamide are prepared from 2-(2,5-Dichloro-pyrimidin-4-ylamino)-N-ethyl-benzamide and the corresponding aniline following the procedure of Example 7A




Exptl No.	Rx	R _f (solvent) or MS	NMR (400MHz) , δ (ppm)
9-1		0.35 (n-hexane: AcOEt=1:2)	CDCl ₃ : 1.27 (t, 3H), 3.10-3.15 (m, 4H), 3.47-3.58 (m, 2H), 3.85-3.93 (m, 4H), 3.89 (s, 3H), 6.08-6.17 (m, 1H), 6.48 (dd, 1H), 6.53 (d, 1H), 7.05-7.11 (m, 1H), 7.42-7.53 (m, 2H), 8.08 (s, 1H), 8.12 (d, 1H), 8.67 (d, 1H), 10.94 (brs, 1H).




9-2		MS (ESI) m/z 497, 499 (M+1) ⁺	CDCl ₃ : 1.26 (t, 3H, J = 7.56Hz), 2.37 (s, 3H), 2.57-2.62 (m, 4H), 3.15-3.20 (m, 4H), 3.49 (dq, 2H, J = 7.56, 1.52 Hz), 3.87 (s, 3H), 6.11-6.16 (m, 1H), 6.49 (dd, 1H, J = 8.56, 2.52 Hz), 6.55 (d, 1H, J = 2.52 Hz), 7.05-7.10 (m, 1H), 7.23 (s, 1H), 7.41-7.50 (m, 2H), 8.07 (s, 1H), 8.08 (d, 1H, J = 8.56Hz), 8.65-8.69 (m, 1H), 10.93 (s, 1H)
9-3		m/z 564, 566 (M+1) ⁺	DMSO-d ₆ : 1.26 (t, 3H, J = 7.56Hz), 1.40-1.50 (m, 2H), 1.56-1.64 (m, 4H), 1.67-1.82 (m, 2H), 1.88-1.97 (m, 2H), 2.33-2.44 (m, 1H), 2.52-2.57 (m, 4H), 2.63-2.73 (m, 2H), 3.51 (dq, 2H, J = 7.56, 1.52 Hz), 3.62-3.69 (m, 2H), 3.86 (s, 3H), 6.10-6.15 (m, 1H), 6.49 (dd, 1H, J = 8.56, 2.52 Hz), 6.55 (d, 1H, J = 2.52 Hz), 7.05-7.10 (m, 1H), 7.23 (s, 1H), 7.43-7.50 (m, 2H), 8.05-8.11 (m, 1H), 8.07 (s, 1H), 8.65-8.69 (m, 1H), 10.91 (s, 1H)
9-4		0.39 (MeOH: CH ₂ Cl ₂ =1:4)	DMSO-d ₆ : 1.19 (t, 3H), 1.52-1.68 (m, 2H), 1.71-1.79 (m, 4H), 1.92-2.05 (m, 2H), 2.12-2.23 (m, 1H), 2.76-2.85 (m, 2H), 3.65-3.73 (m, 2H), 3.82 (s, 3H), 6.54 (dd, 1H), 6.69 (d, 1H), 7.13 (m, 1H), 7.45 (d, 1H), 7.79 (dd, 1H), 8.15 (s, 1H), 8.15-8.18 (m, 1H), 8.60-8.68 (m, 1H), 8.74-8.83 (m, 1H).
9-5		R _f (Hexane: AcOEt = 1:2): 0.30	CDCl ₃ : 1.27 (t, 3H), 3.08-3.14 (m, 4H), 3.52 (q, 2H), 3.71-3.90 (m, 7H), 6.05-6.18 (m, 1H), 6.47 (dd, 1H), 6.53 (dd, 1H), 7.08 (dd, 1H), 7.41-7.53 (m, 2H), 8.08 (s, 1H), 8.12 (d, 1H), 8.67 (d, 1H), 10.94 (s, 1H).

9-6		R _f (AcOEt:Me OH= 4:1) 0.050	DMSO: 1.11 (t, 3H), 1.60-1.69 (m, 1H), 1.88-1.96 (m, 2H), 2.19 (s, 3H), 2.55-2.68 (m, 2H), 3.30-3.45 (m, 2H), 3.75 (s, 3H), 4.33-4.43 (m, 1H), 6.54 (dd, 1H), 6.65 (d, 1H), 7.07 (dd, 1H), 7.30 (dd, 1H), 7.43 (d, 1H), 7.71 (dd, 1H), 8.11 (s, 1H), 8.20 (s, 1H), 8.54 (br.d, 1H), 8.75 (dd, 1H), 11.49 (s, 1H).
9-7		0.44 (CH ₂ Cl ₂ :M eOH=8:2)	CDCl ₃ : 1.34 (t, 3H), 1.62-1.68 (m, 2H), 1.93-2.18 (m, 8H), 2.37-2.40 (br, 2H), 2.74-2.86 (br, 3H), 3.20-3.23 (m, 2H), 3.34 (br, 2H), 3.53 (q, 2H), 3.85 (s, 3H), 6.47 (dd, 1H), 6.76 (d, 1H), 7.04-7.08 (m, 1H), 7.30 (dd, 1H), 7.53 (s, 1H), 8.00 (d, 1H), 8.13-8.17 (m, 1H), 8.22 (d, 1H), 8.42-8.53 (br, 1H), 10.91 (s, 1H), 11.59-11.75 (br, 1H)


The following 2-[5-Chloro-2-(substituted phenylamino)-pyrimidin-4-ylamino]-6,N-dimethylbenzamide are prepared from 2-(2,5-Dichloro-pyrimidin-4-ylamino)-6,N-dimethyl-benzamide and the corresponding aniline following the procedure of Example 7A







Exptl No.	Rx	Identification
--------------	----	----------------

10-1		NMR (400MHz, DMSO-d ₆ , δ): 1.58-1.68(m, 2H), 1.87-1.96(m, 2H), 2.13-2.22(m, 2H), 2.18(s, 3H), 2.18(s, 3H), 2.29(s, 3H), 2.57-2.65(m, 2H), 2.76(d, 3H), 3.75(s, 3H), 4.29-4.37(m, 1H), 6.45(dd, 1H), 6.61(d, 1H), 6.98(d, 1H), 7.18(dd, 1H), 7.47(d, 1H), 7.89(d, 1H), 8.02(s, 1H), 8.07 (s, 1H), 8.37-8.43(m, 1H), 8.49(s, 1H). Rf: 0.39 (MeOH: CH ₂ Cl ₂ =1:4).
10-2		NMR (400MHz, DMSO-d ₆ , δ): 1.35-1.42 (m, 2H), 1.45-1.60 (m, 6H), 1.75-1.85 (m, 2H), 2.29 (s, 3H), 2.30-2.35 (m, 1H), 2.43-2.50 (m, 4H), 2.57-2.66 (m, 2H), 2.76 (d, 3H, J = 5.0Hz), 3.65-3.74 (m, 2H), 3.76 (s, 3H), 6.40 (dd, 1H, J = 9.0, 2.0 Hz), 6.59 (d, 1H, J = 2.0 Hz), 6.98 (d, 1H, J = 7.6 Hz), 7.20 (dd, 1H, J = 7.6, 7.6 Hz), 7.43 (d, 1H, J = 9.0 Hz), 7.91-7.94 (m, 1H), 7.93 (s, 1H), 8.06 (s, 1H), 8.36-8.42 (m, 1H) 8.47 (s, 1H). MS (ESI) <i>m/z</i> 564, 566 (M+1) ⁺
10-3		DMSO-d ₆ : 2.29(s, 3H), 2.77(d, 3H), 3.07-3.11(m, 4H), 3.73-3.76(m, 4H), 3.77(s, 3H), 6.41(dd, 1H), 6.63(d, 1H), 7.00 (d, 1H), 7.21(dd, 1H), 7.49(d, 1H), 7.93(d, 1H), 7.96(s, 1H), 8.07(s, 1H), 8.37-8.42(m, 1H), 8.49(s, 1H). MS <i>m/z</i> 483 [M+1] ⁺

The following 2-[5-Chloro-2-(substituted phenylamino)-pyrimidin-4-ylamino]-5-fluoro-N-methylbenzamide are prepared from 2-(2,5-Dichloro-pyrimidin-4-ylamino)-5-fluoro-N-methylbenzamide and the corresponding aniline following the procedure of Example 7A



Expl No.	Rx	Identification
----------	----	----------------

11-1		NMR (400MHz, DMSO-d ₆ , δ): 2.79(d, 3H), 3.10-3.15(m, 4H), 3.74-3.78(m, 7H), 6.50(dd, 1H), 6.66(d, 1H), 7.13-7.20 (m, 1H), 7.41(d, 1H), 7.57(dd, 1H), 8.09(s, 1H), 8.14(s, 1H), 8.55-8.65(m, 1H), 8.75-8.82(m, 1H), 11.39(s, 1H). MS (ESI): m/z 487, 489 (M+1).
11-2		NMR (400MHz, DMSO-d ₆ , δ): 1.68-1.33 (m, 8H), 1.93-1.73 (m, 2H), 2.35-2.60 (m, 1H), 2.62-2.74 (m, 2H), 2.67 (t, 2H), 2.74 (d, 3H), 3.25-3.38 (m, 4H), 3.76 (s, 3H), 3.83-3.71 (m, 2H), 6.48 (dd, 1H), 6.49 (dd, 1H), 6.63 (d, 1H), 7.15 (dd, 1H), 7.36 (d, 1H), 7.57 (dd, 1H), 8.09(s, 1H), 8.12(s, 1H), 8.65-8.55 (m, 1H), 8.78(d, 1H), 11.39 (s, 1H) MS (ESI): m/z 568, 570 (M+1)
11-3		DMSO-d ₆ : 2.80(d, 3H), 3.79(s, 3H), 6.64(d,1H), 7.05-7.20(m, 3H), 7.23(d, 1H), 7.42-7.49(d, 1H), 7.57(dd, 1H), 8.20(s, 1H), 8.62-8.69(m, 1H), 8.75-8.82(m, 1H), 9.17(s, 1H), 11.43(s, 1H). MS m/z 425 [M+1] ⁺
11-4		DMSO-d ₆ : 2.06(s, 3H), 2.79(d, 3H), 3.10-3.14(m, 2H), 3.15-3.19(m, 2H), 3.55-3.62(m, 4H), 3.77(s, 3H), 6.52(dd, 1H), 6.69(d, 1H), 7.15-7.23(m, 1H), 7.43(d, 1H), 7.58(dd, 1H), 8.10(s, 1H), 8.14(s, 1H), 8.56-8.65(m, 1H), 8.75-8.81(m, 1H), 11.39(s, 1H). MS m/z 528 [M+1] ⁺


12-1 Preparation of 7-[5-Chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-2-methyl-2,3-dihydro-isoindol-1-one

Synthetic procedure for 7-(2,5-Dichloro-pyrimidin-4-ylamino)-2-methyl-2,3-dihydro-isoindol-1-one

N-Methyl-7-nirto-2,3-dihydroisoindole-1-one. At room temperature, a solution of methyl 2-bromomethyl-6-nitrobenzoate (1.26 g, 4.63 mmol) in THF (13 mL) is treated with 2M soln. of methylamine in THF (14 mL), stirred for 5 h, diluted with EtOAc (100 mL), washed with sat. aqueous solution of NaHCO₃ (15 mL) and brine (15 mL), dried (MgSO₄), and evaporated. A flash chromatography (30 g of silica gel; CH₂Cl₂/EtOAc 1:1) gives *N-Methyl-7-nirto-2,3-dihydroisoindole-1-one* (0.561 g, 2.92 mmol) in 63%. Yellow solid. *R_f* (CH₂Cl₂/EtOAc 1:1) 0.46. ¹H-NMR (400 MHz, CDCl₃) 3.21 (s), 4.44 (s), 7.63 – 7.69 (m, 2 H), 7.70 – 7.75 (m, 1 H).


7-Amino-N-methyl-2,3-dihydroisoindole-1-one. At room temperature, a solution of *N-Methyl-7-nirto-2,3-dihydroisoindole-1-one* (561.0 mg, 2.92 mmol) in EtOAc (8.4 mL) is treated with SnCl₂·2H₂O (2.68 g), stirred at 80°C under reflux for 5 h, and treated with 30 mL of 5N NaOH at 0°C. After the both layers are separated, the aqueous layer is extracted with EtOAc (2 x 8 mL), the combined extracts are washed with brine (5 mL), dried (MgSO₄), and evaporated to give *7-Amino-N-methyl-2,3-dihydroisoindole-1-one* (455.9 g, 2.81 mmol) in 96%. Yellow solid. *R_f* (CH₂Cl₂/EtOAc 1:1) 0.53. ¹H-NMR (400 MHz, CDCl₃) 3.12 (s), 4.28 (s), 5.20 (br. s), 6.56 (d, *J* = 8.0), 6.68 (d, *J* = 8.0), 7.21 (dd, *J* = 8.0, 8.0).

7-(4-Amino-2,5-dichloropyrimidin-4-yl)amino-N-methyl-2,3-dihydroisoindole-1-one. At 0°C, a solution of *7-Amino-N-methyl-2,3-dihydroisoindole-1-one* (232.6 mg, 1.43 mmol) in DMF (2.0 mL) is treated with 60% NaH (89.8 mg), stirred at the same temperature for 1.5 h, treated with a solution of 2,4,5-trichloropyrimidine (0.557 g) in DMF (3.5 mL), stirred for 1 h, and warmed to room temperature. After furthermore stirring for 13 h, the mixture is treated with sat. aqueous NH₄Cl (6 mL), and the resulting brown precipitates are collected by a filtration, followed by washing with H₂O, hexane, and CH₃CN to give *7-(4-Amino-2,5-dichloropyrimidin-4-yl)amino-N-methyl-2,3-dihydroisoindole-1-one* (130.2 g, 0.416 mmol) in 26%. Brown solid. *R_f* (CH₂Cl₂/EtOAc 1:1) 0.50. ¹H-NMR (400 MHz, CDCl₃): 3.22 (s), 4.43 (s), 7.15 (d, *J* = 8.0), 7.59 (dd, *J* = 8.0, 8.0), 8.24 (s), 8.71 (d, *J* = 8.0), 11.05 (br. s).




The following 7-[5-Chloro-2-(substituted phenylamino)-pyrimidin-4-ylamino]-2-methyl-2,3-dihydro-isoindol-1-one are prepared from 7-(2,5-Dichloro-pyrimidin-4-ylamino)-2-methyl-2,3-dihydro-isoindol-1-one and the corresponding aniline following the procedure of Example 7A.





7-[5-Chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-2-methyl-2,3-dihydro-isoindol-1-one




$^1\text{H-NMR}$ (400MHz, DMSO- d_6 , δ): 3.07 (s, 3H), 3.13-3.17 (m, 4H), 3.75 (s, 3H), 3.34-3.78 (m, 4H), 4.46 (s, 2H), 6.54 (dd, 1H, $J = 8.6, 2.5$ Hz), 6.67 (d, 1H, $J = 2.5$ Hz), 7.15 (d, 1H, $J = 7.6$ Hz), 7.25-7.34 (m, 1H) 7.36 (d, 1H, $J = 8.6$ Hz), 8.13 (s, 1H), 8.36 (s, 1H), 8.37-8.50 (m, 1H) 10.57 (s, 1H). MS (ESI) m/z 481. 483 ($M+1$) $^+$


The following 7-(5-Chloro-2-(subst.phenylamino)-pyrimidin-4-ylamino)-2-methyl-2,3-dihydro-isoindol-1-ones are prepared from 7-(2,5-Dichloro-pyrimidin-4-ylamino)-2-methyl-2,3-dihydro-isoindol-1-one and the corresponding aniline following the procedure of Example 2:




Exptl No.	Rx	Mass(m/z)	NMR (400MHz) δ (ppm)
12-2		494 [M+1] ⁺	DMSO-d6: 2.24(s, 3H), 2.45-2.50(m, 4H), 3.07(s, 3H), 3.15-3.19(m, 4H), 3.74(s, 3H), 4.46(s, 2H), 6.52(dd, 1H), 6.66(d, 1H), 7.15 (d, 1H), 7.25-7.36(m, 2H), 8.12(s, 1H), 8.35(s, 1H), 8.35-8.45(m, 1H), 10.57(s, 1H)
12-3		495 [M+1] ⁺	DMSO-d6: 1.48-1.57(m, 2H), 1.83-1.88(m, 2H), 2.83-2.90(m, 2H), 3.07(s, 3H), 3.51-3.60(m, 2H), 3.61-3.70(m, 2H), 3.73(s, 3H), 4.46(s, 2H), 4.69(d, 1H), 6.52(dd, 1H), 6.64(d, 1H), 7.14 (d, 1H), 7.25-7.35(m, 2H), 8.12(s, 1H), 8.33(s, 1H), 8.35-8.45(m, 1H), 10.57(s, 1H)
12-4		577 [M+1] ⁺	DMSO-d6: 1.48-1.59(m, 2H), 1.83-1.88(m, 2H), 2.14(s, 3H), 2.25-2.39(m, 4H), 2.42-2.60(m, 5H), 2.66-2.73(m, 2H), 3.07(s, 3H), 3.73-3.77(m, 2H), 3.74(s, 3H), 4.46(s, 2H), 6.52(dd, 1H), 6.64(d, 1H), 7.14 (d, 1H), 7.25-7.34(m, 2H), 8.12(s, 1H), 8.34(s, 1H), 8.35-8.45(m, 1H), 10.57(s, 1H)
12-5		562 [M+1] ⁺	DMSO-d6: 1.35-1.65(m, 8H), 1.73-1.85(m, 2H), 2.40-2.59(m, 7H), 3.08(s, 3H), 3.52-3.61(m, 2H), 3.73(s, 3H), 4.47(s, 2H), 6.72(dd, 1H), 6.94(d, 1H), 7.17(d, 1H), 7.34-7.39(m, 2H), 8.21(s, 1H), 8.37(s, 1H), 8.45-8.53(m, 1H), 10.64(s, 1H)

12-6		MS m/z 536	DMSO-d ₆ : 2.19-2.42 (m, 4H), 2.65-2.89 (m, 3H), 3.07 (s, 3H), 3.11-3.30 (m, 1H), 3.48-3.61 (m, 2H), 3.62-3.71 (m, 1H), 3.75 (s, 3H), 3.75-3.83 (m, 2H), 4.47 (s, 2H), 6.48-6.52 (m, 1H), 6.66 (d, 1H), 7.15 (d, 1H), 7.26-7.37 (m, 2H), 8.13 (s, 1H), 8.35 (s, 1H), 8.42 (brs, 1H), 10.57 (s, 1H).
------	---	------------------	--

The following 7-(5-Chloro-2-(subst.phenylamino) -pyrimidin-4-ylamino)-2-ethyl-2,3-dihydroisindol-1-ones are prepared from 7-(2,5-Dichloro-pyrimidin-4-ylamino)-2-ethyl-2,3-dihydro-isindol-1-one and the corresponding aniline following the procedure of Example 2:



Exptl No.	Rx	Mass(m/z)	NMR (400MHz) δ (ppm)
13-1		508 [M+1] ⁺	DMSO-d ₆ : 1.19(t, 3H), 2.24(s, 3H), 2.47-2.51(m, 4H), 3.15-3.21(m, 4H), 3.54(q, 2H), 3.74(s, 3H), 4.48(s, 2H), 6.54(dd, 1H), 6.65(d, 1H), 7.15 (d, 1H), 7.26-7.36(m, 2H), 8.12(s, 1H), 8.34(s, 1H), 8.37-8.48(m, 1H), 10.58(s, 1H)

Example 7B: 2-[5-Chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzamide (alternative synthesis to Example 7A)


To a suspension of 2-[5-chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-benzoic acid (5.5 g, 12.1 mmol) in 100 mL of THF are added Et₃N (2.06 mL, 14.8 mmol) and isobutyl chloroformate (1.7 mL, 12.8 mmol) at -5°C. After stirring at the same temperature for 30 min, the reaction mixture is further stirred at room temperature for 1 hour and then H₂O is added to the reaction mixture. The resulting precipitate is collected by filtration,

washed with H₂O, and dried under reduced pressure to give an intermediate (4.80 g) (10.96 mmol, 91%) as yellow solid.






NMR (400MHz, DMSO-d₆, δ): 3.10-3.20 (m, 4H), 3.70-3.80 (m, 4H), 3.93 (s, 3H), 6.53 (dd, 1H, J = 9.08, 2.0 Hz), 6.70 (d, 1H, J = 2.0 Hz), 7.49-7.54 (m, 1H), 7.67 (d, 1H, J = 8.56 Hz), 7.89 (s, 1H), 7.85-7.95 (m, 1H), 8.23 (d, 1H, J = 9.08 Hz), 8.26 (d, 1H, J = 8.56Hz), 12.60 (s, 1H).

To a 1M solution of methylamine in THF (560 μ l, 0.56 mmol) is added 82 mg of the obtained intermediate (0.187 mmol) followed by 1M solution of NaHMDS in THF (560 μ l, 0.56 mmol) dropwise. After the reaction mixture is stirred for 10 minutes, 5 mL of H₂O is added and extraction is performed with AcOEt. The organic layer is washed with brine, dried over Na₂SO₄, concentrated under reduced pressure, and purified by silica gel column chromatography (Hexane: AcOEt=1:1 to AcOEt) to give the title compound as a pale yellow solid. Data are given in Example 7A.


By repeating the procedures described above using appropriate starting materials and conditions the following compounds are obtained as identified below.




Exptl No.	Ry	Rf (solvent) or MS	NMR (400MHz) , δ (ppm)
14-1		0.10 (n-hexane: AcOEt=1:1)	CDCl ₃ : 3.02-3.19 (m, 10H), 3.83-3.91 (m, 4H), 3.87 (s, 3H), 6.45 (dd, 1H), 6.52 (d, 1H), 7.09-7.14 (m, 1H), 7.29 (m, 1H), 7.31 (dd, 1H), 7.38-7.45 (m, 1H), 8.06 (s, 1H), 8.14 (d, 1H), 8.39 (d, 1H), 8.97 (s, 1H).
14-2		0.36 (n-hexane: AcOEt=1:2)	CDCl ₃ : 1.27 (d, 6H), 3.09-3.16 (m, 4H), 3.81-3.92 (m, 4H), 3.89 (s, 3H), 4.26-4.37 (m, 1H), 5.93-5.98 (m, 1H), 6.48 (dd, 1H), 6.53 (d, 1H), 7.05-7.11 (m, 1H), 7.42-7.49 (m, 2H), 8.08 (s, 1H), 8.12 (d, 1H), 8.65 (d,

			1H), 10.88 (br.s, 1H).
14-3		505 [M+1] ⁺	DMSO-d ₆ : 2.79(d, 3H), 3.09-3.14(m, 4H), 3.74-3.77(m, 4H), 3.75(s, 3H), 6.49(dd, 1H), 6.65(d, 1H), 7.30 (d, 1H), 7.84(dd, 1H), 8.12(s, 1H), 8.40(s, 1H), 8.65-8.79(m, 2H), 11.39(s, 1H)
14-4		466 [M+1] ⁺	DMSO-d ₆ : 2.70-2.75(m, 2H), 3.04-3.09(m, 2H), 3.12-3.18(m, 4H), 3.74-3.80(m, 4H), 3.75(s, 3H), 6.54(dd, 1H), 6.67(d, 1H), 7.14 (d, 1H), 7.34(d, 1H), 7.37-7.44(m, 1H), 8.17(s, 1H), 8.35-8.50(m, 1H), 8.44(s, 1H), 10.59(s, 1H)
14-5		R _f (Hexane : AcOEt=1:2) : 0.31	DMSO: 1.18 (t, 3H), 3.11-3.21 (4, 4H), 3.30-3.60 (m, 2H), 3.71-3.85 (m, 7H), 6.50-6.58 (m, 1H), 6.71 (d, 1H), 7.17-7.26 (m, 1H), 7.46 (d, 1H), 7.64 (dd, 1H), 8.14 (s, 1H), 8.19 (s, 1H), 8.57-8.68 (m, 1H), 8.80-8.87 (m, 1H), 11.36 (s, 1H).
14-6		R _f (Hexane : AcOEt=1:1) : 0.051	DMSO: 1.71-1.92 (m, 2H), 1.92-2.06 (m, 2H), 3.08-3.14 (m, 4H), 3.48-3.57 (m, 2H), 3.63-3.75 (m, 2H), 3.84-3.90 (m, 7H), 6.47 (dd, 1H), 6.53 (d, 1H), 7.09 (ddd, 1H), 7.25-7.29 (m, 1H), 7.38-7.44 (m, 1H), 8.06 (s, 1H), 8.15 (d, 1H), 8.45 (dd, 1H), 9.60 (s, 1H).
14-7			¹ H-NMR (400MHz, δ ppm, CDCl ₃): 3.04-3.10 (m, 4H), 3.10-3.16 (m, 4H), 3.63-3.68 (m, 4H), 3.85-3.90 (m, 7H), 6.46 (dd, 1H), 6.53 (d, 1H), 7.20-7.25 (m, 1H), 7.33 (brs, 1H), 7.56-7.62 (m, 1H), 7.85 (dd, 1H), 8.03 (d, 1H), 8.12 (s, 1H), 8.57-8.61 (m, 1H), 9.30 (s, 1H).


The following 2-(5-Chloro-2-(subst. phenylamino)-pyrimidin-4-ylamino)-N-methyl-5-pyrrolidin-1-yl-benzamides are prepared from 2-(5-Chloro-2-methyl-pyrimidin-4-ylamino)-N-methyl-5-pyrrolidin-1-yl-benzamide and the corresponding aniline following the procedure of Example 2:







Expl No.	Rx	Mass(m/z)	NMR (400MHz) δ (ppm)
15-1		551 [M+1] ⁺	DMSO-d6: 1.94-1.99(m, 4H), 2.23(s, 3H), 2.43-2.48(m, 4H), 2.78(d, 3H), 3.11-3.17(m, 4H), 3.22-3.29(m, 4H), 3.76(s, 3H), 6.46(dd, 1H), 6.48-6.53(m, 1H), 6.63(d, 1H), 6.79(d, 1H), 7.44(d, 1H), 7.89(s, 1H), 7.99(s, 1H), 8.24(d, 1H), 8.60(d, 1H), 10.88(s, 1H)
15-2		566 [M+1] ⁺	DMSO-d6: 1.60-1.70(m, 2H), 1.90-2.00(m, 6H), 2.12-2.20(m, 2H), 2.18(s, 3H), 2.60-2.65(m, 2H), 2.78(d, 3H), 3.22-3.28(m, 4H), 3.75(s, 3H), 4.25-4.37(m, 1H), 6.49-6.55(m, 2H), 6.62(d, 1H), 6.80(d, 1H), 7.53(d, 1H), 7.90(s, 1H), 8.00(s, 1H), 8.24(d, 1H), 8.58-8.63(m, 1H), 10.88(s, 1H)

15-3		538 [M+1] ⁺	DMSO-d ₆ : 1.94-1.99(m, 4H), 2.78(d, 3H), 3.09-3.15(m, 4H), 3.22-3.27(m, 4H), 3.73-3.77(m, 4H), 3.76(s, 3H), 6.47(dd, 1H), 6.47-6.53(m, 1H), 6.65(d, 1H), 6.79(d, 1H), 7.47(d, 1H), 7.90(s, 1H), 7.99(s, 1H), 8.24(d, 1H), 8.60(d, 1H), 10.88(s, 1H)
------	---	---------------------------	---


The following 2-[5-Chloro-2-(4-fluoro-2-methoxy-phenylamino)-pyrimidin-4-ylamino]-5-subst.-N-methyl-benzamide are prepared from the corresponding aniline following the procedure of Example 2:



Exptl No.	Ry	Mass(m/z)	NMR (400MHz) δ (ppm)
16-1		487 [M+1] ⁺	DMSO-d ₆ : 2.79(d, 3H), 3.11-3.15(m, 4H), 3.74-3.81(m, 4H), 3.81(s, 3H), 6.76(ddd, 1H), 6.95-7.05(m, 2H), 7.21(d, 1H), 7.72(dd, 1H), 8.08(s, 1H), 8.09(s, 1H), 8.33(d, 1H), 8.63-8.73(m, 1H), 11.17(s, 1H)
16-2		500 [M+1] ⁺	DMSO-d ₆ : 2.24(s, 3H), 2.45-2.52(m, 4H), 2.79(d, 3H), 3.13-3.18(m, 4H), 3.81(s, 3H), 6.75(ddd, 1H), 6.94-7.02(m, 2H), 7.20(d, 1H), 7.73(dd, 1H), 8.03-8.11(m, 2H), 8.30(d, 1H), 8.60-8.70(m, 1H), 11.14(s, 1H)
16-3		432 [M+1] ⁺	DMSO-d ₆ : 2.79(d, 3H), 3.80-3.81(m, 6H), 6.75(ddd, 1H), 6.90-7.02(m, 2H), 7.27(d, 1H), 7.67(dd, 1H), 8.10(s, 1H), 8.16(s, 1H), 8.39(d, 1H), 8.70-8.76(m, 1H), 11.20(s, 1H)


16-4		568 [M+1] ⁺	DMSO-d ₆ : 1.35-1.62(m, 8H), 1.78-1.85(m, 2H), 2.30-2.40(m, 1H), 2.41-2.52(m, 4H), 2.60-2.70(m, 2H), 2.78(d, 3H), 3.70-3.80(m, 2H), 3.81(s, 3H), 6.75(ddd, 1H), 6.95-7.02(m, 2H), 7.20(d, 1H), 7.72(dd, 1H), 8.05-8.08(m, 2H), 8.28(d, 1H), 8.63-8.69(m, 1H), 11.12(s, 1H)
------	---	---------------------------	---

Example 16B




CDCl₃: 3.01-3.10 (m, 4H), 3.63-3.68 (m, 4H), 3.89 (s, 3H), 6.59 (ddd, 1H), 6.66 (dd, 1H), 7.20-7.26 (m, 1H), 7.36 (s, 1H), 7.57-7.63 (m, 1H), 7.84 (dd, 1H), 8.09-8.14 (m, 1H), 8.14 (s, 1H), 8.53 (d, 1H), 9.30 (s, 1H).

Example 16C




CDCl₃: 3.56-3.65 (m, 2H), 3.88 (s, 3H), 5.11-5.19 (m, 1H), 6.50-6.56 (m, 1H), 6.61-6.66 (m, 1H), 7.25-7.29 (m, 1H), 7.38 (brs, 1H), 7.58-7.62 (m, 1H), 7.97 (dd, 1H), 8.02-8.10 (m, 1H), 8.15 (s, 1H), 8.41 (dd, 1H), 8.81 (s, 1H).

The following 2-(5-Chloro-2-(subst.phenylamino)-pyrimidin-4-ylamino)-5-fluoro-N-methylbenzamide are prepared from 2-(2,5-Dichloro-pyrimidin-4-ylamino)-5-fluoro-N-methylbenzamide and the corresponding aniline following the procedure of Example 2:














Exptl No.	Rx	Mass(m/z)	NMR (400MHz) δ (ppm)
18-1		595 [M+1] ⁺	DMSO-d ₆ : 2.06 (s, 3H), 2.78 (d, 3H), 3.05-3.18 (m, 8H), 3.53-3.64 (m, 4H), 3.68-3.77 (m, 4H), 3.77 (s, 3H), 6.51 (dd, 1H), 6.69 (d, 1H), 6.88 (br.d, 1H), 7.20 (d, 1H), 7.43 (d, 1H), 7.99-8.03 (m, 2H), 8.34 (br.d, 1H), 8.63-8.71 (m, 1H), 11.15 (s, 1H).






The following 2-[5-chloro-2-(subst.phenylamino)-pyrimidin-4-ylamino]-N-isopropyl-benzene-sulfonamides are prepared from 2-(5-chloro-2-chloro-pyrimidin-4-ylamino)-N-isopropyl-benzenesulfonamide and the corresponding aniline following the procedure of Example 7A













ExptlNo.	Rx	R _f (solvent) Or MS	NMR (400MHz), δ (ppm)
----------	----	-----------------------------------	------------------------------





19-1		0.39 (MeOH: CH ₂ Cl ₂ =1:4)	DMSO-d ₆ : 0.94(d, 6H), 1.75-1.84(m, 1H), 2.07-2.16(m, 1H), 2.33(s, 3H), 2.98-3.04(m, 1H), 3.22-3.36(m, 5H), 3.42-3.47(m, 1H), 3.74(s, 3H), 6.05(dd, 1H), 6.18(d, 1H), 7.18(dd, 1H), 7.25(d, 1H), 7.35-7.45(m, 1H), 7.77-7.82(m, 1H), 7.70-8.10(m, 1H), 8.09-8.17 (m, 2H), 8.45-8.63(m, 1H), 9.34(s, 1H)
19-2		0.40 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 1.00(d, 6H), 1.13(t, 3H), 1.83-1.92(m, 1H), 2.23-2.30(m, 1H), 2.70-2.78(m, 2H), 3.08-3.13(m, 1H), 3.27-3.54 (m, 5H), 3.85(s, 3H), 4.33(d, 1H), 6.05(d, 1H), 6.13(s, 1H), 7.13(bs, 1H), 7.18-7.22(m, 1H), 7.52-7.56(m, 1H), 7.83-7.86(m, 1H), 7.95-7.98(m, 1H), 8.09(s, 1H), 8.47-8.49(m, 1H), 8.89(s, 1H)
19-3		0.30 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 0.93 (d, 6H), 1.05-1.09(m, 1H), 1.48-1.99(m, 6H), 2.16(s, 3H), 2.61-2.67(m, 1H), 2.80-2.83(m, 1H), 3.75(s, 3H), 3.80-3.89(m, 2H), 6.44-6.47(m, 1H), 6.62-6.63(m, 1H), 7.18-7.22(m, 1H), 7.42-7.46(m, 1H), 7.80-7.89(m, 2H), 8.17(s, 1H), 8.23(s, 1H), 8.42-8.44(m, 1H), 8.89(s, 1H)
19-4		0.69 (MeOH: CH ₂ Cl ₂ =1:3)	DMSO-d ₆ : 0.94(d, 6H), 1.45-1.57(m, 2H), 1.80-1.88(m, 2H), 2.14(s, 3H), 2.25-2.35(m, 4H), 2.45-2.55(m, 4H), 2.62-2.70(m, 2H), 3.28-3.37(m, 1H), 3.68-3.74(m, 2H), 3.75(s, 3H), 6.44(dd, 1H, <i>J</i> =8.82, 2.0Hz), 6.61(d, 1H, <i>J</i> =2.0Hz), 7.21(dd, 1H), 7.37(d, 1H), 7.45(dd, 1H), 7.81(dd, 1H, <i>J</i> =1.82, 1.52Hz), 7.84-7.92(m, 1H), 8.12-8.20(m, 1H), 8.16(s, 1H), 8.43 - 8.51(m, 1H), 9.31(s, 1H)
19-5		0.35 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 0.93(d, 6H), 2.23(s, 3H), 2.45-2.48(m, 4H), 3.12-3.15(m, 4H), 3.75(s, 3H), 6.42-6.45(m, 1H), 6.63 (s, 1H), 7.19-7.23(m, 1H), 7.38-7.47(m, 2H), 7.80-7.89(m, 2H), 8.16(s, 1H), 8.46-8.48(m, 1H), 9.34(s, 1H)



19-6		0.45 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 0.99(d, 6H), 3.40-3.49(m, 1H), 3.88(s, 3H), 4.29-4.31(d, 1H), 6.51-6.56(m, 1H), 6.62-6.65 (m, 1H), 7.24-7.28(m, 1H), 7.37(s, 1H), 7.56-7.60(m, 1H), 7.98-8.15(m, 3H), 8.34-8.37(m, 1H), 8.89(s, 1H)
19-7		0.28 (<i>n</i> -hexane: AcOEt=1:1)	DMSO-d ₆ : 0.93(d, 6H), 1.59-1.67(m, 2H), 1.90-1.93(m, 2H), 2.10-2.24(m, 5H), 2.60-2.67(m, 2H), 3.74(s, 3H), 4.33-4.37(m, 1H), 6.47-6.50(m, 1H), 6.63(d, 1H), 7.18-7.22(m, 1H), 7.41-7.45(m, 2H), 7.79-7.87(m, 2H), 8.16(s, 1H), 8.21(s, 1H), 8.41 - 8.43(m, 1H), 9.29(s, 1H)
19-8		0.25 (<i>n</i> -hexane: AcOEt=1:1)	DMSO-d ₆ : 0.93(d, 6H), 3.09-3.12(m, 4H), 3.74-3.76(m, 7H), 6.43-6.46(m, 1H), 6.64(s, 1H), 7.19-7.23(m, 1H), 7.41-7.48(m, 2H), 7.80(d, 1H), 7.82(d, 1H), 8.17(s, 1H), 8.46-8.48(m, 1H), 9.31(s, 1H)
19-9		0.56 (MeOH: CH ₂ Cl ₂ =1:4)	DMSO-d ₆ : 0.93(d, 6H), 1.89-1.90(m, 1H), 2.30(bs, 6H), 3.13-3.50(m, 6H), 3.74(s, 3H), 6.10(d, 1H), 6.22(s, 1H), 7.16-7.20(m, 1H), 7.25-7.27(m, 1H), 7.40(bs, 1H), 7.79-7.81(m, 1H), 7.86-7.88(m, 1H), 8.12(s, 1H), 8.15(s, 1H), 8.51(s, 1H), 9.34(s, 1H)
19-10		0.45 (MeOH: CH ₂ Cl ₂ =1:4)	CDCl ₃ : 0.99(d, 12H), 2.27(s, 2H), 2.31(s, 6H), 2.96(s, 2H), 3.39-3.48(m, 1H), 3.83(s, 3H), 4.30(d, 1H), 6.09-6.12(m, 1H), 6.19(d, 1H), 7.11(s, 1H), 7.19-7.23(m, 1H), 7.51-7.57(m, 1H), 7.76-7.79(m, 1H), 7.95(d, 1H), 8.09(s, 1H), 8.46-8.49(m, 1H), 8.88(s, 1H)
19-11		0.30 (<i>n</i> -hexane: AcOEt=1:1)	DMSO-d ₆ : 0.93(d, 6H), 2.96-2.99(m, 4H), 3.74-3.76(m, 7H), 6.67-6.72(m, 1H), 7.21-7.25(m, 1H), 7.31-7.34(m, 1H), 7.44-7.48(m, 1H), 7.80-7.83(m, 1H), 7.88(d, 1H), 8.21(s, 1H), 8.42(d, 1H), 8.58(s, 1H), 9.30 (s, 1H)

19-12		0.42 (MeOH: CH ₂ Cl ₂ =1:4)	DMSO-d ₆ : 0.94(d, 6H), 1.68-1.76(m, 1H), 1.99-2.07(m, 1H), 2.29(s, 3H), 3.05-3.49(m, 6H), 3.75(s, 3H), 6.36-6.40(m, 1H), 7.10-7.37(m, 3H), 7.70-7.80(m, 1H), 8.08-8.39(m, 3H), 9.24(s, 1H)
19-13		0.50 (MeOH: CH ₂ Cl ₂ =1:4)	CDCl ₃ : 1.01(d, 6H), 1.94-1.96(m, 1H), 2.01(s, 3H), 2.29-2.37(m, 1H), 3.19-3.58(m, 5H), 3.86(s, 3H), 4.42(d, 1H), 4.59-4.63(m, 1H), 5.70(d, 1H), 6.05-6.08(m, 1H), 6.15-6.16(m, 1H), 7.17-7.24(m, 2H), 7.53-7.57(m, 1H), 7.90(d, 1H), 7.91-7.98(m, 1H), 8.09(s, 1H), 8.47(d, 1H), 8.91(s, 1H)
19-14		0.53 (MeOH: CH ₂ Cl ₂ =1:4)	CDCl ₃ : 1.00(d, 6H), 2.04(s, 3H), 2.05-2.29(m, 2H), 2.96(s, 3H), 3.19-3.54(m, 5H), 3.86(s, 3H), 4.57-4.63(m, 1H), 5.39-5.46(m, 1H), 6.07-6.09(m, 1H), 6.16(d, 1H), 7.18-7.26(m, 2H), 7.53-7.57(m, 1H), 7.89-7.98(m, 2H), 8.08(s, 1H), 8.47(d, 1H), 8.94(d, 1H)
19-15		0.56 (MeOH: CH ₂ Cl ₂ =1:4)	DMSO-d ₆ : 0.93(d, 6H), 1.48-1.56(m, 2H), 1.65-1.75(m, 4H), 1.90-1.93(m, 2H), 2.05-2.15(m, 1H), 2.45-2.55(m, 5H), 2.69-2.75(m, 2H), 3.61(d, 2H), 3.74(s, 1H), 6.42-6.51(m, 1H), 6.61(d, 1H), 7.18-7.22(m, 1H), 7.37(d, 1H), 7.43-7.47(m, 1H), 7.80(d, 1H), 7.81-7.89(m, 1H), 8.16(d, 1H), 8.46-8.48(m, 1H), 9.31(s, 1H)
19-16		0.56 (MeOH: CH ₂ Cl ₂ =1:4)	DMSO-d ₆ : 0.92(d, 6H), 1.65-1.75(m, 4H), 1.88-2.00(m, 4H), 2.39-2.43(m, 2H), 2.60-2.65(m, 2H), 3.03-3.07(m, 1H), 3.03-3.40(m, 2H), 3.70(s, 3H), 3.77-3.78(m, 1H), 6.09(d, 1H), 6.23(s, 1H), 7.13-7.17(m, 1H), 7.23-7.25(m, 1H), 7.30-7.42(m, 1H), 7.78(d, 1H), 7.86(d, 1H), 8.10(s, 1H), 8.13(s, 1H), 8.40-8.50(m, 1H), 9.31(s, 1H)


19-17		0.23 (<i>n</i> -hexane: AcOEt=1:1)	DMSO-d ₆ : 0.93(d, 6H), 1.24-1.57(m, 4H), 1.69-1.78(m, 2H), 1.98-2.04(m, 1H), 2.15-2.33(m, 5H), 2.70-2.80(m, 1H), 3.74(s, 3H), 3.91-3.94(m, 1H), 4.05-4.09(m, 1H), 6.46-6.49(m, 1H), 6.63(d, 1H), 7.18-7.22(m, 1H), 7.42-7.46(m, 2H), 7.80(d, 1H), 7.89(d, 1H), 8.17(s, 1H), 8.25(s, 1H), 8.42-8.44(m, 1H), 9.31(s, 1H)
19-18		0.48 (<i>n</i> -hexane: AcOEt=1:1)	DMSO-d ₆ : 0.93(d, 6H), 1.03(t, 3H), 1.13(t, 3H), 1.42-1.81(m, 4H), 2.57-2.83(m, 4H), 3.17-3.41(m, 4H), 3.65-3.75(m, 1H), 3.80(s, 3H), 4.21(bs, 1H), 6.42-6.47(m, 2H), 6.51(d, 1H), 6.63(d, 1H), 7.18-7.22(m, 1H), 7.38-7.47(m, 2H), 7.80-7.82(m, 1H), 7.89(d, 1H), 8.16(s, 1H), 8.47-8.49(m, 1H), 9.31(s, 1H)
19-19		0.44 (CH ₂ Cl ₂ :MeOH=9:1)	CDCl ₃ : 1.45-1.62 (m, 2H), 1.72-1.78 (m, 1H), 1.82-1.90 (m, 1H), 2.40-2.46 (m, 1H), 2.61-2.75 (m, 2H), 3.75-3.70 (m, 2H), 3.76 (s, 3H), 6.45 (dd, 1H), 6.62 (d, 1H), 6.85 (s, 1H), 7.19-7.23 (m, 1H), 7.36-7.48 (m, 3H), 7.80-7.82 (m, 1H), 7.85-7.93 (br, 1H), 8.16 (s, 2H), 8.43-8.52 (m, 1H), 9.31 (s, 1H)
19-20		Ms : 547	DMSO-d ₆ : 0.94 (d, 6H), 1.73-1.82 (m, 1H), 2.23-2.33 (m, 4H), 2.34-2.41 (m, 1H), 2.54-2.62 (m, 1H), 2.62-2.69 (m, 1H), 2.77-2.82 (m, 1H), 3.25-3.35 (m, 1H), 3.74 (s, 3H), 4.85-4.92 (m, 1H), 6.4 (dd, 1H), 6.57 (d, 1H), 7.16-7.24 (m, 1H), 7.38-7.51 (m, 1H), 7.81 (d, 1H), 7.82-7.94 (m, 1H), 8.16 (s, 1H), 8.22 (brs, 1H), 8.38-8.48 (m, 1H), 9.3 (brs, 1H)
19-21		Ms : 579	DMSO-d ₆ : 0.92 (d, 6H), 1.61-1.71 (m, 2H), 1.86-1.96 (m, 2H), 2.12-2.22 (m, 5H), 2.57-2.64 (m, 2H), 3.2-3.4 (m, 1H), 3.77 (s, 3H), 4.27-4.35(m, 1H), 6.86 (dd, 1H), 7.19-7.27 (m, 1H), 7.39-7.46 (m, 1H), 7.81 (dd, 1H), 7.84-7.92 (m, 1H), 8.21 (s, 1H), 8.36-8.42 (m, 1H), 8.62 (s, 1H), 9.28 (s, 1H)


19-22		Ms : 549	DMSO-d6 : 0.90 (s, 6H), 0.94 (d, 6H), 2.9 (d, 2H), 3.24 (d, 2H), 3.25-3.35(m, 1H), 3.27-3.36 (m, 1H), 3.68 (s, 3H), 4.58 (t, 1H), 5.3 (t, 1H), 6.16 (dd, 1H), 6.39 (d, 1H), 7.13 (d, 1H), 7.15-7.21 (m, 1H), 7.35-7.45 (m, 1H), 7.8 (dd, 1H), 7.83-7.92 (m, 1H), 8.09 (s, 1H), 8.11 (s, 1H), 8.45-8.57 (m, 1H), 9.33 (s, 1H)
19-23		Rf : 0.51 (n-hexane : AcOEt=1:1)	DMSO-d6 : 0.94 (d, 6H), 1.22 (s, 6H), 3.25-3.35 (m, 1H), 3.36 (d, 2H), 3.68 (s, 3H), 4.73-4.79 (brs, 1H), 4.81 (t, 1H), 6.29 (dd, 1H), 6.44 (d, 1H), 7.14-7.22 (m, 2H), 7.38-7.46 (m, 1H), 7.8 (dd, 1H), 7.85-7.9 (m, 1H), 8.1 (s, 1H), 8.13 (s, 1H), 8.45-8.55 (m, 1H), 9.32 (s, 1H)
19-24		Ms : 577	DMSO-d6 : 0.93 (d, 6H), 0.96 (s, 6H), 2.22 (s, 6H), 3.25-3.35 (m, 1H), 3.7 (s, 3H), 3.75 (s, 3H), 6.46 (dd, 1H), 6.62 (d, 1H), 7.16-7.23 (m, 1H), 7.38-7.47 (m, 1H), 7.81 (dd, 1H), 7.85-7.9 (m, 1H), 8.17 (s, 1H), 8.23 (s, 1H), 8.38-8.48 (m, 1H), 9.31 (s, 1H)
19-25		Ms : 521	DMSO-d6 : 0.94 (d, 6H), 3.12 (t, 4H), 3.25-3.35 (m, 1H), 3.75 (t, 4H), 6.73 (dd, 1H), 6.85 (dd, 1H), 7.16-7.24 (m, 1H), 7.25-7.32 (m, 1H), 7.38-7.47 (m, 1H), 7.8 (dd, 1H), 7.88 (d, 1H), 8.18 (s, 1H), 8.42-8.52 (m, 1H), 8.86 (s, 1H), 9.36 (s, 1H)
19-26		Ms : 565	DMSO-d6 : 0.93 (d, 6H), 2.4-2.56 (m, 4H), 2.69 (t, 2H), 3.25-3.38 (m, 1H), 3.59 (t, 4H), 4.11 (t, 1H), 6.75 (dd, 1H), 6.93 (dd, 1H), 7.16-7.23 (m, 1H), 7.3-7.4 (m, 1H), 7.4-7.38 (m, 1H), 7.8 (dd, 1H), 7.88 (d, 1H), 8.19 (s, 1H), 8.36-8.5 (m, 1H), 8.92 (s, 1H), 9.34 (s, 1H)







19-27		Ms : 614	DMSO-d6 : 0.93 (d, 6H), 1.3-1.62 (m, 8H), 1.75-1.85 (m, sH), 2.26-2.4 (m, 1H), 2.4-2.58 (m, 4H), 3.28-3.38 (m, 1H), 3.68-3.78 (m, 5H), 6.42 (dd, 1H), 6.64 (d, 1H), 7.18-7.24 (m, 1H), 7.42-7.5 (m, 2H), 7.77 (d, 1H), 7.82 (dd, 1H), 8.13 (s, 1H), 8.17 (s, 1H), 8.4-8.5 (m, 1H), 9.36 (s, 1H)
19-28		Rf : 0.5 (MeOH: CH2Cl2=3: 7)	DMSO-d6 : 0.93 (d, 6H), 1.6-1.7 (m, 2H), 1.88-1.98 (m, 2H), 2.17-2.35 (m, 5H), 2.6-2.73 (m, 2H), 3.25-3.4 (m, 1H), 4.34-4.44 (m, 1H), 6.75 (dd, 1H), 6.93 (dd, 1H), 7.16-7.23 (m, 1H), 7.29-7.36 (m, 1H), 7.37-7.47 (m, 1H), 7.8 (dd, 1H), 7.89 (d, 1H), 8.19 (s, 1H), 8.36-8.46 (m, 1H), 8.92 (s, 1H), 9.31 (s, 1H)
19-29		Ms : 577	DMSO-d6 : 0.93 (d, 6H), 2.45-2.55 (m, 4H), 2.7 (t, 2H), 3.25-3.35 (m, 1H), 3.59 (t, 3H), 3.76 (s, 3H), 4.1 (t, 1H), 6.48 (dd, 1H), 6.65 (d, 1H), 7.18-7.24 (m, 1H), 7.4-7.5 (m, 2H), 7.82 (dd, 1H), 7.88 (d, 1H), 8.17 (s, 1H), 8.24 (s, 1H), 8.4-8.48 (m, 1H), 9.31 (s, 1H)
19-30		Ms : 590	DMSO-d6 : 0.93 (d, 6H), 2.15 (s, 3H), 2.2-2.4 (m, 4H), 2.4-2.6 (m, 4H), 2.69 (t, 2H), 3.25-3.35 (m, 1H), 3.75 (s, 3H), 4.08 (t, 2H), 6.47 (dd, 1H), 6.64 (d, 1H), 7.18-7.24 (m, 1H), 7.41-7.49 (m, 2H), 7.81 (dd, 1H), 7.86-7.91 (m, 1H), 8.17 (s, 1H), 8.24 (s, 1H), 8.39-8.46 (m, 1H), 9.31 (s, 1H)






19-31		Ms : 588	DMSO-d ₆ : 0.94 (d, 6H), 2.19-2.36 (m, 4H), 2.66-2.85 (m, 3H), 3.15-3.21 (m, 1H), 3.73-3.8 (m, 5H), 6.43 (dd, 1H), 6.63 (d, 1H), 7.18-7.25 (m, 1H), 7.4 (d, 1H), 7.43-7.5 (m, 1H), 7.81 (dd, 1H), 7.89 (d, 1H), 8.16 (s, 1H), 8.17 (s, 1H), 8.42-8.52 (m, 1H), 9.32 (s, 1H)
19-32		Ms : 560	CDCl ₃ : 1.01(s, 6H), 1.45-1.56 (m, 2H), 2.03-2.11 (m, 2H), 2.11-2.2 (m, 2H), 2.31 (s, 3H), 2.78-2.87 (m, 2H), 3.22-3.31 (m, 1H), 3.39-3.5 (m, 1H), 3.82 (s, 3H), 4.5-4.6 (m, 1H), 6.13 (dd, 1H), 6.21 (d, 1H), 7.16 (s, 1H), 7.18-7.24 (m, 1H), 7.5-7.57 (m, 1H), 7.82 (d, 1H), 7.97 (dd, 1H), 8.16 (s, 1H), 8.08 (s, 1H), 8.46 (d, 1H), 8.92 (s, 1H)






The following 2-[5-chloro-2-(subst.phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzene-sulfonamides are prepared from 2-(5-chloro-2-chloro-pyrimidin-4-ylamino)-N-methyl-benzenesulfonamide and the corresponding aniline following the procedure of Example A














ExptNo.	Rx	R _f (solvent) or MS	NMR (400MHz), δ (ppm)
20-1		0.50 (AcOEt)	CDCl ₃ : 2.63(d, 3H), 3.14(t, 4H), 3.87-3.90(m, 7H), 4.64(m, 1H), 6.45(dd, 1H), 6.55(d, 1H), 7.23-7.26(m, 1H), 7.51-7.55(m, 1H), 7.91(d, 1H), 7.95(dd, 1H), 8.06(s, 1H), 8.47(d, 1H), 9.26(s, 1H)







20-2		m/z 546, 548 (M+1)	DMSO-d6: 2.06 (s, 3H), 2.43 (s, 3H), 3.10 (m, 2H), 3.16 (m, 2H), 3.59-3.62 (m, 4H), 3.77 (s, 3H), 6.49 (dd, 1H), 6.68 (d, 1H), 7.21-7.25 (m, 1H), 7.42 (d, 1H), 7.49 (dd, 1H), 7.75-7.77 (m, 1H), 7.78(s, 1H), 8.16(s, 1H), 8.21 (s, 1H), 8.50(d, 1H), 9.35 (s, 1H)
20-3		0.27 (n-hexane: AcOEt=3:1)	CDCl3: 2.65(d, 3H), 4.45-4.49(m, 1H), 6.99-7.04(m, 1H), 7.17-7.28(m, 4H), 7.56-7.60(m, 1H), 7.96-7.98(m, 1H), 8.18(s, 1H), 8.31-8.34(m, 1H), 8.41-8.44(m, 1H), 9.14(s, 1H)
20-4		0.27 (n-hexane: AcOEt=3:1)	CDCl3: 2.65(d, 3H), 4.54-4.58(m, 1H), 6.53(dd, 1H), 6.98-7.02(m, 1H), 7.11-7.15(m, 2H), 7.24-7.28(m, 1H), 7.35(bs, 1H), 7.57-7.61(m, 1H), 7.95-7.98(m, 1H), 8.16(s, 1H), 8.29-8.32(m, 1H), 8.42-8.46(m, 1H), 9.14(s, 1H)
20-5		0.46 (MeOH: CH2Cl2=1:4)	CDCl3: 1.95-2.00(m, 5H), 2.29-2.37(m, 1H), 2.62(d, 3H), 3.20-3.78(m, 4H), 3.86(s, 3H), 4.60-4.64(m, 2H), 5.68-5.69(m, 1H), 6.09-6.16(m, 2H), 7.15(bs, 1H), 7.19-7.23(m, 1H), 7.54-7.58(m, 1H), 7.88-7.95(m, 2H), 8.06(s, 1H), 8.55-8.57(m, 1H), 9.08(s, 1H)
20-6		518 [M+1]+	DMSO-d6: 2.23(s, 3H), 2.43(s, 3H), 2.45-2.50(m, 4H), 3.12-3.17(m, 4H), 3.76(s, 3H), 6.45(dd, 1H), 6.63(d, 1H), 7.22(dd, 1H), 7.37(d, 1H), 7.45-7.50(m, 1H), 7.74-7.78(m, 1H), 7.76(d, 1H), 8.15(s, 1H), 8.19(s, 1H), 8.46-8.53(m, 1H), 9.35(bs, 1H)
20-7		504 [M+1]+	DMSO-d6: 2.43(s, 3H), 2.80-2.89(m, 4H), 2.99-3.07(m, 4H), 3.76(s, 3H), 6.44(dd, 1H), 6.61(d, 1H), 7.18-7.24(m, 1H), 7.37(d, 1H), 7.44-7.50(m, 1H), 7.76(dd, 1H), 8.15(s, 1H), 8.18(s, 1H), 8.45-8.55(m, 1H), 9.20-9.45(m, 1H)


20-8		586 [M+1] ⁺	DMSO-d ₆ : 1.35-1.43(m, 2H), 1.45-1.61(m, 6H), 1.75-1.85(m, 2H), 2.30-2.40(m, 1H), 2.43(d, 3H), 2.42-2.55(m, 4H), 2.60-2.70(m, 2H), 3.68-3.77(m, 2H), 3.75(s, 3H), 6.45(dd, 1H), 6.62(d, 1H), 7.21(dd, 1H), 7.36(d, 1H), 7.43-7.51(m, 1H), 7.73-7.81(m, 1H), 7.75(dd, 1H), 8.15(s, 1H), 8.17(s, 1H), 8.45-8.52(m, 1H), 9.34(bs, 1H)
20-9		569 [M+1] ⁺	DMSO-d ₆ : 1.85-1.95(m, 2H), 2.15(s, 3H), 2.18(t, 2H), 2.22-2.40(m, 8H), 2.43(s, 3H), 4.17(t, 2H), 6.65(d, 1H), 7.06(dd, 1H), 7.20(d, 1H), 7.22(ddd, 1H), 7.25(d, 1H), 7.39-7.47(m, 2H), 7.72-7.82(m, 1H), 7.77(dd, 1H), 8.26(s, 1H), 8.52(d, 1H), 9.22(s, 1H), 9.36(s, 1H)
20-10		556 [M+1] ⁺	DMSO-d ₆ : 1.85-1.95(m, 2H), 2.19(t, 2H), 2.25-2.35(m, 4H), 2.43(s, 3H), 3.55-3.60(m, 4H), 4.19(t, 2H), 6.66(d, 1H), 7.06(dd, 1H), 7.17-7.24(m, 1H), 7.21(d, 1H), 7.27(d, 1H), 7.39-7.45(m, 1H), 7.44(d, 1H), 7.70-7.80(m, 1H), 7.76(dd, 1H), 8.26(s, 1H), 8.52(d, 1H), 9.21(s, 1H), 9.36(s, 1H)
20-11		R _f (Hexane : AcOEt=1:1) 0.29	DMSO-d ₆ : 2.64 (d, 3H), 2.87-2.96 (m, 4H), 3.65-3.74 (m, 4H), 3.86 (s, 3H), 4.41-4.51 (m, 1H), 6.50 (dd, 1H), 6.81 (d, 1H), 7.55-7.64 (m, 2H), 7.96 (d, 1H), 8.01 (s, 1H), 8.19 (s, 1H), 8.49 (d, 1H), 9.07 (s, 1H).
20-12		MS 535	DMSO-d ₆ : 2.64 (d, 3H), 3.05 (bs, 4H), 3.59 (bs, 3H), 3.87(bs, 3H), 3.89 (bs, 4H), 4.52-4.48 (m, 1H), 6.57(bs, 1H), 7.25-7.20(m, 1H), 7.44-7.32 (m, 1H), 7.63-7.52 (m, 1H), 7.94(bs, 1H), 8.06 (d, 1H), 8.25(s, 1H), 8.48(d, 1H), 9.06(bs, 1H)

20-13		MS 548	DMSO-d6: 2.17 (bs, 3H), 2.63 (d, 3H), 2.68 (bs, 4H), 3.10(bs, 4H), 3.57 (s, 3H), 4.54-4.46 (m, 1H), 6.59(bs, 1H), 7.27-7.18(m, 1H), 7.37 (bs, 1H), 7.62-7.55 (m, 1H), 7.94(bs, 1H), 7.95 (d, 1H), 8.16(s, 1H), 8.48(d, 1H), 9.04(bs, 1H)
20-14		MS 546	DMSO-d6: 1.06 (t, 3H), 1.86 (dd, 2H), 2.37 (s, 3H), 2.62-2.59 (m, 4H), 2.64(d, 3H), 4.00-3.97 (m, 4H), 4.62-4.54 (m, 1H), 6.44 (dd, 1H), 6.54(d, 1H), 7.27-7.22(m, 1H), 7.34(bs, 1H), 7.58-7.54(m, 1H), 7.95(dd, 1H), 8.02(d, 1H), 8.11(s, 1H), 8.53(d, 1H), 9.07(bs, 1H)
20-15		LC-MS 545	DMSO-d6: 1.46-1.62 (m, 2H), 1.72-1.79 (m, 1H), 1.82-1.90 (m, 1H), 2.38-2.46 (m, 1H), 2.43 (s, 3H), 2.62-2.76 (m, 2H), 3.59-3.69 (m, 2H), 3.43 (s, 3H), 6.47 (dd, 1H), 6.63 (d, 1H), 6.82-6.89 (br, 1H), 7.21 (dd, 1H), 7.32-7.41 (m, 2H), 7.44-7.52 (m, 1H), 7.71-7.82 (m, 2H), 8.15 (s, 1H), 8.15-8.20 (br, 1H), 8.44-8.53 (m, 1H), 9.28-9.38 (m, 1H)
20-16		0.24 (CH ₂ Cl ₂ :MeOH=8:2)	DMSO-d6: 1.47-1.55 (m, 2H), 1.80-1.91 (m, 2H), 2.16 (s, 3H), 2.25-2.41 (m, 5H), 2.42-2.48 (m, 3H), 2.61-2.73 (m, 2H), 3.68-3.79 (m, 5H), 6.45 (dd, 1H), 6.62 (d, 1H), 7.21 (dd, 1H), 7.34 (d, 1H), 7.45-7.49 (m, 1H), 7.73-7.80 (m, 2H), 8.15 (s, 1H), 8.20 (s, 1H), 8.45-8.54 (m, 1H), 9.34 (s, 1H)
20-17		LC-MS 518	DMSO-d6: 1.76-1.84 (m, 1H), 2.08-2.16 (m, 1H), 2.33 (s, 3H), 2.42 (s, 3H), 3.00-3.03 (m, 1H), 3.23-3.27 (m, 3H), 3.42-3.46 (m, 1H), 3.74 (s, 3H), 6.06 (dd, 1H), 6.18- 6.20 (m, 1H), 7.17-7.23 (m, 1H), 7.38-7.48 (br, 1H), 7.72-7.77 (m, 1H), 8.12 (s, 1H), 8.17-8.21 (br, 1H), 8.46-8.58 (br, 1H), 9.30-9.40 (br, 1H)


20-18		LC-MS 601	DMSO-d6: 1.36-1.49 (m, 2H), 1.69-1.76 (m, 2H), 2.13 (s, 3H), 2.15-2.23 (m, 1H), 2.24-2.36 (br, 4H), 2.39-2.48 (m, 5H), 2.43 (s, 3H), 3.27-3.40 (m, 2H), 3.74 (s, 3H), 6.62 (dd, 1H), 6.90 (d, 1H), 7.22-7.26 (m, 1H), 7.41-7.46 (m, 1H), 7.49-7.53 (m, 1H), 7.55-7.86 (br, 1H), 7.77 (dd, 1H), 8.16 (s, 1H), 8.25 (s, 1H), 8.42 (d, 1H), 9.28 (s, 1H)
20-19		LC-MS 519	DMSO-d6: 1.37-1.46 (m, 2H), 1.69-1.75 (m, 2H), 2.43 (s, 3H), 2.53-2.61 (m, 2H), 3.18-3.26 (m, 2H), 3.40-3.74 (m, 2H), 4.62 (d, 1H), 6.62 (dd, 1H), 6.90 (d, 1H), 7.22-7.26 (m, 1H), 7.42-7.46 (br, 1H), 7.48-7.55 (m, 1H), 7.77-7.80 (m, 2H), 8.13-8.18 (br, 1H), 8.25 (s, 1H), 8.40-8.45 (m, 1H), 9.25-9.30 (m, 1H)
20-20		LC-MS 532	DMSO-d6: 1.66-1.76 (m, 1H), 2.00-2.07 (m, 1H), 2.14 (s, 6H), 2.43 (s, 3H), 2.68-2.76 (m, 1H), 2.87-2.91 (m, 1H), 2.99-3.10 (m, 2H), 3.24-3.28 (m, 1H), 3.71 (s, 3H), 6.25 (dd, 1H), 6.90 (d, 1H), 7.00-7.03 (m, 1H), 7.21-7.24 (m, 1H), 7.40-7.45 (m, 1H), 7.78-7.83 (m, 2H), 8.19 (s, 1H), 8.24 (s, 1H), 8.46 (d, 1H), 9.27-9.36 (br, 1H)
20-21		Ms : 549	DMSO-d6: 2.37-2.47 (m, 4H), 2.48-2.53 (m, 3H), 2.64 (t, 2H), 3.57 (t, 3H), 3.77 (s, 3H), 3.92 (t, 2H), 6.61(dd, 1H), 6.93 (d, 1H), 7.28 (dd, 1H), 7.56-7.63 (m, 2H), 7.75-7.85 (m, 2H), 7.74-7.84 (m, 2H), 8.14 (s, 1H), 8.29 (s, 1H) 8.46 (d, 1H), 9.33(s, 1H)
20-22		Ms:562	DMSO-d6: 2.20 (s, 3H), 2.3-2.5 (m, 11H), 2.64 (t, 2H), 3.77 (s, 3H), 3.91 (t, 2H), 6.61(dd, 1H), 6.94 (d, 1H), 7.25-7.31(m, 1H), 7.57 (d, 1H), 7.58-7.64 (m, 1H), 7.74-7.84 (m, 2H), 8.12 (brs, 1H), 8.28 (s, 1H) 8.46 (d, 1H), 9.33(brs, 1H)



20-23		Ms:438	DMSO-d6: 2.42-2.45 (m, 3H), 3.83 (s, 2H), 6.8 (ddd, 1H), 7.02 (dd, 1H), 7.3-7.36 (m, 1H), 7.58-7.64 (m, 1H), 7.74-7.8 (m, 1H), 7.82 (dd, 1H), 7.85 (dd, 1H), 8.18 (brs, 1H), 8.31 (s, 1H), 8.41 (d, 1H), 9.3 (brs, 1H)
20-24		Ms:438	DMSO-d6: 2.41-2.45 (m, 3H), 3.79 (s, 2H), 6.74 (ddd, 1H), 7.0 (dd, 1H), 7.22-7.28 (m, 1H), 7.49-7.55 (m, 1H), 7.6 (dd, 1H), 7.75-7.8 (m, 2H), 8.21 (s, 1H), 8.37 (brs, 1H), 8.39-8.45 (m, 1H), 9.34 (brs, 1H)
20-25		Ms:547	DMSO-d6: 1.24-1.38 (m, 2H), 1.64-1.8 (m, 3H), 1.83-1.92 (m, 2H), 2.16 (s, 3H), 2.41-2.45(m, 3H), 2.76-2.83 (m, 2H), 3.75 (s, 3H), 3.84 (d, 2H), 6.48 (dd, 1H), 6.64 (d, 1H), 7.2-7.25 (m, 1H), 7.41 (d, 1H), 7.43-7.5 (m, 1H), 7.74-7.8 (m, 2H), 8.16 (s, 1H), 8.26 (brs, 1H), 8.44-8.5 (m, 1H), 9.34 (brs, 1H)
20-26		Ms:547	DMSO-d6: 1.18-1.3 (m, 2H), 1.56-1.7 (m, 3H), 1.8-1.88 (m, 2H), 2.15 (s, 3H), 2.41-2.45(m, 3H), 2.73-2.8 (m, 2H), 3.75 (s, 3H), 3.65 (d, 2H), 3.77 (s, 3H), 6.57 (dd, 1H), 6.93 (d, 1H), 7.25 (dd, 1H), 7.51-7.6 (m, 2H), 7.7-7.9 (m, 2H), 8.09 (brs, 1H), 8.28 (s, 1H), 8.45 (d, 1H), 9.31 (brs, 1H)
20-27		Ms:533	DMSO-d6: 1.62-1.72 (m, 2H), 1.9-1.99 (m, 2H), 2.3-2.35 (m, 5H), 2.41-2.45(m, 3H), 2.64-2.74 (m, 2H), 3.75 (s, 3H), 4.35-4.43 (m, 1H), 6.52 (dd, 1H), 6.65 (d, 1H), 7.19-7.25 (m, 1H), 7.41 (d, 1H), 7.43-7.49 (m, 1H), 7.74-7.8 (m, 2H), 8.16 (s, 1H), 8.27 (brs, 1H), 8.42-8.5 (m, 1H), 9.34 (brs, 1H)
20-28		Ms:547	DMSO-d6: 0.96-1.2 (m, 2H), 1.75-1.9 (m, 1H), 2.2-2.3 (m, 1H), 2.35-2.45 (m, 1H), 2.41-2.45(m, 2H), 2.43(d, 3H), 2.6-3.0 (m, 3H), 3.76 (s, 3H), 4.85-5.0 (m, 1H), 6.43-6.49 (m, 1H), 6.57-6.64 (m, 1H), 7.18-7.25 (m, 1H), 7.39-7.52 (m, 2H), 7.73-7.83 (m, 2H), 8.17 (s, 1H), 8.27 (brs, 1H), 8.44-8.51 (m, 1H), 9.35 (brs, 1H)



20-29		Ms:519	DMSO-d6: 1.74-1.83 (m, 1H), 2.23-2.31 (m, 1H), 2.28 (s, 3H), 2.35-2.4 (m, 1H), 2.41-2.45(m, 3H), 2.58-2.63 (m, 1H), 2.63-2.7 (m, 1H), 2.78-2.83 (m, 1H), 3.75 (s, 3H), 4.86-4.92 (m, 1H), 6.43 (dd, 1H), 6.58 (d, 1H), 7.19-7.25 (m, 1H), 7.41 (d, 1H), 7.44-7.51 (m, 1H), 7.73-7.83 (m, 2H), 8.16 (s, 1H), 8.26 (brs, 1H), 8.43-8.52 (m, 1H), 9.34 (brs, 1H)
20-30		Ms : 533	DMSO-d6: 1.04 (t, 3H), 1.74-1.82 (m, 1H), 2.23-2.33 (m, 1H), 2.47-2.5(m, 6H), 2.62-2.72 (m, 2H), 2.8-2.87 (m, 1H), 3.75 (s, 3H), 4.86-4.92 (m, 1H), 6.44 (dd, 1H), 6.59 (d, 1H), 7.19-7.25 (m, 1H), 7.41 (d, 1H), 7.44-7.51 (m, 1H), 7.73-7.8 (m, 2H), 8.16 (s, 1H), 8.26 (brs, 1H), 8.44-8.51 (m, 1H), 9.34 (brs, 1H)
20-31		Ms : 518	DMSO-d6: 2.23(s, 3H), 2.38-2.47 (m, 7H), 2.87-2.93 (m, 4H), 3.75 (s, 3H), 6.63 (dd, 1H), 6.93 (d, 1H), 7.22-7.28 (m, 1H), 7.42 (d, 1H), 7.48-7.54 (m, 1H), 7.76-7.84 (m, 1H), 8.2 (s, 1H), 8.25(s, 1H), 8.43 (dd, 1H) 9.29 (s, 1H)
20-32		Ms : 586	DMSO-d6: 1.35-1.55 (m, 8H), 1.66-1.75 (m, 2H), 2.23(s, 3H), 2.41-2.45 (m, 3H), 3.74 (s, 3H), 6.63 (dd, 1H), 6.91 (d, 1H), 7.21-7.28 (m, 1H), 7.44 (d, 1H), 7.48-7.54 (m, 1H), 7.76-7.87 (m, 1H), 8.16 (s, 1H), 8.25 (s, 1H), 8.43 (dd, 1H) 9.29 (s, 1H)
20-33		Ms : 518	DMSO-d6: 1.62-1.71 (m, 1H), 1.95-2.04 (m, 1H), 2.23-2.27 (m, 3H), 2.39-2.43 (m, 3H), 2.93-3.1 (m, 2H), 3.13-3.26 (m, 2H), 3.71 (s, 3H), 6.19 (dd, 1H), 6.88 (d, 1H), 7.07-7.13 (m, 1H), 7.13-7.2 (m, 1H), 7.4-7.48 (m, 1H), 7.75 (dd, 1H), 8.06 (brs, 1H), 8.18 (s, 1H), 8.4 (d, 1H)
20-34		Ms : 546	DMSO-d6: 2.02 (m, 1H), 2.42-2.46 (m, 3H), 2.71-2.91 (m, 4H), 3.44-3.51 (m, 4H), 3.76 (s, 3H), 6.66 (dd, 1H), 6.94 (d, 1H), 7.21-7.27 (m, 1H), 7.75-7.85 (m, 2H), 8.19 (s, 1H), 8.26 (s, 1H), 8.41 (d, 1H), 9.28 (brs, 1H).

20-35		MS (ESI) 464 (M+H)	HPLC Retention time (min) 2.68
-------	---	-----------------------	-----------------------------------


The following 2-[5-chloro-2-(subst.phenylamino)-pyrimidin-4-ylamino]-N-sec-butyl-benzene-sulfonamides are prepared from 2-(5-chloro-2-chloro-pyrimidin-4-ylamino)-N-sec-butyl-benzenesulfonamide and the corresponding aniline following the procedure of Example 7A






ExptNo.	Rx	R _f (solvent) or MS	NMR (400MHz) , δ (ppm)
21-1		0.35 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 0.62(t, 3H), 0.88(d, 3H), 1.22-1.29(m, 2H), 2.23(s, 3H), 2.45-2.47(m, 4H), 3.05-3.14(m, 5H), 3.75(s, 3H), 6.40-6.43(m, 1H), 6.62(s, 1H), 7.18-7.22(m, 1H), 7.39-7.47(m, 2H), 7.80-7.82(m, 1H), 8.15-8.16(m, 2H), 8.44-8.46(m, 1H), 9.32(s, 1H)
21-2		0.30 (<i>n</i> -hexane: AcOEt=3:1)	DMSO-d ₆ : 0.62(t, 3H), 0.87(d, 3H), 1.17-1.26(m, 2H), 3.03-3.10(m, 1H), 3.79(s, 3H), 6.66-6.71(m, 1H), 6.96-7.00(m, 1H), 7.21-7.25(m, 1H), 7.47-7.51(m, 1H), 7.60-7.64(m, 1H), 7.79-7.83(m, 2H), 8.21(s, 1H), 8.31(s, 1H), 8.35-8.37(m, 1H), 9.29(s, 1H)

21-3		0.30 (<i>n</i> -hexane: AcOEt=1:1)	DMSO-d ₆ : 0.61(t, 3H), 0.87(d, 3H), 1.21-1.29(m, 2H), 1.58-1.67(m, 2H), 1.86-1.93(m, 2H), 2.14-2.20(m, 5H), 2.59-2.67(m, 2H), 3.06-3.08(m, 1H), 3.74(s, 3H), 4.32-4.36(m, 1H), 6.46-6.48(m, 1H), 6.63(d, 1H), 7.17-7.21(m, 1H), 7.40-7.50(m, 2H), 7.79-7.81(m, 2H), 8.16(s, 1H), 8.21(bs, 1H), 8.35-8.42(m, 1H), 9.29(s, 1H)
21-4		0.30 (<i>n</i> -hexane: AcOEt=1:1)	DMSO-d ₆ : 0.61(t, 3H), 0.87(d, 3H), 1.22-1.29(m, 2H), 2.43-2.47(m, 2H), 2.61-2.63(m, 1H), 2.68-2.70(m, 2H), 3.04-3.11(m, 1H), 3.56-3.60(m, 5H), 3.75(s, 3H), 3.93-3.96(m, 1H), 4.08-4.11(m, 2H), 6.45-6.47(m, 1H), 6.64(d, 1H), 7.18-7.22(m, 1H), 7.43-7.46(m, 2H), 7.80-7.82(m, 2H), 8.17(s, 1H), 8.21(s, 1H), 8.42-8.44(m, 1H), 9.31(s, 1H)


The following 2-[5-chloro-2-(subst.phenylamino)-pyrimidin-4-ylamino]-N-*iso*-butyl-benzene-sulfonamides are prepared from 2-(5-chloro-2-chloro-pyrimidin-4-ylamino)-N-*sec*-butyl-benzenesulfonamide and the corresponding aniline following the procedure of Example 7A









ExptNo.	Rx	R _f (solvent) or MS	NMR (400MHz) , δ (ppm)
22-1		0.30 (<i>n</i> -hexane: AcOEt=3:1)	DMSO-d ₆ : 0.69(d, 6H), 1.52-1.59(m, 1H), 2.57-2.58(m, 2H), 3.82(s, 3H), 6.75-6.80(m, 1H), 6.99-7.02(m, 1H), 7.29-7.33(m, 1H), 7.56-7.60(m, 1H), 7.82-7.93(m, 3H), 8.14(bs, 1H), 8.31(s, 1H), 8.33(s, 1H), 9.23(s, 1H)




22-2		0.30 (<i>n</i> -hexane: AcOEt=3:1)	CDCl ₃ : 0.74(d, 6H), 1.57-1.64(m, 1H), 2.72-2.76(m, 2H), 3.88(s, 3H), 4.55-4.56(m, 1H), 6.52-6.57(m, 1H), 6.62-6.65(m, 1H), 7.24-7.28(m, 2H), 7.36(bs, 1H), 7.56-7.60(m, 1H), 7.95-8.08(m, 1H), 8.10-8.14(m, 2H), 8.36-8.39(m, 1H), 8.98(bs, 1H)
22-3		0.54 (AcOEt)	DMSO-d ₆ : 0.73(d, 6H), 1.55-1.62(m, 1H), 2.56-2.59(m, 2H), 3.10-3.12(m, 4H), 3.74-3.76(m, 7H), 6.43-6.46(m, 1H), 6.65(d, 1H), 7.20-7.24(m, 1H), 7.40-7.48(m, 2H), 7.76-7.78(m, 1H), 7.90-7.95(m, 1H), 8.16(s, 1H), 8.17(s, 1H), 8.43-8.45(m, 1H), 9.32(s, 1H)

The following 2-[5-chloro-2-(subst.phenylamino)-pyrimidin-4-ylamino]-N-(1-ethyl-propyl)-benzenesulfonamides are prepared from 2-(5-chloro-2-chloro-pyrimidin-4-ylamino)-N-(1-ethyl-propyl)-benzenesulfonamide and the corresponding aniline following the procedure of Example 7A




ExptNo.	Rx	R _f (solvent) or MS	NMR (400MHz) , δ (ppm)
23-1		0.46 (MeOH: CH ₂ Cl ₂ =3:7)	DMSO-d ₆ : 0.58(t, 6H), 1.14-1.34(m, 4H), 1.58-1.68(m, 2H), 1.87-1.96(m, 2H), 2.12-2.22(m, 2H), 2.18(s, 3H), 2.57-2.65(m, 2H), 2.86-2.96(m, 1H), 3.75(s, 3H), 4.30-4.39(m, 1H), 6.46(dd, 1H), 6.63(d, 1H), 7.19(dd, 1H), 7.39-7.48(m, 2H), 7.75-7.84(m, 2H), 8.18(s, 1H), 8.20 (s, 1H), 8.39(m, 1H), 9.33(bs, 1H)




23-2		0.35 (<i>n</i> -hexane: AcOEt=1:1)	CDCl ₃ : 0.59(t, 6H), 1.14-1.34(m, 4H), 2.23(s, 3H), 2.45-2.47(m, 4H), 2.90-2.95(m, 1H), 3.11-3.14(m, 4H), 3.76 (s, 3H), 6.39-6.42(m, 1H), 6.62(s, 1H), 7.18-7.22(m, 1H), 7.41-7.46(m, 2H), 7.76-7.82(m, 2H), 8.12(s, 1H), 8.16(s, 1H), 8.43-8.44(m, 1H), 9.35(s, 1H)
23-3		0.41 (MeOH: CH ₂ Cl ₂ =1:4)	DMSO-d ₆ : 0.59(t, 6H), 1.16-1.35(m, 4H), 1.75-1.89(m, 1H), 2.08-2.15(m, 1H), 2.32(s, 3H), 2.90-3.02(m, 2H), 3.21-3.45(m, 4H), 3.73(s, 3H), 6.02(dd, 1H), 6.18(d, 1H), 7.16(dd, 1H), 7.27(d, 1H), 7.35-7.45(m, 1H), 7.77-7.82(m, 2H), 8.10 (s, 1H) 8.12 (s, 1H), 8.45-8.55(m, 1H), 9.38(s, 1H)
23-4		0.41 (MeOH: CH ₂ Cl ₂ =1:4)	DMSO-d ₆ : 0.60(t, 6H), 1.04(t, 3H), 1.17-1.35(m, 4H), 1.76-1.83(m, 1H), 2.10-2.15(m, 1H), 2.56-2.64(m, 2H), 2.91-3.01(m, 2H), 3.21-3.47(m, 4H), 3.74(s, 3H), 6.02(dd, 1H), 6.18(d, 1H), 7.14-7.17(m, 1H), 7.28(d, 1H), 7.35-7.45(m, 1H), 7.77-7.82(m, 2H), 8.11 (s, 1H) 8.12 (s, 1H), 8.45-8.55(m, 1H), 9.38(s, 1H)
23-5		0.25 (<i>n</i> -hexane: AcOEt=1:1)	DMSO-d ₆ : 0.58(t, 6H), 1.06-2.16(m, 11H), 2.16(s, 3H), 2.62-2.67(m, 1H), 2.81-2.94(m, 2H), 3.75(s, 3H), 3.80-3.89(m, 2H), 6.41-6.44(m, 1H), 6.62(d, 1H), 7.17-7.21(m, 1H), 7.42-7.47(m, 2H), 7.77-7.82(m, 2H), 8.18(s, 1H), 8.19(s, 1H), 8.35 -8.42(m, 1H), 9.35(s, 1H)
23-6		Ms : 561	DMSO-d ₆ : 0.59 (t, 6H), 1.14-1.38 (m, 4H), 2.87-2.98 (m, 1H), 3.1 (t, 4H), 3.72-3.79 (m, 7H), 6.42 (dd, 1H), 6.64 (d, 1H), 7.18-7.24 (m, 1H), 7.42-7.5 (m, 2H), 7.77 (d, 1H), 7.81 (dd, 1H), 8.13 (s, 1H), 8.17 (s, 1H), 8.4-8.5 (m, 1H), 9.36 (s, 1H)


23-7		Ms : 575	DMSO-d6 : 0.58 (t, 6H), 1.13-1.37 (m, 4H), 1.72-1.82 (m, 1H), 2.21-2.31 (m, 4H), 2.32-2.4 (m, 1H), 2.54-2.61 (m, 1H), 2.62-2.68 (m, 1H), 2.75-2.82 (m, 1H), 2.87-2.97 (m, 1H), 3.75 (s, 3H), 4.84-4.91 (m, 1H), 6.37 (dd, 1H), 6.56 (d, 1H), 7.14-7.24 (m, 1H), 7.38-7.52 (m, 2H), 7.72-7.86 (m, 1H), 8.12-8.25 (m, 2H), 8.34-8.45 (m, 1H), 9.33 (brs, 1H)
23-8		Ms : 605	DMSO-d6 : 0.58 (t, 6H), 1.14-1.36 (m, 4H), 2.43-2.53 (m, 4H), 2.69 (t, 2H), 2.89-2.95 (m, 1H), 3.59 (t, 4H), 3.76 (s, 3H), 4.09 (t, 1H), 6.45 (dd, 1H), 6.64 (d, 1H), 7.17-7.23 (m, 1H), 7.41-7.52 (m, 2H), 7.78 (d, 1H), 7.81 (dd, 1H), 8.18 (s, 1H), 8.19 (s, 1H), 8.36-8.46 (m, 1H), 9.35 (s, 1H)
23-9		Ms : 618	DMSO-d6 : 0.58 (t, 6H), 1.14-1.37 (m, 4H), 2.15 (s, 1H), 2.25-2.4 (m, 4H), 2.45-2.55 (m, 4H), 2.68 (t, 2H), 2.88-2.97 (m, 1H), 3.76 (s, 3H), 4.07 (t, 1H), 6.44 (dd, 1H), 6.64 (d, 1H), 7.15-7.23 (m, 1H), 7.41-7.51 (m, 2H), 7.7-7.84 (m, 2H), 8.12-8.22 (m, 1H), 8.34-8.44 (m, 1H), 9.34 (s, 1H)


The following 2-[5-chloro-2-(subst.phenylamino)-pyrimidin-4-ylamino]-N-iso-butyl-benzene-sulfonamides are prepared from 2-(5-chloro-2-chloro-pyrimidin-4-ylamino)-N-cyclobutyl-benzenesulfonamide and the corresponding aniline following the procedure of Example 7A




ExpNo.	Rx	Rf (solvent) or MS	NMR (400MHz) , δ (ppm)
--------	----	-----------------------	-------------------------------


24-1		0.35 (<i>n</i> -hexane: AcOEt=1:1)	DMSO-d ₆ : 1.37-1.48(m, 2H), 1.69-1.91(m, 4H), 3.09-3.12(m, 4H), 3.63-3.74(m, 1H), 3.76(s, 3H), 6.43-6.45(m, 1H), 6.63(d, 1H), 7.18-7.22(m, 1H), 7.41-7.47(m, 2H), 7.76-7.78(m, 1H), 8.17-8.24(m, 3H), 8.46(d, 1H), 9.33(s, 1H)
24-2		0.46 (MeOH: CH ₂ Cl ₂ =3:7)	DMSO-d ₆ : 1.37-1.93(m, 10H), 2.18(s, 3H), 2.59-2.62(m, 1H), 3.60-3.74(m, 1H), 3.77(s, 3H), 4.32-4.36(m, 1H), 6.46-6.49(m, 1H), 6.62(d, 1H), 7.16-7.20(m, 1H), 7.41-7.44(m, 2H), 7.75-7.77(m, 1H), 8.16(s, 1H), 8.22 (bs, 1H), 8.40-8.42(m, 1H), 9.30(bs, 1H)
24-3		0.46 (MeOH: CH ₂ Cl ₂ =1:4)	CDCl ₃ : 1.45-1.75(m, 5H), 1.94-2.06(m, 6H), 2.29-2.37(m, 1H), 3.21-3.56(m, 4H), 3.72-3.81(m, 1H), 3.86(s, 3H), 4.55-4.65(m, 1H), 4.90(d, 1H), 5.72(d, 1H), 6.07(bs, 1H), 6.15(bs, 1H), 7.18-7.22(m, 2H), 7.52-7.56(m, 1H), 7.89-7.94(m, 2H), 8.08(s, 1H), 8.50(d, 1H), 9.00(s, 1H)










Exptl No.	Rx	Ms	NMR (400MHz), δ (ppm)
25-1		Ms : 559	DMSO-d ₆ : 1.2-1.38 (m, 4H), 1.4-1.65 (m, 4H), 3.11 (t, 4H), 3.42-3.5 (m, 1H), 3.7-3.8 (m, 7H), 6.44 (dd, 1H), 6.64 (d, 1H), 7.18-7.26 (m, 1H), 7.38-7.5 (m, 2H), 7.81 (d, 1H), 7.88-7.96 (m, 1H), 8.16 (s, 1H), 8.17 (s, 1H), 8.4-8.5 (m, 1H), 9.34 (s, 1H)





25-2		Ms : 587	DMSO-d ₆ : 1.2-1.38 (m, 4H), 1.42-1.6 (m, 6H), 1.88-1.98 (m, 2H), 2.1-2.25 (m, 5H), 2.55-2.65 (m, 2H), 3.4-3.5 (m, 1H), 3.74 (s, 3H), 4.3-4.4 (m, 1H), 6.48 (dd, 1H), 6.63 (d, 1H), 7.18-7.24 (m, 1H), 7.38-7.47 (m, 1H), 7.77-7.82 (m, 1H), 7.88-7.96 (m, 1H), 8.17 (s, 1H), 8.22 (s, 1H), 8.36-8.46 (m, 1H), 9.31 (s, 1H)
------	---	----------	--






The following 5-Chloro-N²-(substituted phenyl)-N⁴-[2-(propane-1-sulfonyl)-phenyl]-pyrimidine-2,4-diamine are prepared from (2,5-Dichloro-pyrimidin-4-yl)-[2-(propane-1-sulfonyl)-phenyl]-amine and the corresponding aniline following the procedure of Example 7A













Exptl No.	Rx	R _f (solvent), MS or Mp	NMR (400MHz), δ (ppm)
26-1		0.58 (AcOEt)	CDCl ₃ : 0.97(t,3H), 1.72-1.82(m, 2H), 3.08-3.14(m, 6H), 3.87-3.89(m, 7H), 6.46(dd, 1H), 6.53(d,1H), 7.24-7.28(m,1H), 7.30(s, 1H), 7.60-7.64(m, 1H), 7.94(dd, 1H), 8.05(d, 1H), 8.15(s, 1H), 8.59(d, 1H), 9.40(s, 1H)
26-2		0.57 (MeOH: AcOEt=1:4)	CDCl ₃ : 0.98(t, 3H), 1.85-1.68(m, 2H), 2.15(s, 3H), 3.16-3.07(m, 6H), 3.67-3.62(m, 2H), 3.81-3.78(m, 2H), 3.89(s, 3H), 6.47(d, 1H), 6.55(d, 1H), 7.36-7.33(m, 1H), 7.62 (dd, 1H), 7.95(dd, 1H), 8.08(d, 1H), 8.15(s, 1H), 8.58(d, 1H), 9.41(s, 1H)






26-3		0.13 (MeOH: AcOEt=1:4)	CDCl ₃ : 0.97(t, 3H), 1.43-1.52(m, 2H), 1.52-1.67 (m, 4H), 1.69-1.72(m, 4H), 1.90-1.98(m, 2H), 2.34-2.46(m, 1H), 2.51-2.59(m, 4H), 2.64-2.74(m, 2H), 3.11(dd, 2H), 3.64-3.73(m, 2H), 3.87(s, 3H), 6.47(dd, 1H), 6.56 (d, 1H), 7.24-7.33(m, 1H), 7.62(dd, 1H), 7.94(dd, 1H), 8.00(d, 1H), 8.14(s, 1H), 8.59(d, 1H), 9.39(bs, 1H).
26-4		0.22 (AcOEt)	CDCl ₃ : 0.97(t, 3H), 1.45(d, 1H), 1.68-1.82(m, 4H), 2.0-2.1(m, 2H), 2.91(ddd, 2H), 3.10(ddd, 2H), 3.46-3.51 (m, 2H), 3.84-3.92(m, 1H), 3.88 (s, 1H), 6.48(dd, 1H), 6.57(d, 1H), 7.23-7.32 (m, 1H), 7.62(dd, 1H), 7.94(dd, 1H), 8.02 (dd, 1H), 8.14(s, 1H), 8.59(d, 1H), 9.39(bs, 1H)
26-5		0.1 (AcOEt)	CDCl ₃ : 0.97(t, 3H), 1.71-1.82(m, 2H), 1.86-1.98(m, 2H), 2.01-2.08(m, 2H), 2.25-2.37(m, 1H), 2.75 (ddd, 2H), 3.10(ddd, 2H), 3.63-3.66(m, 2H), 3.88(s, 3H), 5.25-5.40(m, 1H), 5.40-5.58 (m, 1H), 6.48(dd, 1H), 6.57(d, 1H), 7.22-7.34 (m, 1H), 7.62(ddd, 1H), 7.93 (d, 1H), 7.94 (dd, 1H), 8.02 (d, 1H), 8.14(s, 1H), 8.59(d, 1H), 9.40(m, 1H)
26-6		MS 587	CDCl ₃ : 0.97 (t, 3H), 1.77 (ddd, 2H), 2.00-1.85 (m, 4H), 2.27-2.18(m, 1H), 2.72(ddd, 2H) 3.12-3.08 (m, 2H), 3.69-3.61(m, 2H), 3.58-3.46(m, 1H), 3.64 (t, 2H), 3.80(t, 2H), 3.88(s, 3H), 5.56-5.46 (m, 1H), 6.47(dd, 1H), 6.55(d, 1H), 7.32-7.23 (m, 1H), 7.30(bs, 1H), 7.64-7.60(m, 1H), 7.94(dd, 1H), 8.02(d, 1H), 8.14(s, 1H), 8.59(d, 1H), 9.40(s, 1H)
26-7		MS 587	CDCl ₃ : 0.98 (t, 3H), 1.46(bs, 6H) 1.82-1.73 (m, 2H), 2.17(s, 3H), 3.58-3.46(m, 1H), 2.95-2.84 (m, 2H), 3.12-3.08 (m, 2H), 3.90(s, 3H), 6.48(dd, 1H), 6.52(d, 1H), 7.30-7.22 (m, 1H), 7.31(bs, 1H), 7.66-7.60(m, 1H), 7.95(dd, 1H), 8.06(d, 1H), 8.15(s, 1H), 8.59(d, 1H), 9.43(s, 1H)







26-8		MS 573	CDCl ₃ : 0.97 (t, 3H), 1.19(t, 3H), 1.77 (ddd, 2H), 2.41(m, 2H), 3.18-3.09(m, 6H), 3.68-3.64 (m, 2H), 3.85-3.78 (m, 2H), 3.89(s,3H), 6.47(dd, 1H), 6.55(d, 1H), 7.29-7.25 (m, 1H), 7.34(bs, 1H), 7.64-7.60(m, 1H), 7.95(dd, 1H), 8.07(d, 1H), 8.15(s, 1H), 8.58(d, 1H), 9.41(s, 1H)
26-9		MS 587	CDCl ₃ : 0.97 (t, 3H), 1.17(d, 3H) 1.76 (ddd, 2H), 2.88-2.81(m, 2H), 3.18-3.05(m, 6H), 3.74-3.67 (m, 2H), 3.86-3.78 (m, 2H), 3.89(s,3H), 6.47(dd, 1H), 6.55(d, 1H), 7.29-7.20(m, 1H), 7.34(bs, 1H), 7.64-7.60(m, 1H), 7.95(dd, 1H), 8.07(d, 1H), 8.15(s, 1H), 8.58(d, 1H), 9.41(s, 1H)
26-10		MS 517	CDCl ₃ : 0.97 (t, 3H), 1.76 (ddd, 2H), 2.86(d, 3H), 3.14-3.08(m, 2H), 3.13(t, 4H), 3.55 (t, 4H), 3.89(s,3H), 4.48-4.39 (m, 1H), 6.46(dd, 1H), 6.55(d, 1H), 7.29-7.21(m, 1H), 7.34(bs, 1H), 7.64-7.60(m, 1H), 7.95(dd, 1H), 8.06(d, 1H), 8.15(s, 1H), 8.58(d, 1H), 9.41(s, 1H)
26-11		MS 587	CDCl ₃ : 0.98 (t, 3H), 1.51 (s, 6H), 1.82-1.72 (m, 1H), 2.13 (s, 3H), 3.12-3.08 (m, 2H), 3.26(s, 2H), 3.44 (t, 2H), 3.74(t, 2H), 3.88(s,3H), 5.56-5.46 (m, 1H), 6.45(dd, 1H), 6.51(d, 1H), 7.00(bs, 1H), 7.62-7.58 (m, 1H), 7.64-7.60(m, 1H), 7.93(d, 1H), 7.96(dd, 1H), 8.13(s, 1H), 8.62(d, 1H), 9.42(s, 1H)



26-12		MS 559	CDCl ₃ : 0.98 (t, 3H), 1.81-1.71 (m, 3H), 1.95-1.84 (m, 3H), 2.68-2.63(m, 1H), 3.12-3.08 (m, 4H), 3.28(d, 2H), 3.89(s,3H), 5.45-5.38 (m, 1H), 6.53(dd, 1H), 6.59(d, 1H), 6.71-6.62 (m, 1H), 7.28-7.21 (m, 1H), 7.35(bs, 1H), 7.65-7.61(m, 1H), 7.95(dd, 1H), 8.08(d, 1H), 8.15(s, 1H), 8.58(d, 1H), 9.41(s, 1H)
26-13		MS 559	CDCl ₃ : 0.98 (t, 3H), 1.81-1.71 (m, 3H), 1.95-1.84 (m, 3H), 2.68-2.63(m, 1H), 3.12-3.08 (m, 4H), 3.28(d, 2H), 3.89(s,3H), 5.45-5.38 (m, 1H), 6.53(dd, 1H), 6.59(d, 1H), 6.71-6.62 (m, 1H), 7.28-7.21 (m, 1H), 7.35(bs, 1H), 7.65-7.61(m, 1H), 7.95(dd, 1H), 8.08(d, 1H), 8.15(s, 1H), 8.58(d, 1H), 9.41(s, 1H)
26-14		MS 559	CDCl ₃ : 0.98 (t, 3H), 1.85-1.74 (m, 3H), 2.00-1.86 (m, 3H), 2.70-2.51(m, 1H), 3.13-3.08 (m, 4H), 3.29-3.27 (m, 2H), 3.89(s,3H), 5.46-5.37 (m, 1H), 6.53(dd, 1H), 6.59(d, 1H), 6.69-6.56 (m, 1H), 7.29-7.19 (m, 1H), 7.34(bs, 1H), 7.65-7.61(m, 1H), 7.95(dd, 1H), 8.08(d, 1H), 8.15(s, 1H), 8.58(d, 1H), 9.41(s, 1H)
26-15		MS 559	CDCl ₃ : 0.99(t, 3H), 1.79-1.72 (m, 2H), 2.92 (t, 2H), 2.98(t, 2H), 3.16-3.12 (m, 2H), 3.53(t, 2H), 3.67(t, 2H), 3.87(s,3H), 6.54(dd, 1H), 6.82 (d, 1H), 7.29-7.19 (m, 1H), 7.56(bs, 1H), 7.67-7.62(m, 1H), 7.96(dd, 1H), 8.07(d, 1H), 8.21(s, 1H), 8.59(dd, 1H), 9.46(s, 1H)
26-16		MS 561	CDCl ₃ : 0.98 (t, 3H), 1.81-1.72 (m, 2H), 2.49 (t, 4H), 2.66 (t, 2H), 3.12-3.08(m, 2H), 3.18 (t, 2H), 3.74(t, 4H), 3.86(s,3H), 6.53(dd, 1H), 6.20(dd, 1H), 6.26 (d, 1H), 7.13(bs, 1H), 7.25-7.21 (m, 1H), 7.62-7.57(m, 1H), 7.87(dd, 1H), 7.93(dd, 1H), 8.12(s, 1H), 8.62(d, 1H), 9.40(s, 1H)

26-17		MS 518	CDCl ₃ : 0.98 (t, 3H), 1.78-1.73 (m, 2H), 2.49 (t, 4H), 2.66 (t, 2H), 2.94-2.92 (m, 4H), 3.15-3.11(m, 2H), 3.76-3.73(m, 4H), 3.88(s,3H), 6.52(dd, 1H), 6.82(d, 1H), 7.28-7.24 (m, 1H), 7.57(bs, 1H), 7.25-7.21 (m, 1H), 7.68-7.63(m, 1H), 7.95(dd, 1H), 8.02(d, 1H), 8.20(s, 1H), 8.56(d, 1H), 9.41(s, 1H)
26-18		MS 559	CDCl ₃ : 0.97 (t, 3H), 1.81-1.72 (m, 2H), 2.08-2.00 (m, 2H), 2.49 (t, 4H), 2.66 (t, 2H), 2.40 (t, 2H), 3.59 (t, 2H), 3.69(t, 2H), 3.87(s,3H), 6.41 (dd, 1H), 6.51(d, 1H), 7.29-7.25 (m, 2H), 7.65-7.60(m, 1H), 7.95(dd, 1H), 8.05(d, 1H), 8.15(s, 1H), 8.56(d, 1H), 9.41(s, 1H)
26-19		MS 587	CDCl ₃ : 0.98 (t, 3H), 1.82-1.73 (m, 2H), 2.14 (s, 3H), 3.12-3.08 (m, 2H), 3.55-3.45(m, 2H), 3.66-3.56 (m, 4H), 3.79-3.68 (m, 2H), 3.95(s,3H), 6.95(dd, 1H), 7.03 (d, 1H), 7.32-7.28(m, 1H), 7.69-7.64 (m, 1H), 7.71(s, 1H), 7.97(dd, 1H), 8.22 (s, 1H), 8.39(d, 1H), 8.52(d, 1H), 9.46(s, 1H)
26-20		MS 546	CDCl ₃ : 0.97 (t, 3H), 1.82-1.73 (m, 2H), 3.12-3.08 (m, 2H), 3.80-3.58(m, 8H), 3.94(s,3H), 6.94(dd, 1H), 7.02 (d, 1H), 7.32-7.28(m, 1H), 7.69-7.64 (m, 1H), 7.32-7.28(m, 1H), 7.97(dd, 1H), 8.21 (s, 1H), 8.34(d, 1H), 8.52(d, 1H), 9.45(s, 1H)
26-21		MS 615	CDCl ₃ : 0.97 (t, 3H), 1.82-1.72 (m, 2H), 2.71 (t, 3H), 3.05 (s, 2H), 3.10 (m, 2H), 3.18 (t, 4H), 3.88(s,3H), 4.17-4.08 (m, 1H), 6.47 (dd, 1H), 6.54 (d, 1H), 6.99-6.89(m, 1H), 7.28-7.24 (m, 1H), 7.31(bs, 1H), 7.65-7.60 (m, 1H), 7.32-7.28(m, 1H), 7.95(dd, 1H), 8.05 (d, 1H), 8.15(s, 1H), 8.59(d, 1H), 9.41(s, 1H)


26-22		MS 530	CDCl ₃ : 0.98 (t, 3H), 1.80-1.74 (m, 2H), 3.12-3.08 (m, 2H), 3.45-3.42 (m, 2H), 3.55-3.53 (m, 2H), 3.87 (s, 2H), 3.89(s,3H), 5.98-5.89 (m, 1H), 6.44 (dd, 1H), 6.50(d, 1H), 7.35-7.19 (m, 2H), 7.62-7.58(m, 1H), 7.95(dd, 1H), 8.09 (d, 1H), 8.15(s, 1H), 8.57(d, 1H), 9.43(s, 1H)
26-23		MS 558	CDCl ₃ : 0.97 (t, 3H), 1.10 (s, 3H), 1.12 (s, 3H), 1.80-1.74(m, 2H), 2.80-2.63 (m, 5H), 3.12-3.08 (m, 2H), 3.19-3.17 (m, 4H), 3.87(s,3H), 6.48 (dd, 1H), 6.56(d, 1H), 7.30-7.23 (m, 2H), 7.62-7.58(m, 1H), 7.94(dd, 1H), 8.00 (d, 1H), 8.14(s, 1H), 8.59(d, 1H), 9.40(s, 1H)
26-24		MS 544	CDCl ₃ : 0.98 (t, 3H), 1.81-1.72 (m, 2H), 2.03-1.91 (m, 1H), 2.28-2.19 (m, 1H), 2.33 (s, 6H), 2.92-2.84 (m, 1H), 3.12-3.08 (m, 2H), 3.17(t, 1H), 3.35(ddd, 1H), 3.51-3.42 (m, 2H), 3.87 (s, 3H), 6.11 (dd, 1H), 6.14 (d,1H), 7.09 (s, 1H), 7.26-7.20(m, 1H), 7.60-7.56 (m, 1H), 7.85(d, 1H), 7.92(dd, 1H), 8.11(s, 1H), 8.38(d, 1H), 9.41(s, 1H)
26-25		MS 530	CDCl ₃ : 0.98 (t, 3H), 1.82-1.71 (m, 2H), 1.96-1.86 (m, 1H), 2.33-2.20 (m, 1H), 2.51 (s, 1H), 3.17-3.08 (m, 3H), 3.35-3.30 (m, 1H), 3.54-3.30 (m, 3H), 3.87 (s, 3H), 6.12 (dd, 1H), 6.16 (d,1H), 7.09 (s, 1H), 7.32-7.21(m, 1H), 7.58 (dd, 1H), 7.85(d, 1H), 7.92(dd, 1H), 8.11(s, 1H), 8.64(d, 1H), 9.40(s, 1H)
26-26		MS 546	CDCl ₃ : 0.98 (t, 3H), 1.83-1.71 (m, 2H), 1.98-1.81 (m, 2H), 2.16-2.02(m, 2H), 2.53-2.28 (m, 5H), 2.87-2.72 (m, 2H), 3.12-3.08 (m, 2H), 3.88 (s, 3H), 4.32 (bs, 3H), 6.44 (dd, 1H), 6.53(d, 1H), 7.32-7.25 (m, 2H), 7.63-7.59 (m, 2H), 7.94(dd, 1H), 8.04 (d, 1H), 8.15(s, 1H), 8.57(d, 1H), 9.42(s, 1H)


26-27		MS 545	CDCl ₃ : 0.97 (t, 3H), 1.38-1.30 (m, 1H), 1.49-1.40 (m, 2H), 1.70-1.62 (m, 1H), 1.83-1.72 (m, 2H), 1.89 (d, 2H), 2.74-2.10 (m, 2H), 3.12-3.08 (m, 2H), 3.57 (d, 2H), 3.63 (d, 2H), 3.90 (s, 3H), 6.50 (d, 1H), 6.58 (s, 1H), 7.34-7.24 (m, 2H), 7.64-7.60 (m, 1H), 7.94 (dd, 1H), 8.02 (d, 1H), 8.14 (s, 1H), 8.60 (dd, 1H), 9.40 (s, 1H)
26-28		MS 517	CDCl ₃ : 0.97 (t, 3H), 1.88-1.65 (m, 3H), 2.05-1.97 (m, 2H), 2.21-2.08 (m, 1H), 2.67-2.55 (m, 4H), 2.78-2.71 (m, 2H), 3.12-3.08 (m, 2H), 3.61 (d, 2H), 3.87 (s, 3H), 6.47 (dd, 1H), 6.56 (d, 1H), 7.28-7.23 (m, 2H), 7.64-7.60 (m, 1H), 7.94 (dd, 1H), 7.99 (d, 1H), 8.13 (s, 1H), 8.60 (dd, 1H), 9.39 (s, 1H)
26-29		MS 585	CDCl ₃ : 0.97 (t, 3H), 1.89-1.65 (m, 8H), 2.03 (d, 2H), 2.20-2.10 (m, 1H), 2.68-2.58 (m, 4H), 2.78-2.72 (m, 2H), 3.12-3.08 (m, 2H), 3.61 (d, 2H), 3.87 (s, 3H), 6.47 (dd, 1H), 6.56 (d, 1H), 7.30-7.23 (m, 2H), 7.64-7.60 (m, 1H), 7.94 (dd, 1H), 7.99 (dd, 1H), 8.13 (s, 1H), 8.60 (dd, 1H), 9.39 (s, 1H)
26-30		MS 451, 453	0.97 (t, 3H), 1.71-1.82 (m, 2H), 3.06-3.14 (m, 2H), 3.89 (s, 1H), 6.60 (ddd, 1H), 6.66 (dd, 1H), 7.25-7.30 (m, 1H), 7.35 (br.s, 1H), 7.63 (dd, 1H), 7.95 (dd, 1H), 8.09-8.18 (m, 1H), 8.17 (s, 1H), 8.52 (dd, 1H), 9.42 (s, 1H).
26-31		MS 647, 649	0.98 (t, 3H), 1.71-1.83 (m, 2H), 2.18 (s, 3H), 2.47-2.64 (m, 4H), 2.72-2.84 (m, 2H), 3.08-3.15 (m, 2H), 3.42-3.54 (m, 2H), 3.58-3.69 (m, 2H), 3.84 (s, 3H), 3.94-4.03 (m, 2H), 6.45-6.51 (m, 1H), 6.78 (d, 1H), 7.22-7.27 (m, 1H), 7.60 (s, 1H), 7.67-7.74 (m, 1H), 7.93-7.97 (m, 1H), 7.73 (d, 1H), 8.02 (s, 1H), 8.54 (d, 1H), 9.33 (s, 1H).






26-32		m.p. 139,4	400MHz, CDCl ₃ , δ (ppm): 0.98 (t; 3H), 1.55-1.90 (m; 6H), 2.38 (s; 3H), 2.45-2.80 (m; 6H), 3.13 (m; 2H), 3.47 (m; 2H), 3.84 (s; 3H), 6.54 (dd; 1H), 6.79 (d; 1H), 7.23 (dd; 1H); 7.51 (s; 1H), 7.64 (dd; 1H), 7.92 (d; 1H), 8.00 (s; 1H), 8.19 (s; 1H), 8.57 (d; 1H), 9.41 (s; 1H).
26-33		m.p. 163,4	400MHz, CDCl ₃ , δ (ppm): 0.98 (t; 3H), 1.50-1.90 (m; 6H), 2.24 (bs; 1H), 2.45-2.65 (m; 6H), 3.12 (m; 2H), 3.45 (m; 2H), 3.77 (m; 4H), 3.85 (s; 3H), 6.55 (dd; 1H); 6.79 (d; 1H), 7.24 (dd; 1H), 7.52 (s; 1H), 7.64 (dd; 1H), 7.93 (d; 1H), 8.01 (s; 1H), 8.20 (s; 1H), 9.42 (s; 1H).
26-34		m.p. 232,9	400MHz, CDCl ₃ , δ (ppm): 1.00 (s; 3H), 1.78 (m; 2H), 2.83 (s; 3H), 3.03 (m; 2H), 3.12 (m; 2H), 3.38-3.60 (m; 8H), 3.88 (s; 3H), 6.56 (m; 1H), 6.82 (d; 1H), 7.29 (m; 1H), 7.60 (s; 1H), 7.64 (m; 1H), 7.95 (d; 1H), 8.12 (s; 1H), 8.20 (s; 1H), 8.59 (d; 1H), 9.50 (s; 1H).
26-35		m.p. 197,3	400MHz, CDCl ₃ , δ (ppm): 0.99 (t; 3H), 1.43 (m; 1H), 1.63 (m; 2H), 1.77 (m; 2H), 1.90 (m; 2H), 2.70 (m; 2H), 3.13 (m; 2H), 3.28 (m; 2H), 3.75 (s; 1H), 3.84 (s; 3H), 6.55 (m; 1H), 6.80 (d; 1H), 7.24 (m; 1H), 7.53 (s; 1H), 7.64 (s; 1H), 7.93 (d; 1H), 8.02 (s; 1H), 8.20 (s; 1H), 8.58 (d; 1H), 9.41 (s; 1H).
26-36		m.p. 147,6	400MHz, CDCl ₃ , δ (ppm): 1.00 (t; 3H), 1.78 (m; 2H), 3.12 (m; 2H), 3.56 (m; 1H), 3.87 (s; 3H), 6.53 (dd; 1H), 6.80 (d; 1H), 7.30 (dd; 1H), 7.52 (s; 1H), 7.64 (m; 1H), 7.95 (dd; 1H), 8.08 (s; 1H), 8.20 (s; 1H), 8.60 (d; 1H), 9.48 (s; 1H).
26-37		m.p. 143.2	500MHz, CDCl ₃ , δ (ppm): 0.96 (t; 3H), 1.70 (m; 2H), 2.11 (m; 1H), 2.39 (m; 1H), 2.75 (s; 3H), 3.02 (m; 1H), 3.22 (m; 2H), 3.43 (d; 2H), 3.82 (s; 3H), 3.86 (m; 1H), 6.40 (dd; 1H), 6.94 (d; 1H), 7.34 (ddd; 1H), 7.47 (s; 1H), 7.63 (ddd; 1H), 7.93 (dd; 1H), 8.18 (s; 1H), 8.51




			(d; 1H).
26-38		m.p. 133.5	400MHz, CDCl ₃ , δ (ppm): 1.00 (t; 3H), 1.70-1.95 (m; 6H), 2.63 (s; 1H), 2.92 (s; 1H), 3.00-3.25 (m; 5H), 3.89 (s; 3H), 5.42 (s; 1H), 6.70 (s; 1H), 6.83 (m; 2H), 7.25 (m; 1H), 7.55 (s; 1H), 7.63 (m; 1H), 8.95 (m; 1H), 8.15 (s; 1H), 8.23 (s; 1H), 8.54 (d; 1H), 9.45 (s; 1H).
26-39		m.p. 188.8	400MHz, CDCl ₃ , δ (ppm): 0.99 (t; 3H), 1.70-1.90 (m; 3H), 2.08 (m; 1H), 2.28 (s; 6H), 2.83 (s; 1H), 3.00-3.23 (m; 4H), 3.37 (m; 1H), 3.83 (s; 3H), 6.19 (dd; 1H), 6.83 (d; 1H), 7.23 (dd; 1H), 7.50 (s; 1H), 7.59 (m; 2H), 7.93 (d; 1H), 8.19 (s; 1H), 8.60 (d; 1H), 9.42 (s; 1H).

The following 5-Chloro-N²-(substituted phenyl)-N⁴-[2-ethanesulfonyl-phenyl]-pyrimidine-2,4-diamine are prepared from (2,5-Dichloro-pyrimidin-4-yl)-[2-ethanesulfonyl-phenyl]-amine and the corresponding aniline following the procedure of Example 7A



Exptl No.	Rx	R _f (solvent) or MS	NMR (400MHz), δ (ppm) or Retention time min. (HPLC)
27-1		0.53 (AcOEt)	CDCl ₃ : 1.28(t,3H), 3.12-3.19(m, 6H), 3.87-3.89(m, 7H), 6.45(dd, 1H), 6.53(d,1H), 7.24-7.28(m,1H), 7.31(s, 1H), 7.60-7.64(m, 1H), 7.95(dd, 1H), 8.04(d, 1H), 8.14(s, 1H), 8.58(d, 1H), 9.39(s, 1H)

27-2	 <chem>COC1=CC=C(C=C1)N2CCN(CC2)C3CCCCN3</chem>	585 (M+H)	2.38
27-3	 <chem>COC1=CC=C(C=C1)N2C=NC=N2</chem>	486 (M+H)	3.07
27-4	 <chem>COC1=CC=C(C=C1)N2CCN(CC2)C3CCOCC3</chem>	587 (M+H)	2.29
27-5	 <chem>CC(=O)N1CCN(CC1)C2=CC=C(OC)C=C2</chem>	545 (M+H)	2.59
27-6	 <chem>NC(=O)C1CCN(CC1)C2=CC=C(OC)C=C2</chem>	545 (M+H)	2.45

27-7		531 (M+H)	2.25
27-8		545 (M+H)	2.45
27-9		600 (M+H)	2.17

HPLC condition

Column: YMC CombiScreen ODS-A (5um, 12nm), 50 x 4.6 mm I.D.


Flow rate: 2.0 ml/min





Eluent: A) TFA/water (0.1/100), B) TFA/acetonitrile (0.1/100)






Gradient: 5-100%B (0-5min)





Detection: UV at 215nm





The following 5-Chloro-N²-(substituted phenyl)-N⁴-[2-(propane-2-sulfonyl)-phenyl]-pyrimidine-2,4-diamine are prepared from (2,5-Dichloro-pyrimidin-4-yl)-[2-(propane-2-sulfonyl)-phenyl]-amine and the corresponding aniline following the procedure of Example 7A











ExpIN o.	Rx	Rf (solvent) or MS	NMR (400MHz) , δ (ppm) or Retention time min. (HPLC)
28-1		0.2 (AcOEt)	CDCl ₃ : 1.31 (d, 6H), 1.85-1.73 (m, 1H), 1.86-1.98 (m, 3H), 2.62-2.70 (m, 1H), 3.11-3.13 (m, 2H), 3.21-8.28 (m, 1H), 3.28 (m, 2H), 3.88 8s, 3H), 5.41 (brs, 1H), 6.53 (d, 1H), 6.59 (d, 1H), 6.64 (brs, 1H), 7.28-7.34 (m, 1H), 7.34 (s, 1H), 7.60-7.67 (m, 1H), 7.91 (dd, 1H), 8.08 (d, 1H), 8.13 (s, 1H), 8.60 (d, 1H), 9.55 (s, 1H).
28-2		MS m/z 561, 563 (M+1).	CDCl ₃ : 1.31(d, 6H), 2.64 (t, 2H), 2.68-2.77 (m, 4H), 3.19(t, 4H), 3.17-3.28(m, 1H), 3.68(t, 2H), 3.88(s, 3H), 6.48(dd, 1H), 6.55 (d, 1H), 7.23-7.32(m, 1H), 7.62(ddd, 1H), 7.91(dd, 1H), 8.04(dd, 1H), 8.12(s, 1H), 8.60(d, 1H), 9.54(bs, 1H)
28-3		0.55 (AcOEt)	CDCl ₃ : 1.31(d, 6H), 3.12-3.14(m, 4H), 3.21-3.27(m, 1H), 3.87-3.89(m, 7H), 6.46(dd, 1H), 6.53(d,1H), 7.23-7.27(m, 1H), 7.30(s, 1H), 7.59-7.64(m, 1H), 7.91(dd, 1H), 8.05(d, 1H), 8.14(s, 1H), 8.60(d, 1H), 9.55(s, 1H)
28-4		0.37 (AcOEt)	CDCl ₃ : 1.32(d, 6H), 3.21-3.27(m, 1H), 4.00(s, 1H), 7.11(dd, 1H), 7.26-7.27(m, 1H), 7.29-7.33(m, 1H), 7.64(s, 1H), 7.66-7.71(m, 1H), 7.95(dd, 1H), 8.10(s, 1H), 8.21(s, 1H), 8.46(d, 1H), 8.50(s, 1H), 8.54(d, 1H), 9.59(s, 1H)






28-5		0.03 (AcOEt)	CDCl ₃ : 1.31(d, 6H), 1.67-1.77(m, 2H), 1.95-2.05 (m, 2H), 2.39-2.48 (m, 1H), 2.48-2.61(m, 2H), 2.63-2.78(m, 8H), 3.24 (sept, 1H), 3.71-3.63 (m, 2H), 3.87(s, 3H), 6.47(dd, 1H), 6.55 (d, 1H), 7.21-7.28(m, 1H), 7.61(ddd, 1H), 7.91(dd, 1H), 8.00(dd, 1H), 8.12(s, 1H), 8.60(d, 1H), 9.53(bs, 1H)
28-6		502 (M+H)	2.84
28-7		478 (M+H)	4.53
28-8		MS 599	CDCl ₃ : 1.31(d, 6H), 1.51-1.42(m, 2H), 1.67-1.53(m, 4H), 1.81-1.68(m, 2H), 1.96-1.89(m, 2H), 2.47-2.36(m, 1H), 2.57-2.54(m, 4H), 2.69(dd, 2H) 3.24(sept, 1H), 3.67(d, 1H), 3.87(s, 1H), 6.48 (dd, 1H), 6.56 (d, 1H), 7.31-7.21(m, 1H), 7.63-7.59 (m, 1H), 8.00(d, 1H), 8.12(s, 1H), 8.60(d, 1H), 9.55(s, 1H)
28-9		MS 585	CDCl ₃ : 1.26 (t, 3H), 1.31(d, 6H), 1.74-1.68(m, 2H), 1.85-1.76(m, 4H), 2.08-1.98(m, 2H), 2.19-2.10(m, 2H), 2.67-2.58(m, 4H), 2.79-2.72(m, 2H), 3.24(sept, 1H) 3.61(d, 2H), 3.87(s, 3H), 6.48(dd, 1H), 6.56 (d, 1H), 7.29-7.22 (m, 1H), 7.62(dd, 1H), 7.90(dd, 1H), 7.99(d, 1H), 8.12(s, 1H), 8.60(d, 1H), 9.53(s, 1H)


28-10		MS 559	CDCl ₃ : 1.31(d, 6H), 1.59-1.37(m, 2H), 1.81-1.69(m, 1H), 1.87(d, 2H), 2.73-2.67(m, 2H), 3.28-3.21(m, 1H), 3.37(s, 3H), 3.61(d, 1H), 3.87(s, 3H), 6.49(dd, 1H), 6.57(s, 1H), 7.31-7.21(m, 1H), 7.64-7.60 (m, 1H), 7.91(dd, 1H), 8.00(d, 1H), 8.60(d, 1H) 9.53(s, 1H)
28-11		MS 558	CDCl ₃ : 1.31(d, 6H), 2.15(s, 3H), 3.12(ddd, 4H), 3.24(sept, 1H), 3.64 (t, 2H), 3.80(t, 2H), 3.89(s, 3H), 6.47(dd, 1H), 6.55(d, 1H), 7.29-7.24(m, 1H), 7.33(bs, 1H), 7.62(m, 1H), 7.92(dd, 1H), 8.08(d, 1H), 8.14(s, 1H), 8.60(d, 1H) 9.55(s, 1H)
28-12		MS 544	CDCl ₃ : 1.16, (t, 3H), 1.31(d, 6H), 2.56-2.44(b, 2H), 2.71-2.60 (m, 4H), 3.28-3.17(m, 5H), 3.88(s, 3H), 6.48(dd, 1H), 6.58(d, 1H), 7.30-7.22(m, 1H), 7.63-7.58(m, 1H), 7.90(dd, 1H), 8.01(d, 1H), 8.12(s, 1H), 8.60(d, 1H) 9.54(s, 1H)
28-13		MS 601	CDCl ₃ : 1.31(d, 6H, J=6.55), 1.75-1.63(m, 2H), 2.00-1.91(m, 2H), 2.37-2.27(m, 1H), 2.60 (t, 4H, J=4.79), 2.74-2.59(m, 2H), 3.24(sept, 1H), 3.66 (d, 2H, J=12.1), 3.75(t, 4H, J=4.53), 3.88(s, 3H), 6.48(dd, 1H, J=2.52, 8.56), 6.56(d, 1H, J=2.52), 7.33-7.22(m, 1H), 7.64-7.59 (m, 1H), 7.91(dd, 1H, J=8.05, 1.51), 8.01(d, 1H, J=8.56), 8.12(s, 1H), 8.61(d, 1H, J=7.55) 9.54(s, 1H)


28-14		MS 559	CDCl ₃ : 1.11 (d, 6H, J=6.55), 1.31(d, 6H, J=7.05), 2.82-2.68(m, 5H), 3.20-3.17(m, 4H), 3.28-3.17(m, 1H), 3.87(s, 3H), 6.48(dd, 1H, J=2.52, 8.56), 6.56(d, 1H, J=2.52), 7.33-7.24(m, 1H), 7.62-7.58(m, 1H), 7.90(dd, 1H, J=), 8.01(d, 1H, J=8.56), 8.12(s, 1H), 8.60(d, 1H, J=8.56) 9.54(s, 1H)
28-15		MS 559	CDCl ₃ : 1.31 (d, 6H, J=7.05), 1.97-1.85(m, 2H), 2.17-1.98(m, 2H), 2.35-2.25(m, 1H), 2.75(m, 2H), 3.24(sept, 1H), 3.65(d, 2H), 3.88(s, 3H), 5.30 (bs, 1H), 5.48(bs, 1H), 6.48(dd, 1H, J=2.51, 8.56), 6.56(d, 1H, J=2.52), 7.33-7.21(m, 1H), 7.62 (m, 1H), 7.91(dd, 1H, J=1.51, 8.06), 8.03(dd, 1H, J=3.02, 8.56), 8.13(s, 1H), 8.60(d, 1H, J=8.57), 9.54(s, 1H)
28-16		MS 532	CDCl ₃ : 1.31 (d, 6H, J=7.06), 1.46-1.43(m, 1H), 1.79-1.68(m, 2H), 2.08-1.99(m, 2H), 2.99-2.88(m, 2H), 3.24(sept, 1H), 3.51-3.45(m, 2H), 3.91-3.80(m, 1H), 3.88(s, 3H), 6.49(dd, 1H, J=2.52, 8.56), 6.57(d, 1H, J=2.52), 7.34-7.23(m, 1H), 7.64-7.60(m, 1H), 7.91(dd, 1H, J=1.51, 8.06), 8.02(dd, 1H, J=3.02, 9.06), 8.13(s, 1H), 8.60(d, 1H, J=8.06) 9.53 (s, 1H)
28-17		MS 532	CDCl ₃ : 1.31 (d, 6H, J=6.96), 2.18-2.12(m, 2H), 3.24(sept, 1H), 3.37-3.32(m, 2H), 3.39(s, 3H), 3.43(d, 1H, J=8.56), 3.51(dd, 1H, J=5.04, 10.6), 3.87(s, 3H), 4.17-4.09 (m, 1H) 6.13 (dd, 1H, J=2.51, 8.56), 6.16(d, 1H, J=2.52), 7.09(bs, 1H), 7.31-7.21(m, 1H), 7.60-7.56 (m, 1H), 7.85(d, 1H, J=8.56), 7.89(dd, 1H, J=1.51, 8.06), 8.10(s, 1H), 8.65(d, 1H, J=9.06) 9.54 (s, 1H)




28-18		MS 546	CDCl ₃ : 1.31 (d, 6H, J=7.05), 1.82-1.70(m, 2H), 2.08-1.99(m, 2H), 2.96-2.87(m, 2H), 3.24(sept, 1H), 3.41-3.33(m, 1H), 3.40(s, 3H), 3.51-3.42(m, 2H), 3.87(s, 3H), 6.49(dd, 1H, J=2.52, 9.07), 6.57(d, 1H, J=2.52), 7.32-7.22(m, 1H), 7.64-7.60 (m, 1H), 7.91(dd, 1H, J=3.02, 9.06), 8.12(s, 1H), 8.60(d, 1H, J=8.56) 9.53 (s, 1H)
28-19		0.33 (AcOEt)	CDCl ₃ : 1.31 (d, 6H, J=7.05), 1.82-1.70(m, 2H), 2.08-1.99(m, 2H), 2.96-2.87(m, 2H), 3.24(sept, 1H), 3.41-3.33(m, 1H), 3.40(s, 3H), 3.51-3.42(m, 2H), 3.87(s, 3H), 6.49(dd, 1H, J=2.52, 9.07), 6.57(d, 1H, J=2.52), 7.32-7.22(m, 1H), 7.62(m, 1H), 7.91(dd, 1H, J=3.02, 9.06), 8.12(s, 1H), 8.60(d, 1H, J=8.56) 9.53 (s, 1H)
28-20		MS 544	CDCl ₃ : 1.31 (d, 6H), 1.66-1.53(m, 2H), 2.10-2.01(m, 2H), 2.51 (s, 3H), 2.70-2.13(m, 1H), 2.83-2.74(m, 2H), 3.24(Sept, 1H), 3.63-3.55(m, 2H), 3.87(s, 3H), 4.34-4.25(m, 1H), 6.48(dd, 1H), 6.56(d, 1H), 7.34-7.24(m, 1H), 7.64-7.60(m, 1H), 7.90(dd, 1H), 8.00(d, 1H), 8.12(s, 1H), 8.60(dd, 1H), 9.53(s, 1H)
28-21		MS 531	CDCl ₃ : 1.30 (s, 3H), 1.32 (s, 3H), 2.33-2.22(m, 1H), 2.54(s, 3H), 3.37-3.20(m, 3H), 3.57-3.44(m, 3H), 3.86(s, 3H), 6.12(dd, 1H), 6.16(d, 1H), 7.14-7.08(m, 1H), 7.30-7.20(m, 1H), 7.65-7.58(m, 1H), 7.93-7.87(m, 1H), 8.10(s, 1H), 8.64(d, 1H) 9.54 (s, 1H)


28-22		MS 545	CDCl ₃ : 1.30 (s, 3H), 1.32 (s, 3H), 2.03-1.89(m, 1H), 2.30-2.18(m, 1H), 2.34(s, 6H), 2.96-2.83(m, 1H), 3.29-3.16(m, 2H), 3.40-3.34(m, 1H), 3.53-3.43(m, 2H), 3.87(s, 3H), 6.11(dd, 1H,) 6.13(dd, 1H), 7.08(bs, 1H), 7.31-7.21(m, 1H), 7.60-7.56(m, 1H), 7.85(d, 1H), 7.89(dd, 1H), 8.10(s, 1H), 8.66(d, 1H) 9.54(s, 1H)
28-23		MS 545	CDCl ₃ : 1.31 (s, 3H), 1.32 (s, 3H), 3.05(s, 3H), 3.24(sept, 1H), 3.50-3.43(m, 4H), 3.85(s, 2H), 3.89(s, 3H), 6.11(dd, 1H,) 6.43(dd, 1H), 6.50(d, 1H), 7.31-7.28(m, 1H), 7.64-7.60(m, 1H), 7.92(dd, 1H), 8.09(d, 1H), 8.13(s, 1H), 8.58(d, 1H) 9.55(s, 1H)
28-24		0.05 (AcOEt/Me OH=4/1)	CDCl ₃ : 1.30 (s, 3H), 1.32 (s, 3H), 1.92-1.83(m, 1H), 2.17-1.95(m, 1H), 2.43-2.27(m, 2H), 2.79-2.71(m, 4H), 3.15-2.97(m, 4H), 3.23-3.16(m, 4H), 3.24(sept, 1H), 3.87(s, 3H), 6.11(dd, 1H,) 6.47(dd, 1H), 6.55(d, 1H), 7.33-7.23(m, 1H), 7.63-7.59(m, 1H), 7.95(dd, 1H), 8.01(dd, 1H), 8.12(s, 1H), 8.60(d, 1H) 9.54(s, 1H)
28-25		MS 600	CDCl ₃ : 1.30 (s, 3H), 1.32 (s, 3H), 1.80-1.70(m, 2H), 2.01-1.93(m, 2H), 2.49-2.28(m, 12H), 2.76-2.62(m, 4H), 3.04-2.96(m, 4H), 3.16-3.05(m, 2H), 3.24(sept, 1H), 3.72-3.63(m, 2H), 3.87(s, 3H), 6.48(dd, 1H), 6.55(d, 1H), 7.31-7.23(m, 1H), 7.66-7.589(m, 1H), 7.91(dd, 1H), 8.01(d, 1H), 8.12(s, 1H), 8.60(d, 1H) 9.53(s, 1H)

28-26		MS 573	CDCl ₃ : 1.30 (s, 3H), 1.32 (s, 3H), 2.59-2.43 (m, 4H), 2.78-2.73(m, 1H), 3.00-2.86(m, 2H), 3.38-3.20(m, 3H), 3.54-2.45(m, 1H), 3.73(dd, 1H), 3.84-3.77(m, 1H), 3.94-3.87(m, 1H), 3.88(s, 3H), 6.46(dd, 1H), 6.53(d, 1H), 7.32-7.23(m, 1H), 7.31 (bs, 1H), 7.63-7.52(m, 1H), 7.91(dd, 1H), 8.04(d, 1H), 8.13(s, 1H), 8.60(d, 1H) 9.54(s, 1H)
28-27		MS 559	CDCl ₃ : 1.30(s, 3H), 1.32 (s, 3H), 1.82-1.73 (m, 1H), 1.97-1.84(m, 3H), 2.73-2.51 (m, 1H), 3.12(t, 2H), 3.31-3.20 (m, 3H), 3.90(s, 3H), 5.46-5.37(m, 1H), 6.53(dd, 1H), 6.59(d, 1H), 6.68-6.62 (m, 1H), 7.28-7.21 (m, 1H), 7.33(bs, 1H), 7.65-7.61(m, 1H), 7.92(dd, 1H), 8.08(d, 1H), 8.14(s, 1H), 8.60(d, 1H), 9.55(s, 1H)
28-28		MS 559	CDCl ₃ : 1.30(s, 3H), 1.32 (s, 3H), 1.82-1.73 (m, 1H), 1.97-1.84(m, 3H), 2.73-2.51 (m, 1H), 3.12(t, 2H), 3.31-3.20 (m, 3H), 3.90(s, 3H), 5.46-5.37(m, 1H), 6.53(dd, 1H), 6.59(d, 1H), 6.68-6.62 (m, 1H), 7.28-7.21 (m, 1H), 7.33(bs, 1H), 7.65-7.61(m, 1H), 7.92(dd, 1H), 8.08(d, 1H), 8.14(s, 1H), 8.60(d, 1H), 9.55(s, 1H)
28-29		MS 413	CDCl ₃ : 1.31 (s, 3H), 1.33(s, 3H), 2.92(t, 4H), 3.28(sept, 1H) 3.73(t, 4H), 3.87(s, 3H), 6.51(dd, 1H), 6.82(d, 1H), 7.32-7.23 (m, 1H), 7.57(bs, 1H), 7.70-7.64(m, 1H), 7.92(dd, 1H), 8.01(bs, 1H), 8.12(s, 1H), 8.60(d, 1H), 9.53(s, 1H)
28-30		MS 493	CDCl ₃ : 1.30 (s, 3H), 1.33(s, 3H), 3.25(sept, 1H) 3.60 (bs, 3H), 3.89(s, 3H), 6.59(s, 1H), 7.27-7.18 (m, 1H), 7.61(dd, 1H), 7.83(bs, 1H), 7.90(dd, 1H), 8.15 (s, 1H), 8.55(d, 1H), 9.55(s, 1H)




28-31		MS 445	CDCl ₃ : 1.31(d, 6H), 1.59-1.37(m, 2H), 1.81-1.69(m, 1H), 1.87(d, 2H), 2.73-2.67(m, 2H), 3.28-3.21(m, 1H), 3.37(s, 3H), 3.61(d, 1H), 3.87(s, 3H), 6.49(dd, 1H), 7.025(bs, 1H), 7.28-7.23(m, 1H), 7.64-7.59(m, 1H), 7.93-7.89(m, 2H), 8.15(s, 1H), 8.57(dd, 1H) 9.56(s, 1H)
-------	---	-----------	---





Exptl No.	Rx	HPLC Retention time (min)	Mass (ESI) m/z
29-1		3.30	546 (M+H)
29-2		2.82	627 (M+H)
29-3		3.07	587 (M+H)





Expl No.	Rx	HPLC Retention time (min)	Mass (ESI) m/z
30-1	<p>Chemical structure of Rx for entry 30-1: 4-methoxy-N-(4-methylphenyl)morpholine.</p>	2.82	516 (M+H)
30-2	<p>Chemical structure of Rx for entry 30-2: 1-(4-methoxy-N-(4-methylphenyl)pyrrolidin-2-yl)propan-1-one.</p>	2.65	557 (M+H)
30-3	<p>Chemical structure of Rx for entry 30-3: 1-(4-methoxy-N-(4-methylphenyl)pyrrolidin-2-yl)propan-1-one.</p>	2.50	557 (M+H)
30-4	<p>Chemical structure of Rx for entry 30-4: 1-(4-methoxy-N-(4-methylphenyl)pyrrolidin-2-yl)propan-1-one.</p>	3.10	498 (M+H)



30-5		2.30	543 (M+H)
30-6		2.52	557 (M+H)
30-7		2.23	612 (M+H)




Expl No.	Rx	HPLC Retention time (min)	Mass (ESI) m/z
31-1		3.15	423 (M+H)

31-2		2.62	490 (M+H)
------	---	------	-----------




Exptl No.	Rx	MS	NMR (400MHz) in CDCl ₃ , δ (ppm)
32-1		585.3	1.03 (s, 3H), 1.04(s, 3H), 2.15(s, 3H), 2.32(sept, 1H) 3.00(d, 2H) 3.10(t, 2H), 3.13(t, 2H), 3.64(t, 2H), 3.79(t, 2H), 3.89(s, 3H), 6.45(dd, 1H), 6.55(d, 1H), 7.34-7.26 (m, 1H), 7.52(bs, 1H), 7.64-7.60(m, 1H), 7.97(dd, 1H), 8.07(d, 1H), 8.15(s, 1H), 8.54(d, 1H), 9.32(s, 1H)
32-2		532 (M+H)	3.17






- 113 -






Expl No.	Rx	MS	NMR (400MHz) in CDCl ₃ , δ (ppm)
33-1		585.3	1.66-1.52 (m, 2H), 1.92-1.73 (m, 4H), 2.12-2.03 (m, 2H), 2.15(s, 3H), 3.00(d, 2H) 3.11(t, 2H), 3.14(t, 2H), 3.58-3.46(m, 1H), 3.64 (t, 2H), 3.80(t, 2H), 3.89(s, 3H), 6.48(dd, 1H), 6.55(d, 1H), 7.30-7.24 (m, 1H), 7.52(bs, 1H), 7.63-7.58(m, 1H), 7.94(dd, 1H), 8.08(d, 1H), 8.14(s, 1H), 8.60(d, 1H), 9.54(s, 1H)
33-2		544 (M+H)	3.15





Expl No.	Rx	HPLC Retention time (min)	Mass (ESI) m/z
34-1		3.15	532 (M+H)

34-2		3.34	558 (M+H)
34-3		3.35	546 (M+H)
34-4		3.32	546 (M+H)
34-5		3.09	566 (M+H)
34-6		2.87	552 (M+H)


Ex No		MS	NMR (400MHz), CDCl ₃ , δ ppm
34-7		MS 435, 436	1.05 (t, 3H), 1.69-1.78 (m, 2H), 2.86-2.95 (m, 1H), 3.16-3.25 (m, 1H), 6.57-6.68 (m, 2H), 7.17 (dd, 1H), 7.35-7.39 (m, 1H), 7.50 (dd, 1H), 8.13 (s, 1H), 8.16-8.21 (m, 1H), 8.48 (d, 1H), 10.14 (s, 1H)
34-8		MS 549, 551	0.94 (t, 3H), 1.69-1.80 (m, 2H), 2.38 (s, 3H), 2.55-2.64 (m, 4H), 3.02-3.08 (m, 2H), 3.22-3.29 (m, 4H), 3.88 (s, 3H), 6.55 (ddd, 1H), 6.60-6.66 (m, 1H), 7.13-7.18 (m, 1H), 7.34 (br.s, 1H), 7.44 (d, 1H), 8.10 (s, 1H), 8.10-8.23 (m, 2H), 8.88 (s, 1H).



35-1		567 [M+1] ⁺	DMSO-d ₆ : 2.24 (s, 3H), 2.45-2.50 (m, 4H), 2.78 (d, 3H), 3.10-3.17 (m, 8H), 3.74-3.79 (m, 7H), 6.49 (dd, 1H), 6.66 (d, 1H), 6.85-6.89 (m, 1H), 7.18 (d, 1H), 7.40 (d, 1H), 7.98-8.02 (m, 2H), 8.29 (br.d, 1H), 8.60-8.66 (m, 1H), 11.17 (s, 1H).
35-2		505 [M+1] ⁺	DMSO-d ₆ : 2.24(s, 3H), 2.46-2.50(m, 4H), 2.79(d, 3H), 3.13-3.17(m, 4H), 3.78(s, 3H), 6.69(d, 1H), 6.87(dd, 1H), 7.07-7.17(m, 2H), 7.19-7.23(m, 2H), 7.54(d, 1H), 8.13(s, 1H), 8.45(s, 1H), 8.65-8.75(m, 1H), 9.04(s, 1H), 11.19(s, 1H)

Example 36 (Intermediates for Left anilines)

36-1 Preparation of 2-amino-N-methyl-benzamide




To a suspension of 16.3 g (100 mmol) of isatoic anhydride in 100mL of H₂O is added portionwise 100mL of 2N methylamine - tetrahydrofuran solution (200 mmol) at room temperature. The reaction mixture is stirred for 1 hour and then extracted with AcOEt. The organic layer is washed with H₂O and brine, dried over Na₂SO₄, and concentrated under

reduced pressure to give 13.79 g of desired product, 2-amino-*N*-methyl-benzamide (92 mmol, 92%) as colorless solid.

NMR (400MHz, CDCl₃, δ): 2.97 (d, 3H, J = 4.52 Hz), 5.49 (bs, 1H), 6.07 (bs, 1H), 6.64 (ddd, 1H, J = 8.04, 7.56, 1.0 Hz), 6.68 (dd, 1H, J = 8.32, 1.0 Hz), 7.20 (ddd, 1H, J = 8.32, 7.56, 1.52 Hz), 7.29 (dd, 1H, J = 8.04, 1.52 Hz).

36-2

2-(2,5-Dichloro-pyrimidin-4-ylamino)-*N*-methyl-benzamide




To a solution of 15.0 g (99.8 mmol) of 2-amino-*N*-methyl-benzamide in DMF (300mL) are added 2, 4, 5-trichloropyrimidine (23.8 g, 130 mmol) and potassium carbonate (17.9 g, 130 mmol). The reaction mixture is stirred at 75°C for 5 hours, cooled to room temperature, and then poured into H₂O (600mL). The resulting precipitate is collected by a filtration followed by washing with 50% aqueous CH₃CN (200mL) and dried under reduced pressure (40°C, 10 hours) to give desired 2-(2,5-dichloro-pyrimidin-4-yl-amino)-*N*-methyl-benzamide as ivory solid (26.4 g, 88.9 mmol, 89%).

NMR (400MHz, DMSO-d₆, δ): 2.81 (d, 3H, J = 4.52 Hz), 7.22 (dd, 1H, J = 8.56, 8.04 Hz), 7.60 (ddd, 1H, J = 8.56, 8.56, 1.0 Hz), 7.81 (dd, 1H, J = 8.04, 1.0 Hz), 8.48 (s, 1H), 8.52 (d, 1H, J = 8.56 Hz) 8.80-8.90 (m, 1H), 12.18 (s, 1H).


According the manner described above, the following compounds are prepared.

36-3


2-(5-Bromo-2-chloro-pyrimidin-4-ylamino)-*N*-methyl-benzamide



NMR (400MHz, DMSO- d_6 , δ): 2.81(d, 3H), 7.23(ddd, 1H, $J=7.54, 7.54, 1.0\text{Hz}$), 7.59(ddd, 1H, $J=7.93, 8.06, 1.52\text{Hz}$), 7.79(dd, 1H, $J=7.8, 1.52\text{Hz}$), 8.47(dd, 1H $J=8.06, 1.0\text{Hz}$), 8.55(s, 1H), 8.81-8.87(m, 1H), 12.0(brs, 1H). Rf: 0.46 (n-Hexane: AcOEt=7:3).

36-42-(2,5-Dichloro-pyrimidin-4-ylamino)-N-ethyl-benzamide


NMR (400MHz, CDCl_3 , δ): 1.28 (t, $d=7.04$, 3H), 3.48-3.57 (m, 2H), 6.22 (br. s, 1H), 7.11-7.17 (m, 1H), 7.51 (dd, $J=1.0, 8.04$, 1H), 7.53-7.61 (m, 1H), 8.22 (s, 1H), 8.69-8.74 (m, 1H), 11.66 (br. s, 1H). Rf : 0.60 (Hexane :AcOEt=1 :1).

36-5Preparation of 2-(5-bromo-2-chloro-pyrimidin-4-ylamino)-N-methyl-benzenesulfonamide


A suspension of 5-bromo-2,4-dichloropyrimidine (684 mg, 3.0 mmol) and 2-amino-N-methyl-benzenesulfonamide (559 mg, 3.0 mmol) in N,N-dimethylformamide (10 mL) containing potassium carbonate (830 mg, 6.0 mmol) is stirred at room temperature for 23 hours. Saturated aqueous ammonium chloride is added and the mixture is poured into water and extracted twice with ethyl acetate. The organic layer is washed with brine, dried over sodium sulfate, and evaporated in vacuo. The residue is purified by silica gel column chromatography (n-hexane - ethyl acetate gradient) to afford the title compound as a slightly yellow solid.

$^1\text{H-NMR}$ (CDCl_3 , δ (ppm)): 2.67 (d, 3H), 4.79 (q, 1H), 7.26 (s, 1H), 7.29 (ddd, 1H), 7.66 (ddd, 1H), 7.95 (dd, 1H), 8.37 (s, 1H), 8.48 (d, 1H), 9.52 (s, 1H). Rf (n-hexane : ethyl acetate = 10:3): 0.33.

According to the manner described above, the following compound is prepared.

36-62-(2,5-Dichloro-pyrimidin-4-ylamino)-N-methyl-benzenesulfonamide


$^1\text{H-NMR}$ (400MHz, CDCl_3 , δ); 2.67(d, 3H), 4.97-5.04(m, 1H), 7.29(ddd, 1H, $J=7.54$, 7.54, 1.0Hz), 7.66(ddd, 1H, $J=7.93$, 8.08, 1.48Hz), 7.94(dd, 1H, $J=8.04$, 1.52Hz), 8.24(s, 1H), 8.51(dd, 1H $J=8.06$, 1.0Hz), 9.64(brs, 1H). Rf: 0.45 (n-Hexane: AcOEt=4:1).

36-72-(2,5-Dichloro-pyrimidin-4-ylamino)-N-isopropyl-benzenesulfonamide

To a solution of 2-amino-N-isopropyl-benzenesulfonamide (16.1g, 75.1mmol) in DMI (150mL) is added sodium hydride (6.6g, 165.3mmol) portionwise at 0°C . After the mixture is stirred at room temperature for one hour, 2, 4, 5-trichloropyrimidine (20.7g, 112.7mmol) is added at 0°C . After further stirring at room temperature for 5 hrs, water is added and the mixture is extracted with AcOEt three times. Organic layer is washed with brine, dried over sodium sulfate and evaporated under reduced pressure. The residue is purified by silica gel column chromatography (Hexane to Hexane:AcOEt=4:1) to afford the title compound as pale brown solid (10.2g, 38%).

$^1\text{H-NMR}$ (400MHz, CDCl_3 , δ); 1.06(d, 6H), 3.43-3.53(m, 1H), 4.38(d, 1H), 7.29(dd, 1H), 7.66(dd, 1H), 7.98(d, 1H), 8.29(s, 1H), 8.51(d, 1H), 9.51(brs, 1H). Rf: 0.45 (n-Hexane:AcOEt=4:1)


The following compounds are prepared in the same manner described above.



Exptl No.	Rz	Rf (solvent) or MS	NMR (400MHz) , δ (ppm)
36-8		0.45 (n-Hexane: AcOEt=4:1)	DMSO-d ₆ ; 0.63(t, 6H), 0.86(d, 3H), 1.21-1.31(m, 2H), 3.02-3.12(m, 1H), 7.37(dd, 1H), 7.71(dd, 1H), 7.85(d, 1H), 7.89(d, 1H), 8.20(d, 1H), 8.56(s, 1H), 9.51(brs, 1H)
36-9		0.46 (n-Hexane: AcOEt=7:3)	CDCl ₃ ; 0.70(t, 6H), 1.23-1.45(m, 4H), 3.03-3.13(m, 1H), 4.27(d, 1H), 7.27(dd, 1H), 7.65(dd, 1H), 7.98(d, 1H), 8.29(s, 1H), 8.52(d, 1H), 9.59(brs, 1H)

36-10

Preparation of 2-(2-chloro-5-nitro-pyrimidin-4-ylamino)-N-methyl-benzenesulfonamide:




2,4-Dichloro-5-nitro-pyrimidine (1.94 g, 10 mmol) and 2-amino-N-methyl-benzenesulfonamide (1.86 g, 10 mmol) are dissolved in CHCl₃ (30 mL). The reaction mixture is heated at 61°C for 2 h. The solvent is evaporated and the residue is washed with ether to give the title product.

Rf = 0.5 (n-hexane : ethyl acetate = 1:1). ¹H-NMR (400MHz, CDCl₃), δ (ppm): 2.67 (d, 3H), 4.6-4.7 (m, 2H), 7.41 (t, 1H), 7.7 (t, 1H), 8.04 (d, 1H), 8.15 (d, 1H), 9.21 (s, 1H), 11.2 (s, 1H).

36-11

Preparation of (2,5-Dichloro-pyrimidin-4-yl)-[2-(propane-1-sulfonyl)-phenyl]-amine


- 120 -





To a solution of 2-(Propane-1-sulfonyl)-phenylamine (3.69g, 18.5 mmol) of N,N-dimethylformamide (40mL), sodium hydride (1.48g, 37 mmol) is added portionwise at 0°C. After stirring, 2,4,5-trichloropyrimidine (2.1mL, 18.5 mmol) is added. The mixture is stirred at 0°C for 30 minutes and is further stirred at room temperature for 7hrs. After adding saturated aqueous ammonium chloride, the mixture is poured into water and extracted twice with ethyl acetate. The organic layer is washed with brine, dried over sodium sulfate, and evaporated in vacuo. The residue is purified by silica gel column chromatography (n-hexane - ethyl acetate gradient) to afford the title compound as colorless solids.

¹H-NMR (CDCl₃), δ (ppm): 0.99 (t, 3H), 1.77 (d, 2H), 3.07-3.11 (m, 2H), 7.26 (s, 1H), 7.32 (ddd, 1H), 7.73 (ddd, 1H), 7.95 (dd, 1H), 8.31 (s, 1H), 8.61 (dd, 1H), 9.94 (bs, 1H). R_f (n-hexane : ethyl acetate = 3:1): 0.63

According to the manner described above, the following compounds are prepared.



Exptl No.	Rx	Identification
36-12		¹ H-NMR (CDCl ₃), δ (ppm): 1.35(d, 6H), 3.18-3.24(m, 1H), 7.30-7.34(m, 1H), 7.70-7.75(m, 1H), 7.92(dd, 1H), 8.30(s, 1H), 8.63(d, 1H), 10.06(s, 1H). R _f 0.70: (AcOEt)
36-13		NMR (400MHz) in CDCl ₃ , δ (ppm): 1.29(t, 3H), 3.15(q, 1H), 7.31-7.35(m, 1H), 7.71-7.75(m, 1H), 7.96(dd,

		1H), 8.31(s, 1H), 8.60(d, 1H), 9.92(s, 1H). Rf: 0.67 (AcOEt).
36-14		1.01-1.06(m, 2H), 1.32-1.37(m, 2H), 2.49-2.55(m, 1H), 7.29-7.33(m, 1H), 7.69-7.73(m, 1H), 7.91(dd, 1H), 8.31(s, 1H), 8.58(d, 1H), 9.90(s, 1H). Rf 0.69 (AcOEt)
36-15		0.99(t, 6H), 1.72-1.90(m, 4H), 2.76-2.82(m, 1H), 7.26-7.34(m, 1H), 7.69-7.74(m, 1H), 7.92(dd, 1H), 8.30(s, 1H), 8.62(d, 1H), 10.02(s, 1H). Rf: 0.73 (AcOEt)

Example 36-16

Synthesis of substituted amines which are commercially not available:

Preparation of 3-amino-4'-methoxy-4-methylbiphenyl

To a solution of 4-methoxyphenyl-boronic acid (500 mg, 3.29 mmol) in toluene (5.2 mL) and ethanol (1.3 mL), potassium carbonate (910 mg, 6.58 mmol), tetrakis(triphenylphosphine)-palladium (228.1 mg, 0.099 mmol) and 4-bromo-1-methyl-2-nitrobenzene (711 mg, 3.29 mmol) are added and stirred at 100°C for 7 hours. The mixture is poured into water and extracted with ethyl acetate two times. The organic layer is washed with water and then brine, dried over magnesium sulfate, and evaporated in vacuo. The residue is purified with silica gel column chromatography (n-hexane : ethyl acetate = 5 : 1) to afford the 4'-methoxy-4-methyl-3-nitro-biphenyl as a yellow solid.

¹H-NMR (δ, ppm) : 2.62 (s, 3H), 3.86 (s, 3H), 7.02-6.98 (m, 2H), 7.37 (d, 1H), 7.54 (dd, 2H), 7.68 (dd, 1H), 8.18 (d, 1H). Rf (hexane : ethyl acetate = 3:1): 0.40.

A suspension of 4'-methoxy-4-methyl-3-nitrobiphenyl (630 mg, 2.95 mmol) and 10% palladium on charcoal (63 mg, 0.059 mmol) in methanol (6 mL) is stirred under hydrogen atmosphere for 12 hours. Palladium catalyst is removed by filtration and the resulting solution is evaporated in vacuo to afford the title compound.

¹H-NMR (δ, ppm) : 2.20 (s, 3H), 3.84 (s, 3H), 6.87 (d, 1H), 6.89 (dd, 1H), 6.95 (d, 2H), 7.09 (d, 1H), 7.48 (d, 2H). Rf (n-hexane : ethyl acetate = 1:1): 0.50.

Preparation of 4-(3-amino-4-methylbenzoyl)-piperazine-1-carboxylic acid tert-butyl ester

To a solution of 4-methyl-3-nitro-benzoic acid (300 mg, 2.76 mmol), N-butoxycarbonyl-piperazine (340 mg, 1.83 mmol) in DMF (3.0 mL), triethylamine (300 μ L, 3.59 mmol), TBTU (800 mg, 2.49 mmol) and HOAt (270.5 mg, 1.99 mmol) are added and stirred at room temperature for 24 hours. The mixture is poured into water and extracted twice with ethyl acetate. The organic layer is washed with water and then brine, dried over magnesium sulfate, and evaporated in vacuo. The residue is purified with silica gel column chromatography (n-hexane : ethyl acetate = 5 : 1) to afford 4-(4-methyl-3-nitrobenzoyl)-piperazine-1-carboxylic acid tert-butyl ester as a colorless solid.

$^1\text{H-NMR}$ (δ , ppm) : 1.47 (s, 9H), 2.64 (s, 3H), 3.28-3.88 (m, 8H), 7.42 (d, 1H), 7.56 (dd, 1H), 8.03 (d, 1H). Rf (hexane : ethyl acetate = 10:1): 0.13.

The title compound is obtained by reduction with hydrogen over 10% palladium on charcoal in methanol solution.

Preparation of 4-(3-amino-4-methylphenyl)-morpholine

To a solution of 4-bromo-1-methyl-2-nitrobenzene (225 mg, 1.04 mmol), morpholine (125 μ L, 1.25 mmol), and cesium carbonate (474.4 mg, 1.46 mmol) in toluene, palladium diacetate (31.2 mg, 0.139 mmol) and 2-(di-t-butylphosphino)biphenyl (125 mg, 0.403 mmol) are added and stirred at 100°C for 5 hours. After cooling, the mixture is filtered to remove insoluble material. The filtrate is poured into water and extracted with ethyl acetate twice. The organic layer is washed with water and then brine, dried over magnesium sulfate, and evaporated in vacuo. The residue is purified with silica gel column chromatography (n-hexane : ethyl acetate = 5 : 1) to afford 4-(4-methyl-3-nitrophenyl)-morpholine as a yellow solid.


$^1\text{H-NMR}$ (δ , ppm) : 2.50 (s, 3H), 3.17-3.19 (m, 4H), 3.86-3.88 (m, 4H), 7.04 (dd, 1H), 7.21 (d, 1H), 7.47 (d, 1H). Rf (hexane : ethyl acetate = 5:1): 0.20.

The title compound is obtained by reduction with hydrogen over 10% palladium on charcoal in methanol solution.

Example 37: Synthesis of substituted amines which are commercially not available:

37-1

Preparation of 1-(3-Methoxy-4-nitro-phenyl)-piperdin-4-ol











To a suspension of piperidin-4-ol (2.79g, 28 mmol) and potassium carbonate (3.88 g, 28 mmol) in N,N-dimethylformamide (40 mL), 4-Fluoro-2-methoxy-1-nitro-benzene (4.0g, 23 mmol) is added and stirred at room temperature for 24 hours. The mixture is poured into water and the precipitate is collected by a filtration. The resulting solid is dried in vacuo at 50°C to afford 1-(3-methoxy-4-nitro-phenyl)-piperidin-4-ol (5.23g) as yellow solids in 89% yield.





¹H-NMR (400MHz, CDCl₃, δ, ppm) : 1.54(d, 1H), 1.62-1.71(m, 2H), 1.98-2.04(m, 2H), 3.22(ddd, 4H), 3.73-3.80(m, 2H), 3.95(s, 3H), 3.98-4.02(m, 1H), 6.33(d, 1H), 6.43(dd, 1H), 8.00(d, 1H).





By repeating the procedures described above using appropriate starting materials and conditions the following compounds are obtained.





Ex-No	Rx	Identification
37-2		¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 1.53-1.72(m, 2H), 1.80-1.83(m, 4H), 1.99-2.04(m, 2H), 2.24-2.31(m, 1H), 2.54-2.67(m, 4H), 3.03(dt, 2H), 3.84-3.89 (m, 2H), 3.95(s, 3H), 6.31(d, 1H), 6.42(dd, 1H), 8.01(d, 1H). Rf 0.54 (AcOEt)
37-3		¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 1.81-1.91(m, 2H), 1.99-2.04(m, 2H), 2.38-2.48(m, 1H), 3.03(ddd, 2H), 3.91-3.96(m, 2H), 3.95(s, 3H), 5.22-5.41(m, 1H), 5.40-5.53(m, 1H), 6.36(d, 1H), 6.43(dd, 1H), 8.00(d, 1H). Rf 0.15 (AcOEt)

37-4	 <p>Ethyl-[1-(3-methoxy-4-nitro-phenyl)-pyrrolidin-3-yl]-amine</p>	¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 1.15(t, 3H), 1.88-1.96(m, 1H), 2.22-2.30(m, 1H), 2.68-2.77(m, 2H), 3.15-3.18(m, 1H), 3.38-3.44(m, 1H), 3.52-3.62(m, 2H), 3.93(s, 3H), 5.92(d, 1H), 6.07-6.10(m, 1H), 8.00-8.02(m, 1H). R _f 0.65 (<i>n</i> -hexane: AcOEt=1:1).
37-5	 <p>1-(3-Methoxy-4-nitro-phenyl)-4-methyl-piperazine</p>	¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 2.36(s, 3H), 2.52-2.57(m, 4H), 3.40-3.43(m, 4H), 3.95(s, 3H), 6.32(d, 1H, <i>J</i> =2.52Hz), 6.43(dd, 1H, <i>J</i> =9.56, 2.52Hz), 7.99(d, 1H, <i>J</i> =9.08Hz). R _f 0.60 (MeOH : CH ₂ Cl ₂ =4:1).
37-6	 <p>3-(3-Methoxy-4-nitro-phenoxy)methyl-1-methyl-piperidine</p>	¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 1.10-1.19(m, 1H), 1.59-2.18(m, 6H), 2.28(s, 3H), 2.71-2.74(m, 1H), 2.88-2.91(m, 1H), 3.86-3.95 (m, 5H), 6.47-6.52(m, 2H), 7.97-8.00(m, 1H). R _f 0.65 (<i>n</i> -hexane: AcOEt=1:1)
37-7		¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 4.08(s, 3H), 7.30(dd, 1H), 7.58(d, 1H), 8.05(d, 1H), 8.15(s, 1H), 8.67(s, 1H). R _f 0.42 (AcOEt)





37-8		¹ H-NMR (400MHz, CDCl ₃ , δ , ppm): 1.40 – 1.50 (m, 2H), 1.55 – 1.69 (m, 6H), 1.90 – 1.96 (m, 2H), 2.45 – 2.53 (m, 5H), 2.90 – 2.99 (m, 2H), 3.90 – 4.00 (m, 2H), 3.94 (s, 3H), 6.30 (d, 1H, J = 2.5 Hz), 6.41 (dd, 1H, J = 9.0, 2.5 Hz), 7.99 (d, 1H, J = 9.0 Hz)
37-9		¹ H-NMR (400MHz, DMSO-d ₆ , δ , ppm) : 1.95-1.82(m, 2H), 2.15-2.06 (m, 1H), 2.30 (s, 3H), 3.17 (dd, 1H), 3.32-3.23 (m, 1H), 3.56-3.34 (m, 3H), 3.96 (s, 1H), 6.09 (d, 1H), 6.21 (dd, 1H), 7.91 (d, 1H)
37-10		¹ H-NMR (400MHz, CDCl ₃ , δ , ppm): 2.30 - 2.48 (m, 3H), 2.59 - 2.66 (m, 1H), 2.70 - 2.76 (m, 1H), 2.85 - 2.92 (m, 1H), 3.09 - 3.17 (m, 1H), 3.30 - 3.34 (m, 1H), 3.52 - 3.58 (m, 1H), 3.68 - 3.84 (m, 3H), 3.87 - 3.91 (m, 1H), 3.96 (s, 3H), 6.32 (d, 1H, J = 2.5 Hz), 6.42 (dd, 1H, J = 9.6, 2.5 Hz), 8.00 (d, 1H, J = 9.6 Hz)
37-11		¹ H-NMR (400MHz, DMSO-d ₆ , CDCl ₃ , δ , ppm) : 1.90-1.79(m, 1H), 2.25-2.15 (m, 1H), 2.21 (s, 3H), 2.87-2.77 (m, 1H), 3.16 (dd, 1H), 3.42-3.32 (m, 1H), 3.59-3.52 (m, 1H), 3.67-3.61 (m, 1H), 3.91 (s, 3H), 6.13 (d, 1H), 6.24 (dd, 1H), 7.91 (dd, 1H)





37-12		¹ H-NMR (400MHz, CDCl ₃): 1.43-1.00(m, 2H), 1.95-1.81 (m, 2H), 2.94-2.17(m, 2H), 2.96(s, 3H), 3.27 (d, 2H), 3.35(s, 3H), 3.97-3.90 (m, 2H), 3.95(s, 3H), 6.30(d, 1H), 6.42(dd, 1H) 8.00(d, 1H). Rf: 0.25 (AcOEt)
37-13		¹ H-NMR (400MHz, CDCl ₃): 1.14(t, 3H), 2.48(dd, 2H), 2.59(t, 4H), 3.42 (t, 4H), 3.95(s, 3H), 6.32(d, 1H), 6.43(dd, 1H) 8.01(d, 1H). Rf 0.15 (AcOEt)
37-14		¹ H-NMR (400MHz, CDCl ₃): 1.02-0.89 (m, 2H), 2.01-1.94 (m, 2H), 2.52-2.38 (m, 1H), 2.65-2.53 (m, 4H), 3.04-2.94(m, 2H), 3.79-3.69(m, 4H), 3.97-3.88 (m, 2H), 3.95(s, 3H), 6.32(d, 1H), 6.42(dd, 1H) 8.00(d, 1H). Rf 0.10 (AcOEt)
37-15		¹ H-NMR (400MHz, CDCl ₃): 1.08 (s, 3H), 1.09(s, 3H), 2.66(t, 4H), 2.74 (sept, 1H), 3.41 (t, 4H), 3.95(s, 3H), 6.32(d, 1H), 6.42(dd, 1H) 8.00(d, 1H). Rf 0.15 (AcOEt)





37-16		¹ H-NMR (400MHz, CDCl ₃): 1.91-1.81 (m, 2H), 2.06-1.97(m, 2H), 2.48-2.40(m, 1H), 3.07-2.98(m, 2H), 3.97-3.93(m, 2H), 3.93(s, 3H), 5.37-5.30(m, 1H), 5.55-5.43 (m, 1H), 6.33(d, 1H), 6.43(dd, 1H) 8.00(d, 1H). Rf 0.10 (AcOEt)
37-17		¹ H-NMR (400MHz, CDCl ₃): 2.18-2.07 (m, 1H), 2.30-2.22 (m, 1H), 3.38(s, 3H), 3.56-3.44(m, 4H), 3.95 (s, 3H), 4.13 (ddd, 1H), 5.96(d, 1H), 6.12(dd, 1H) 8.03(d, 1H). Rf 0.30 (AcOEt)
37-18		¹ H-NMR (400MHz, CDCl ₃): 1.46(s, 9H), 1.81-1.68(m, 4H), 2.73(bs, 3H), 3.07-2.97(m, 2H), 3.95(s, 3H), 4.03-3.94 (m, 2H), 6.32(d, 1H), 6.43(dd, 1H) 8.00(d, 1H). Rf 0.55 (Hexane:AcOEt)
37-19		¹ H-NMR (400MHz, CDCl ₃): 3.60-3.57(m, 2H), 3.68-3.65(m, 2H), 3.97(s, 3H), 4.07(s, 2H), 6.17(bs, 1H), 6.26(d, 1H), 6.39(dd, 1H) 8.04(d, 1H). Rf 0.85 (AcOEt)






37-20		¹ H-NMR (400MHz, CDCl ₃): 3.08(s, 3H), 3.54(dd, 2H), 3.67(dd, 2H), 3.96 (s, 3H), 4.05(s, 2H), 6.25(d, 1H), 6.38(dd, 1H) 8.03(d, 1H) . Rf 0.30 (AcOEt)
37-21		¹ H-NMR (400MHz, CDCl ₃): 1.73-1.55 (m, 2H), 1.99-1.91 (m, 2H), 2.09(s, 3H), 2.61-2.49 (m, 5H), 3.47(t, 2H), 3.63(t, 2H), 3.99-3.89 (m, 3H), 3.95 (s, 3H), 6.32(d, 1H), 6.42(dd, 1H) 8.01(d, 1H) . Rf 0.10 (AcOEt:MeOH=4:1)
37-22		¹ H-NMR (400MHz, CDCl ₃): 3.90(s, 3H), 3.98(s, 3H), 3.98 (s, 3H), 6.56(s, 1H), 7.59(s, 1H) . Rf 0.605 (AcOEt)
37-23		¹ H-NMR (400MHz, CDCl ₃): 3.25-3.22 (m, 4H), 3.90-3.87 (m, 4H), 3.95(s, 3H), 6.48(s, 1H), 7.57(s, 1H) . Rf 0.060 (Hexane:AcOEt=5:1)







37-24		¹ H-NMR (400MHz, CDCl ₃): 2.37 (s, 3H), 2.61 (bs, 4H), 3.27 (bs, 4H), 3.88 (s, 3H), 3.95(s, 3H), 6.48(s, 1H), 7.56(s, 1H) . Rf 0.10 (AcOEt:MeOH=5:1)
37-25		¹ H-NMR (400MHz, CDCl ₃): 1.09(t, 3H), 1.89(dd, 2H), 2.36(s, 3H), 2.55(t, 4H), 3.39(t, 4H), 4.03(t, 2H), 6.32(d, 1H), 6.42(dd, 1H), 7.98(d, 1H) . Rf 0.12 (AcOEt:MeOH=9:1)
37-26		¹ H-NMR (400MHz, CDCl ₃): 1.36(s, 3H), 1.38 (s, 3H), 2.10 (s, 2H), 2.17(s, 3H), 3.27-2.96 (m, 2H), 3.71 (d, 2H), 3.96 (s, 3H), 6.33(d, 1H), 6.43(dd, 1H), 8.02(d, 1H) . Rf 0.10 (AcOEt)
37-27		¹ H-NMR (400MHz, CDCl ₃): 1.16(s, 3H), 1.18 (s, 3H), 2.50(dd, 2H), 3.02-2.47 (m, 2H), 3.69 (dd, 2H), 3.96 (s, 3H), 6.31(d, 1H), 6.43(dd, 1H), 8.00(d, 1H) . Rf 0.070(AcOEt)
37-28		¹ H-NMR (400MHz, CDCl ₃): 1.16(d, 3H), 2.57(dd, 1H), 3.00-2.89 (m, 4H), 3.18-3.11 (m, 1H), 3.75-3.68 (m, 2H), 3.96 (s, 3H), 6.31(d, 1H), 6.43(dd, 1H), 8.01(d, 1H) . Rf 0.070 (AcOEt)






37-29		¹ H-NMR (400MHz, CDCl ₃): 1.18(t, 3H), 2.40(dd, 2H), 3.47-3.38(m, 4H), 3.71-3.63(m, 2H), 3.85-3.79(m, 2H), 3.96(s, 3H), 6.32(d, 1H), 6.42(dd, 1H), 8.01(d, 1H) . Rf 0.20 (AcOEt)
37-30		¹ H-NMR (400MHz, CDCl ₃): 1.16(s, 3H), 1.18(s, 3H), 2.82(sept, 1H), 3.50-3.37(m, 4H), 3.77-3.65(m, 2H), 3.86-3.78(m, 2H), 3.96(s, 3H), 6.33(d, 1H), 6.43(dd, 1H), 8.01(d, 1H) . Rf 0.48 (AcOEt)
37-31		¹ H-NMR (400MHz, CDCl ₃): 2.86(d, 3H), 3.48-3.45(m, 4H), 3.61-3.58(m, 4H), 3.96(s, 3H), 4.48-4.37 (m, 1H), 6.29(d, 1H), 6.40(dd, 1H), 8.01(d, 1H) . Rf 0.20 (AcOEt)
37-32		¹ H-NMR (400MHz, CDCl ₃): 1.72-1.60(m, 2H), 2.06-1.97(m, 2H), 3.25-3.17 (d, 3H), 3.78-3.70(m, 2H), 3.95(s, 3H), 4.04-3.99(m, 1H), 6.33(d, 1H), 6.43(dd, 1H), 8.00(d, 1H) . Rf 0.20 (AcOEt)


37-33		¹ H-NMR (400MHz, CDCl ₃): 1.53 (s, 6H), 2.14(s, 3H), 3.50(s, 2H), 3.61-3.58(m, 2H), 3.97-3.81(m, 2H), 3.97 (s, 3H), 6.10 (d, 1H), 6.26(dd, 1H), 8.05(d, 1H) . Rf 0.030 (AcOEt)
37-34		¹ H-NMR (400MHz, CDCl ₃): 2.54-2.23 (m, 4H), 2.67 (t, 2H), 3.29-3.23(m, 2H), 3.74(t, 4H), 3.94(s, 3H), 6.07(d, 1H), 6.16 (dd, 1H), 8.00(d, 1H) . Rf 0.15 (AcOEt)
37-35		¹ H-NMR (400MHz, CDCl ₃): 2.10-2.02 (m, 2H), 2.41(t, 2H), 3.56(dd, 2H), 3.71(t, 2H), 3.95(s, 3H), 4.19(t, 2H), 6.49(dd, 1H), 6.55(d, 1H), 7.99(d, 1H) . Rf 0.10 (AcOEt)
37-36		¹ H-NMR (400MHz, CDCl ₃): 2.14(s, 3H), 3.87-3.34(m, 8H), 3.99 (s, 3H), 7.01(dd, 1H), 7.16(d, 1H), 7.88(d, 1H) . Rf 0.25 (AcOEt)

37-37		$^1\text{H-NMR}$ (400MHz, CDCl_3): 3.49-3.37(m, 2H), 3.88-3.55(m, 6H), 3.99 (s, 3H), 7.00(dd, 1H), 7.16(d, 1H), 7.87(d, 1H) . Rf 0.50 (AcOEt)
37-38		$^1\text{H-NMR}$ (400MHz, CDCl_3): 1.17 (s, 3H), 1.19(s, 3H), 2.69 (t, 4H), 3.06(s, 2H), 3.42 (t, 4H), 3.96(s, 3H), 4.13(sept, 1H), 6.34 (d, 1H), 6.44(dd, 1H), 6.90-6.79(m, 1H), 8.00(d 1H) . Rf 0.20 (AcOEt)
37-39		$^1\text{H-NMR}$ (400MHz, CDCl_3): 1.44-1.34 (m, 2H), 1.84-1.77 (m, 1H), 1.94-1.85 (m, 2H), 3.04-2.94 (m, 2H), 3.55 (t, 2H), 3.96-3.57(m, 2H), 3.95(s, 3H), 6.31(d, 1H), 6.42(dd, 1H), 8.00(d, 1H) . Rf 0.30 (AcOEt)
37-40		$^1\text{H-NMR}$ (400MHz, CDCl_3): 1.44-1.34 (m, 2H), 1.84-1.77 (m, 1H), 1.94-1.85 (m, 2H), 3.04-2.94 (m, 2H), 3.55 (t, 2H), 3.96-3.57(m, 2H), 3.95(s, 3H), 6.31(d, 1H), 6.42(dd, 1H), 8.04(d, 1H) . Rf 0.45 (AcOEt)

37-41		¹ H-NMR (400MHz, CDCl ₃) : 4.05 (s, 3H), 7.07 (d, 1H), 7.08 (d, 1H), 7.27-7.26 (m, 1H), 7.33 (t, 1H), 7.92 (s, 1H), 8.04 (d, 1H). Rf: 0.20 (AcOEt)
37-42		¹ H-NMR (400MHz, CDCl ₃) : 2.34 (s, 3H), 2.55-2.37 (m, 4H), 3.86-3.38 (m, 4H), 4.00 (s, 3H), 7.13 (d, 1H), 7.66 (dd, 1H), 7.93 (d, 1H). Rf: 0.30 (AcOEt:MeOH=4:1)
37-43		¹ H-NMR (400MHz, CDCl ₃) : 2.43 (s, 3H), 2.74 (s, 6H), 7.91 (dd, 1H), 7.23 (d, 1H), 7.24 (d, 1H), 7.46 (dd, 1H). Rf: 0.70 (Hexane:AcOEt=5:1)
37-44		¹ H-NMR (400MHz, CDCl ₃) : 2.15 (s, 3H), 3.80-3.48 (m, 2H), 6.87 (dd, 1H), 6.92 (dd, 1H), 7.09 (d, 1H), 7.40 (dd, 2H), 8.54 (dd, 2H).
37-45		¹ H-NMR (400MHz, CDCl ₃) : 3.86 (s, 3H), 4.00 (s, 3H), 6.78 (d, 1H), 6.99 (dd, 2H), 7.14 (d, 1H), 7.48 (dd, 2H), 7.71 (dd, 1H), 8.03 (d, 1H). Rf: 0.30 (Hexane:AcOEt=3:1)


37-46		¹ H-NMR (400MHz, CDCl ₃) :1.44 (t, 3H), 3.10 (t, 4H), 3.86 (t, 4H), 4.13 (q, 2H), 7.01(dd, 1H), 7.08 (dd, 1H), 7.35 (d, 1H). Rf: 0.25 (Hexane:AcOEt=3:1)
37-47		¹ H-NMR (400MHz, CDCl ₃) :1.26 (t, 3H), 3.32 (t, 4H), 3.85 (t, 4H), 4.15 (q, 2H), 6.34(d, 1H), 6.42 (dd, 1H), 7.98 (d, 1H). Rf: 0.45 (Hexane:AcOEt=5:1)
37-48		¹ H-NMR (400MHz, CDCl ₃) :3.45 (s, 3H), 3.77 (dd, 2H), 3.81 (s, 3H), 4.06 (t, 2H), 7.08-7.08(m, 2H), 7.37 (t, 1H). Rf: 0.45 (Hexane:AcOEt=3:1)
37-49		¹ H-NMR (400MHz, CDCl ₃) :2.44 (t, 1H), 3.83 (s, 3H), 3.96 (ddd, 2H), 4.20 (t, 2H), 7.06 (d, 1H), 7.12(dd, 1H), 7.40 (d, 1H). Rf: 0.10 (Hexane:AcOEt=3:1)
37-50		¹ H-NMR (400MHz, CDCl ₃) :1.45 (t, 3H), 3.81 (s, 3H), 4.13 (q, 2H), 7.01(d, 1H), 7.08 (dd, 1H), 7.36 (d, 1H). Rf: 0.20 (Hexane:AcOEt=3:1)
37-51		¹ H-NMR (400MHz, CDCl ₃) :1.35 (s, 3H), 1.36(s, 3H), 3.81(s, 3H), 4.52 (sept, 1H), 7.08-7.01 (m, 2H), 7.31 (d, 1H). Rf: 0.30 (Hexane:AcOEt=3:1)

37-52		¹ H-NMR (400MHz, CDCl ₃) :1.05 (t, 3H), 1.83 (ddd, 2H), 3.81(s, 3H), 4.01(t, 2H), 7.01 (d, 1H), 7.08 (dd, 1H), 7.36 (d, 1H). Rf: 0.35 (Hexane:AcOEt=3:1)
37-53		¹ H-NMR (400MHz, CDCl ₃) :3.86 (s, 6H), 3.79 (s, 3H), 6.91 (dd, 1H), 7.00 (d, 1H), 7.18 (d, 1H). Rf: 0.5 (Hexane:AcOEt=9:1)
37-54		¹ H-NMR (400MHz, CDCl ₃) :4.04(s, 3H), 7.22 (d, 1H), 7.48(dd, 2H), 7.83 (dd, 1H), 8.16 (d, 1H), 8.69 (dd, 2H). Rf: 0.12 (Hexane:AcOEt=1:1)
37-55		¹ H-NMR (400MHz, CDCl ₃) :4.02 (s, 3H), 7.22 (d, 1H), 7.39 (ddd, 1H), 7.77(dd, 1H), 7.85(ddd, 1H), 8.08 (d, 1H), 8.63(dd, 1H), 8.83 (d, 1H). Rf: 0.55 (Hexane:AcOEt=2:1)
37-56		¹ H-NMR (400MHz, CDCl ₃) :4.03 (s, 3H), 7.19 (d, 1H), 7.28-7.24 (m, 1H), 7.72(dd, 1H), 7.80-7.76(m, 1H), 8.25 (dd, 1H), 8.52 (d, 1H), 8.69 (ddd, 1H). Rf: 0.55 (Hexane:AcOEt=2:1)

37-57		mp 90.7 °C; ¹ H-NMR (400MHz, CDCl ₃) δ (ppm): 1.68 (m; 2H), 2.00 (m; 2H), 2.36 (s; 1H), 2.62 (bs; 4H), 2.72 (m; 2H), 3.62 (m; 2H), 3.78 (bs; 4H), 3.90 (s; 3H), 6.99 (d; 1H); 7.13 (dd; 1H), 7.26 (s; 1H); 7.40 (s; 1H).
-------	---	---

38


Preparation of 1-[4-(4-Methoxy-3-nitro-phenyl)-piperazin-1-yl]-ethanone



To a solution of 5-bromo-1-methoxy-2-nitrobenzene (300 mg, 1.29 mmol) in dioxane, 1-acetyl piperazine (400mg, 3.12 mmol), cesium carbonate (1.0g, 3.07 mmol), palladium diacetate (29.0 mg, 0.129 mmol) and 2-(di-*t*-butylphosphino)biphenyl (77 mg, 0.258 mmol) are added and stirred at 100°C for 8 hours. After cooling, the mixture is filtered to remove insoluble material. The filtrate is poured into water and extracted with ethyl acetate twice. The organic layer is washed with water and then brine, dried over magnesium sulfate, and evaporated in vacuo. The residue is purified by silica gel column chromatography (n-hexane : ethyl acetate gradient) to afford 1-[4-(4-Methoxy-3-nitro-phenyl)-piperazin-1-yl]-ethanone (319mg, 44%) as yellow solids. ¹H-NMR (400MHz, CDCl₃, δ, ppm) : 2.14 (s, 3H), 3.63 (ddd, 4H), 3.63 (t, 2H), 3.78 (t, 2H), 3.92 (s, 3H), 7.03 (d, 1H), 7.12 (d, 1H), 7.41 (d, 1H). R_f (ethyl acetate): 0.18

39


Preparation of 1-(3-Methoxy-4-nitro-phenyl)-piperidin-4-one



To a solution of 4-piperidone hydrochloride monohydrate (10.0 g, 0.065mol) in DMF (80 mL) are added 4-Fluoro-2-methoxy-1-nitro-benzene (10.0 g, 0.058 mol) and potassium carbonate

(20.2 g), and the mixture is stirred at 70°C for 20 h. After a filtration, the filtrate is poured into H₂O (ca. 300 mL), and the resulting precipitates are collected by a filtration followed by washing with H₂O for several times to give title compound (8.98 g) in 61% yield. Orange solid. ¹H-NMR (400 MHz, CDCl₃, δ): 2.65-2.62 (4H, *m*), 3.81-3.78 (4H, *m*), 3.98 (3H, *s*), 6.34 (1H, *d*), 6.45 (1H, *dd*), 8.05 (1H, *d*).

40


Preparation of 1-[1-(3-Methoxy-4-nitro-phenyl)-piperidin-4-yl]-4-methyl-piperazine

To a solution of 1-(3-Methoxy-4-nitro-phenyl)-piperidin-4-one (4.96g, 0.020mol) in dichloroethane (50 ml) is added N-methylpiperazine (2.7ml, 0.024 mol) at 0 °C and the mixture is stirred at room temperature. After 4 h, sodium triacetoxy-borohydride (5.04g, 0.024mol) is added and the mixture is further stirred at room temperature for 24 h. After addition of 1N sodium hydroxide at 0 °C, the mixture is poured into water and extracted three times with dichloromethane. The organic layer is combined and extracted three times with 1N hydrochloride. The water layer is basified with 2N sodium hydroxide and extracted three times with dichloromethane. The organic layer is washed with brine, dried over sodium sulfate, and evaporated in vacuo to give the title compound as yellow solids (6.04g) in 91% yield.

¹H-NMR (400 MHz, CDCl₃, δ): 1.70-1.57 (2H, *m*), 2.03-1.93 (2H, *m*), 2.29 (3H, *s*), 2.55-2.38 (5H, *m*), 2.70-2.56 (4H, *m*), 2.97 (2H, *ddd*), 3.97-3.92 (2H, *m*), 3.95 (3H, *s*), 6.31 (1H, *d*), 6.42 (1H, *dd*), 8.00 (1H, *d*).

41

Preparation of 4'-Methoxy-4-methyl-3-nitro-biphenyl




To a solution of 4-methoxyphenyl-boronic acid (500 mg, 3.29 mmol) in toluene (5.2 mL) and ethanol (1.3 mL), potassium carbonate (910 mg, 6.58 mmol), tetrakis(triphenylphosphine)-palladium (228.1 mg, 0.099 mmol) and 4-bromo-1-methyl-2-nitrobenzene (711 mg, 3.29 mmol) are added and stirred at 100°C for 7 hours. The mixture is poured into water and extracted with ethyl acetate two times. The organic layer is washed with water and then brine, dried over magnesium sulfate, and evaporated in vacuo. The residue is purified with silica gel column chromatography (n-hexane : ethyl acetate = 5 : 1) to afford the 4'-methoxy-4-methyl-3-nitrobiphenyl (630mg, 79%) as a yellow solid.

¹H-NMR (400MHz, CDCl₃, δ, ppm) : 2.62 (s, 3H), 3.86 (s, 3H), 7.02-6.98 (m, 2H), 7.37 (d, 1H), 7.54 (dd, 2H), 7.68 (dd, 1H), 8.18 (d, 1H). R_f (hexane : ethyl acetate = 3:1): 0.40.

42

Preparation of 4-(2-Ethoxy-ethoxy)-1-(3-methoxy-4-nitro-phenyl)-piperidine






To a solution of 1-(3-Methoxy-4-nitro-phenyl)-piperidin-4-ol (300mg, 1.2 mmol) in N,N-dimethylformamide (3.0 mL), sodium hydride (1.52g, 3.8 mmol) is added. After stirring, 2-bromoethyl methyl ether (150μl, 1.6 mmol) is added and the mixture is further stirred at 70°C for 15 hours. After addition of saturated aqueous ammonium chloride, the mixture is poured into water and extracted twice with ethyl acetate. The organic layer is washed with brine, dried over sodium sulfate, and evaporated in vacuo. The residue is purified by silica gel column chromatography (n-hexane - ethyl acetate gradient) to afford 4-(2-Methoxy-ethoxy)-1-(3-methoxy-4-nitro-phenyl)-piperidine (111mg, 29%) as a yellow oil.

¹H-NMR (400MHz, CDCl₃, δ, ppm) : 1.52(t, 3H), 1.95-2.00(m, 2H), 1.70-1.79(m, 2H), 3.23(ddd, 2H), 3.58-3.64(m, 2H), 3.65-3.68(m, 2H), 3.64-3.72(m, 2H), 3.95(s, 3H), 6.31(d, 1H), 6.42(dd, 1H), 8.00(d, 1H). R_f 0.53 (n-hexane: AcOEt=1:1).

According the procedure described above using appropriate alkyl halides, the following compounds are prepared.

Ex- No.	Rx	Identification
------------	----	----------------

42-1		¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 2.04-2.21(m, 1H), 2.63(t, 2H), 2.68(t, 2H), 3.42(t, 4H), 3.87(t, 4H), 3.96(s, 3H), 6.33(d, 1H), 6.44(dd, 1H), 8.02(d, 1H). Rf 0.09 (AcOEt).
42-2		¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 1.71-1.79(m, 2H), 1.95-2.02(m, 2H), 3.22(ddd, 2H), 3.40(s, 3H), 3.55-3.57(m, 2H), 3.59-3.73(m, 3H), 3.65-3.67(m, 2H), 3.95(s, 3H), 6.31(d, 1H), 6.42(dd, 1H), 8.00(d, 1H). Rf 0.35 (<i>n</i> -hexane: AcOEt=1:1)

Example : 432-Methoxy-4-(1-methyl-piperidin-4-yloxy)-phenylamine 4-(3-Methoxy-4-nitro-phenoxy)-1-methyl-piperidine


To a solution of 4-Fluoro-2-methoxy-1-nitro-benzene (10.3g, 60 mmol) in toluene (50 mL) and 25% KOH aq.(50mL), 4-hydroxy-1-methylpiperidine (13.8g, 120 mmol) and tetra-*n*-butyl ammonium bromide (3.87g, 12mmol) are added at room temperature. The mixture is heated at 60°C for 1 day. The reaction mixture is cooled to room temperature, poured into ice water and extracted twice with ethyl acetate. The organic layer is successively washed with dil.HCl and brine, dried over sodium sulfate, and evaporated in vacuo to afford the crude compound in quantitative yield (13.4g).

R_f = 0.22 (methanol : dichloromethane = 1:4). ¹H-NMR (400 MHz, CDCl₃, δ, ppm): 1.84-1.92(m, 2H), 2.0-2.1(m, 2H), 2.3-2.4 (m, 2H), 2.33 (s, 3H), 2.65-2.75(m, 2H), 3.94(s, 3H), 4.39-4.46(m, 1H), 6.49 (dd, 1H), 6.99 (d, 1H), 6.54 (d, 1H), 7.99 (d, 1H).

Example : 44


2-Methoxy-4-(2-morpholin-4-yl-ethoxy)-phenylamine

3-Methoxy-4-nitro-phenol



To a solution of 3-Fluoro-4-nitro-phenol (15.7g, 100 mmol) in THF (300 mL) , 30% KOMe in Methanol(49mL, 210mmol) is added at 0°C. The mixture is heated to gentle reflux for 18 hours.

4-[2-(3-Methoxy-4-nitro-phenoxy)-ethyl]-morpholine



To a solution of 3-Methoxy-4-nitro-phenol (1.69g, 10 mmol) in DMF (25 mL) , 4-(2-Chloroethyl)morpholine hydrochloride(2.05g, 11mmol), K₂CO₃(1.52g, 11mmol), KI(332mg, 2mmol) are added at room temperature. The mixture is heated to gentle reflux for 4 hours. The reaction mixture is cooled to room temperature and quenched with water. The resulting mixture is extracted twice with ethyl acetate and then the organic layer is successively washed with water and brine, dried over sodium sulfate, filtered and evaporated in vacuo to afford the crude compound in 90% yield (2.55g).


R_f = 0.11 (AcOEt only). ¹H-NMR (400 MHz, CDCl₃), δ (ppm): 2.56-2.61(m, 4H), 2.83(t, The reaction mixture is cooled to room temperature and quenched slowly with 1NHCl aq at 0°C. The resulting mixture is extracted twice with ethyl acetate and then the organic layer is successively washed with brine, dried over sodium sulfate, filtered and evaporated in vacuo to afford the crude compound in 94% yield(15.9g).

R_f = 0.22 (methanol : dichloromethane = 1:4). ¹H-NMR (400 MHz, CDCl₃), δ (ppm): 3.95(s, 3H), 5.49(s, 1H), 6.44 (dd, 1H, J=8.8, 2.52Hz), 6.54 (d, 1H, J=2.52Hz), 7.96 (d, 1H J=8.6Hz).

3.72-3.76(m, 4H), 3.94(s, 3H), 4.18(t, 2H), 6.51 (dd, 1H, $J=9.08$, 2.52Hz), 6.56 (d, 1H, $J=2.48$ Hz), 8.00 (d, 1H $J=9.08$ Hz).
2H),

Example : 45


2-Methoxy-4-(2-morpholin-4-yl-ethoxy)-phenylamine
Acetic acid 4-methoxy-3-nitro-phenyl ester



To a solution of 4-Methoxyphenol (12.4g, 100 mmol) in AcOH (50 mL), Ac₂O (50mL) is added at room temperature. The mixture is heated to gentle reflux for 1.5 hour. The reaction mixture is cooled to room temperature and c.HNO₃ (d=1.38, 10mL) is added slowly at 0 °C. The mixture is heated to 55°C for 1.5h. The reaction mixture is cooled to room temperature and quenched with water at 0oC. The resulting solid is filtered on Buchner funnel to afford the crude compound in 76% yield (16.0g).


R_f = 0.59 (AcOEt : n-Hexane = 3:7). ¹H-NMR (400 MHz, CDCl₃), δ (ppm): 2.31(s, 3H), 3.96(s, 3H), 7.08 (d, 1H, $J=9.04$ Hz), 7.31 (dd, 1H, $J=9.04$, 3.04Hz), 7.96 (d, 1H $J=3.04$ Hz).

4-Methoxy-3-nitro-phenol




To a solution of Acetic acid 4-methoxy-3-nitro-phenyl ester (1.06g, 5 mmol) in EtOH (20 mL) , 1N NaOH aq (5.5mL) is added at 0°C. The mixture is stirred at room temperature for 2 hours. The reaction mixture is quenched with AcOH and extracted twice with ethyl acetate. The organic layer is successively washed with water and brine, dried over sodium sulfate, filtered and evaporated in vacuo to afford the crude compound in quantitative yield (840mg).

R_f = 0.59 (AcOEt : n-Hexane = 3:7). ¹H-NMR (400 MHz, CDCl₃), δ (ppm): 3.91(s, 3H), 6.99 (d, 1H, $J=9.04$ Hz), 7.17 (dd, 1H, $J=9.04$, 3.00Hz), 7.38 (d, 1H $J=3.04$ Hz).

4-[2-(4-Methoxy-3-nitro-phenoxy)-ethyl]-morpholine





To a solution of 4-Methoxy-3-nitro-phenol (1.01g, 6 mmol) in DMF (15 mL), 4-(2-Chloroethyl)morpholine hydrochloride (1.34g, 7.2mmol), K₂CO₃ (2.49g, 18mmol), KI(2.99g, 18mmol) are added at room temperature. The mixture is heated to 80°C for 4 hours. The reaction mixture is cooled to room temperature and quenched with saturated NH₄Cl solution in water. The resulting mixture is extracted twice with ethyl acetate and then the organic layer is successively washed with water and brine, dried over sodium sulfate, filtered and evaporated in vacuo to afford the crude compound in quantitative yield (1.70g). R_f = 0.14 (AcOEt only). ¹H-NMR (400MHz, DMSO, δ , ppm) : 2.36-2.51 (m, 4H), 2.67 (t, J=5.5, 2H), 3.52-3.60 (m, 4H), 3.86 (s, 3H), 4.11 (t, J=6.0, 2H), 7.25-7.29 (m, 2H), 7.46-7.49 (m, 1H).






Preparation of 2-Methoxy-4-(1-methyl-piperidin-4-yloxy)-phenylamine:





To a solution of 4-(3-Methoxy-4-nitro-phenoxy)-1-methyl-piperidine (3.0g, 11.3 mmol) in ethanol (50 mL), 5% palladium on carbon(300mg) is added under a nitrogen atmosphere. The reaction vessel is fitted with a balloon adapter and charged with hydrogen and evacuated three times until the reaction is under a hydrogen atmosphere. The reaction is allowed to stir overnight. The reaction mixture is filtered through a pad of Celite and washed with methanol. The filtrate is concentrated in vacuo to afford 2-Methoxy-4-(1-methyl-piperidin-4-yloxy)-phenylamine in quantitative yield (2.7g).

R_f = 0.41 (methanol : dichloromethane = 1:1). ¹H-NMR (400 MHz, CDCl₃), δ (ppm): 1.75-1.86(m, 2H), 1.92-2.05(m, 2H), 2.2-2.32 (m, 2H), 2.30 (s, 3H), 3.4-3.7(brs, 2H), 3.82(s, 3H), 4.1-4.2(m, 1H), 6.37(dd, 1H), 6.46 (d, 1H), 6.61 (d, 1H).






By repeating the procedures described above using appropriate starting materials and conditions the following compounds are obtained.






Ex-No.	Rx	Identification
46-1		¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 3.92(s,3H), 3.97(br,2H), 6.75(d,1H), 7.00(dd, 1H), 7.12(d, 1H), 8.06(s, 1H), 8.41(s, 1H). Rf 0.32 (AcOEt)
46-2	 [1-(4-Amino-3-methoxy-phenyl)-pyrrolidin-3-yl]-ethyl-amine	¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 1.13(t, 3H), 1.77-1.86(m, 1H), 2.19-2.27(m,1H), 2.67-2.75(m, 2H), 3.01-3.06(m, 1H), 3.20-3.26(m, 1H), 3.33-3.38(m, 1H), 3.42-3.49(m, 2H), 3.84(s, 3H), 6.04-6.07(m, 1H), 6.14-6.15(m, 1H), 6.64-6.66(m, 1H). Rf 0.2 (AcOEt only)
46-3	 2-Methoxy-4-(4-methyl-piperazin-1-yl)-phenylamine	¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 2.44(s, 3H), 2.70-2.73(m, 4H), 3.13-3.17(m, 4H), 3.48(brs, 2H), 3.84(s, 3H), 6.41(dd, 1H, J=8.5, 2.52Hz), 6.51(d, 1H, J=2.52Hz), 6.64(d, 1H, J=8.5Hz). Rf 0.2 (AcOEt only).
46-4	 2-Methoxy-4-(1-methyl-piperidin-3-ylmethoxy)-phenylamine	¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 1.01-1.12(m, 1H), 1.57-2.13(m, 6H), 2.26(s,3H), 2.74-2.77(m, 1H), 2.93-2.96(m, 1H), 3.47 (bs, 2H), 3.70-3.80(m, 2H), 3.82(s, 3H), 6.31-6.34(m, 1H), 6.44-6.45(m, 1H), 6.60-6.62(m, 1H). Rf 0.2 (AcOEt only)






46-5		$^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.80-1.67 (2H, <i>m</i>), 1.99-1.90 (2H, <i>m</i>), 2.42-2.27 (1H, <i>m</i>), 2.56-2.43 (4H, <i>m</i>), 2.68-2.58 (2H, <i>m</i>), 2.76-2.58 (4H, <i>m</i>), 3.57-3.48 (2H, <i>m</i>), 3.83 (3H, <i>s</i>), 6.41 (1H, <i>dd</i>), 6.52 (1H, <i>d</i>), 6.63 (1H, <i>d</i>). R_f (hexane/acetone 1:1) 0.44.
46-6		$^1\text{H-NMR}$ (400MHz, CDCl_3 , δ , ppm) : 1.83-1.95 (<i>m</i> , 2H), 1.97-2.08 (<i>m</i> , 2H), 2.20-2.31 (<i>m</i> , 1H), 2.60-2.72 (<i>m</i> , 2H), 3.46-3.53 (<i>m</i> , 2H), 3.84 (<i>s</i> , 3H), 5.42-5.60 (<i>m</i> , 1H), 6.43 (<i>dd</i> , 1H), 6.53 (<i>d</i> , 1H), 6.64 (<i>d</i> , 1H).
46-7		$^1\text{H-NMR}$ (400MHz, CDCl_3 , δ , ppm) : 2.13 (<i>s</i> , 3H), 3.01-3.05 (<i>m</i> , 4H), 3.59 (<i>t</i> , 2H), 3.75 (<i>t</i> , 2H), 3.81 (<i>s</i> , 3H), 6.30 (<i>dd</i> , 1H), 6.39 (<i>bs</i> , 1H), 6.71 (<i>d</i> , 1H).
46-8		$^1\text{H-NMR}$ (400MHz, CDCl_3 , δ , ppm) : 1.84-1.97 (<i>m</i> , 2H), 1.98-2.07 (<i>m</i> , 2H), 2.20-2.32 (<i>m</i> , 1H), 2.61-2.72 (<i>m</i> , 2H), 3.47-3.55 (<i>m</i> , 2H), 3.95 (<i>s</i> , 3H), 5.20-5.38 (<i>m</i> , 1H), 5.40-5.56 (<i>m</i> , 2H), 6.43 (<i>d</i> , 1H), 6.53 (<i>bs</i> , 1H), 6.64 (<i>d</i> , 1H).
46-9		$^1\text{H-NMR}$ (400MHz, CDCl_3 , δ , ppm) : 2.59-2.67 (<i>m</i> , 2H), 2.77-2.68 (<i>m</i> , 4H), 3.08-3.15 (<i>m</i> , 4H), 3.49-3.56 (<i>m</i> , 1H), 3.67-3.77 (<i>m</i> , 2H), 3.98 (<i>s</i> , 3H), 6.41-6.43 (<i>m</i> , 1H), 6.52 (<i>bs</i> , 1H), 6.65 (<i>d</i> , 1H).





46-10		¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 1.72-1.96 (m, 2H), 1.98-2.10 (m, 2H), 2.63 (s, 3H), 2.73-2.84 (m, 2H), 3.40 (s, 3H), 3.34-3.42 (m, 2H), 3.44-3.49 (m, 1H), 3.55-3.57 (m, 2H), 3.64-3.66 (m, 2H), 3.83 (s, 3H), 6.41-6.43 (m, 1H), 6.53 (bs, 1H), 6.63 (d, 1H).
46-11		¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 1.22(t, 3H), 1.72-1.84 (m, 2H), 2.00-2.10 (m, 2H), 2.72-2.82 (m, 2H), 3.33-3.38 (m, 2H), 3.43-3.49 (m, 1H), 3.55 (q, 2H), 3.58-3.61 (m, 2H), 3.64-3.66 (m, 2H), 3.83 (s, 3H), 6.41-6.43 (m, 1H), 6.53 (bs, 1H), 6.63 (d, 1H).
46-12		¹ H-NMR (400MHz, CDCl ₃ , δ, ppm) : 2.20 (s, 3H), 3.84 (s, 3H), 6.87 (d, 1H), 6.89 (dd, 1H), 6.95 (d, 2H), 7.09 (d, 1H), 7.48 (d, 2H). R _f (n-hexane : ethyl acetate = 1:1): 0.50.
46-13		¹ H- NMR (400MHz, CDCl ₃ , δ, ppm): 1.49 – 1.59 (m, 3H), 1.70 – 1.95 (m, 6H), 2.00 – 2.20 (m, 2H), 2.60 – 2.90 (m, 7H), 3.50 – 3.60 (m, 3H), 3.83 (s, 3H), 3.85 - 3.91 (m, 1H), 6.41 (dd, 1H, J = 8.0, 2.5 Hz), 6.50 (d, 1H, J = 2.5 Hz), 6.63 (d, 1H, J = 8.0 Hz)





46-14		¹ H-NMR (400MHz, DMSO-d ₆ , δ , ppm) : 1.87-1.79(m, 1H), 2.22 (ddd, 1H), 2.48 (s, 3H), 3.05 (dd, 1H), 3.28-3.21 (m, 1H), 3.40-3.32 (m, 2H), 3.45 (dd, 1H), 3.84 (s, 3H), 6.06 (dd, 1H), 6.15 (d, 1H), 6.66 (d, 1H)
46-15		¹ H-NMR (400MHz, CDCl ₃ , δ , ppm): 2.35 - 2.73 (m, 4H), 2.68 - 2.75 (m, 1H), 2.82 - 2.93 (m, 2H), 3.14 - 3.19 (m, 1H), 3.29 - 3.40 (m, 2H), 3.50 - 3.60 (bs, 2H), 3.69 - 3.78 (m, 2H), 3.84 (s, 3H), 3.85 - 3.91 (m, 1H), 6.40 (dd, 1H, J = 8.0, 2.5 Hz), 6.50 (d, 1H, J = 2.5 Hz), 6.64 (d, 1H, J = 8.0 Hz)
46-16		¹ H-NMR (400MHz, DMSO-d ₆ , δ , ppm) : 1.95-1.85(m, 1H), 2.22-2.14 (m, 1H), 2.31 (s, 3H), 2.89-2.79 (m, 1H), 3.10 (t, 1H), 3.39-3.25 (m, 3H), 3.42 (t, 1H), 3.85 (s, 3H), 6.05 (dd, 1H), 6.14 (d, 1H), 6.67 (d, 1H)
46-17		¹ H-NMR (400MHz, CDCl ₃ , δ , ppm) : 1.68-1.81 (m, 2H), 1.97-2.09 (m, 2H), 2.74-2.87 (m, 2H), 3.31-3.41 (m, 2H), 3.77-3.88 (m, 1H), 3.84 (s, 3H), 6.40-6.48 (m, 1H), 6.65 (bs, 1H), 6.64 (d, 1H).
46-18		¹ H-NMR (400 MHz, CDCl ₃), δ (ppm): 2.55-2.61(m, 4H), 2.80(t, 2H), 3.72-3.77(m, 4H), 3.81(s, 3H), 4.05(t, 2H), 6.24 (dd, 1H, J=8.56, 2.52Hz), 6.34 (d, 1H, J=2.52Hz), 6.68 (d, 1H, J=8.56Hz). R _f = 0.31 (methanol : dichloromethane = 1:9).





46-19		¹ H-NMR (400 MHz, CDCl ₃), δ (ppm): 2.55-2.61(m, 4H), 2.78(t, 2H), 3.72-3.77(m, 4H), 3.82(s, 3H), 4.05(t, 2H), 6.35 (dd, 1H, $J=8.56$, 2.52Hz), 6.47 (d, 1H, $J=2.52$ Hz), 6.63 (d, 1H $J=8.56$ Hz). R _f = 0.61 (methanol : dichloromethane = 1:4).
46-20		¹ H-NMR (DMSO), δ (ppm): 3.84 (s, 3H), 6.95-7.00 (m, 1H), 7.08-7.12 (m, 2H).
46-21		¹ H-NMR (400MHz, CDCl ₃): 1.47-1.34(m, 2H), 1.75-1.63 (m, 1H), 1.86-1.79(m, 2H), 2.64-2.58 (m, 2H), 3.28(d, 2H), 3.61(d, 3H), 3.87(s, 3H), 3.36(s, 1H), 3.49-3.45 (m, 2H), 3.84(s, 3H), 6.43(dd, 1H), 6.53(d, 1H) 6.64(d, 1H)
46-22		¹ H-NMR (400MHz, CDCl ₃): 1.13(t, 3H), 2.49(dd, 2H), 2.68-2.59 (m, 4H), 3.10 (t, 4H), 3.84(s, 3H), 6.43(dd, 1H), 6.53(d, 1H) 6.65(d, 1H)
46-23		¹ H-NMR (400MHz, CDCl ₃): 1.78-1.68 (m, 2H), 1.99-1.89 (m, 2H), 2.36-2.20(m, 1H), 2.67-2.50(m, 6H), 3.56-3.48(m, 2H), 3.79-3.69(m, 4H), 3.84(s, 3H), 6.42(dd, 1H), 6.52(d, 1H) 6.64(d, 1H)







46-24		¹ H-NMR (400MHz, CDCl ₃): 1.08 (s, 3H), 1.10 (s, 3H), 2.69(t, 4H), 2.72-2.68 (m, 1H), 3.08 (t, 4H), 3.83(s, 3H), 6.42(dd, 1H), 6.53(d, 1H) 6.64(d, 1H)
46-25		¹ H-NMR (400MHz, CDCl ₃): 1.96-1.84 (m, 2H), 2.07-1.99 (m, 2H), 2.32-2.28(m, 1H), 2.70-2.60(m, 2H), 3.54-3.47(m, 2H), 3.84(s, 3H), 5.35-5.24(m, 1H), 5.50-5.45 (m, 1H), 6.42(dd, 1H), 6.52(d, 1H) 6.64(d, 1H)
46-26		¹ H-NMR (400MHz, CDCl ₃): 2.18-2.03 (m, 2H), 3.28-3.19 (m, 2H), 3.39-3.31(m, 1H), 3.36(s, 3H), 3.49-3.42 (m, 1H), 3.85 (s, 3H), 6.07(dd, 1H), 6.16(d, 1H), 6.66(d, 1H)
46-27		¹ H-NMR (400MHz, CDCl ₃): 1.48(s, 9H), 1.88-1.71 (m, 2H), 1.97-1.82 (m, 2H), 2.78 (s, 3H), 2.84-2.64(m, 2H), 3.55-3.48(m, 2H), 3.95(s, 3H), 3.84 (s, 3H), 6.43(d, 1H), 6.52(bs, 1H), 6.64(d, 1H)
46-28		¹ H-NMR (400MHz, CDCl ₃): 3.02(s, 3H), 3.33(dd, 2H), 3.44(t, 2H), 3.74 (s, 2H), 3.83(s, 3H), 6.38(dd, 1H), 6.47(d 1H), 6.66(d, 1H)







46-29		¹ H-NMR (400MHz, CDCl ₃): 1.78-1.38 (m, 2H), 1.96-1.89 (m, 2H), 2.30(s, 3H), 2.39-2.31(m, 1H), 2.55-2.42(m, 4H), 2.71-2.56(m, 6H), 3.35-3.49 (m, 2H), 3.83 (s, 3H), 6.41(dd, 1H), 6.52(d, 1H) 6.63(d, 1H)
46-30		¹ H-NMR (400MHz, CDCl ₃): 3.80(s, 3H), 3.82(s, 3H), 3.82 (s, 3H), 6.40(s, 1H), 6.54(s, 1H)
46-31		¹ H-NMR (400MHz, CDCl ₃): 3.20(t, 2H), 4.57(t, 2H), 6.55(dd, 1H), 6.70-6.65(m, 1H), 6.68 (d, 1H). Rf 040 (AcOEt)
46-32		¹ H-NMR (400MHz, CDCl ₃): 2.98 (t, 4H), 3.62 (bs, 2H), 3.79 (s, 3H), 3.81 (s, 3H), 3.87(t, 4H), 6.36(s, 1H), 6.53(s, 1H)
46-33		¹ H-NMR (400MHz, CDCl ₃): 2.37 (s, 3H), 2.61 (t, 4H), 3.27 (t, 4H), 3.88 (s, 3H), 3.95(s, 3H), 6.48(s, 1H), 7.56(s, 1H)







46-34	 <chem>CCOC1=CC=C(N1CCN2CCNCC2)C=C</chem>	¹ H-NMR (400MHz, CDCl ₃): 1.05(t, 3H), 1.83 (ddd, 2H), 2.35(s, 3H), 2.58(t, 4H), 3.07(t, 4H), 3.94(t, 2H), 6.41(dd, 1H), 6.51(d, 1H), 6.65(d, 1H)
46-35	 <chem>CC(=O)N1C(C)CCN(C1)C2=CC=C(N)C=C2OC</chem>	¹ H-NMR (400MHz, CDCl ₃): 1.28(s, 3H), 1.30 (s, 3H), 2.04 (s, 2H), 2.17(s, 3H), 2.84-2.72 (m, 2H), 3.20 (d, 2H), 3.86 (s, 3H), 6.41(d, 1H), 6.46(dd, 1H), 6.66(d, 1H),
46-36	 <chem>CC(=O)N1CCN(C1)C2=CC=C(N)C=C2OC</chem>	¹ H-NMR (400MHz, CDCl ₃): 1.18(t, 3H), 2.39(dd, 2H), 3.07-2.98(m, 4H), 3.61(t, 2H), 3.78(t, 2H), 3.88(s, 3H), 6.41(dd, 1H), 6.51(d, 1H), 6.65(d, 1H)
46-37	 <chem>CC(C)C(=O)N1CCN(C1)C2=CC=C(N)C=C2OC</chem>	¹ H-NMR (400MHz, CDCl ₃): 1.15(s, 3H), 1.16(s, 3H), 2.83(sept, 1H), 3.07-2.98(m, 4H), 3.73-3.64(m, 2H), 3.83-3.76(m, 2H), 3.84(s, 3H), 6.41(dd, 1H), 6.51(d, 1H), 6.65(d, 1H)

46-38	 <chem>CN(C)C(=O)N1CCN(CC1)c2ccc(OC)c(N)c2</chem>	¹ H-NMR (400MHz, CDCl ₃): 2.84(d, 3H), 3.02(t, 4H), 3.51(t, 4H), 3.84(s, 3H), 4.48-4.38(m, 1H), 6.41(dd, 1H), 6.51(d, 1H), 6.65(d, 1H)
46-39	 <chem>CN(C)C(=O)N1CCN(CC1)c2ccc(OC)c(N)c2</chem>	¹ H-NMR (400MHz, CDCl ₃): 1.99-1.81 (m, 2H), 2.23-2.12(m, 2H), 2.69-2.58(m, 2H), 2.84 (d, 3H), 3.54-3.45(m, 2H), 3.84(s, 3H), 5.55-5.45(m, 1H), 6.42(dd, 1H), 6.52(d, 1H), 6.64(d, 1H)
46-40	 <chem>CN(C)C(=O)N1CCN(CC1)c2ccc(OC)c(N)c2</chem>	¹ H-NMR (400MHz, CDCl ₃): 1.53 (s, 6H), 2.11(s, 3H), 3.05(s, 2H), 3.28(t, 2H), 3.64(t, 2H), 3.86 (s, 3H), 6.26 (dd, 1H), 6.33(d, 1H), 6.67(d, 1H)
46-41	 <chem>CN(C)C(=O)N1CCN(CC1)c2ccc(OC)c(N)c2</chem>	¹ H-NMR (400MHz, CDCl ₃): 2.55-2.41 (m, 4H), 2.63 (t, 2H), 3.13(t, 2H), 3.77-3.68(m, 4H), 3.83(s, 3H), 6.15(dd, 1H), 6.25 (d, 1H), 6.62(d, 1H)


46-42	 <chem>COC1=CC=C(N)C=C1OCCN2CCCC2=O</chem>	¹ H-NMR (400MHz, CDCl ₃): 2.05-2.00 (m, 2H), 2.39(t, 2H), 3.57(t, 2H), 3.64 (t, 2H), 3.83(s, 3H), 4.04(t, 2H), 6.32 (dd, 1H), 6.44(d, 1H), 6.63(d, 1H)
46-43	 <chem>CC(=O)N1CCN(C1)C(=O)C2=CC=C(N)C(OC)=C2</chem>	¹ H-NMR (400MHz, CDCl ₃): 2.13 (s, 3H), 3.53-3.46 (m, 2H), 3.65-3.55(m, 4H), 3.71-3.66(m, 2H), 3.88 (s, 3H), 6.67(d, 1H), 6.87(dd, 1H), 6.95(d, 1H)
46-44	 <chem>CC(=O)N1CCN(C1)C(=O)C2=CC=C(N)C(OC)=C2</chem>	¹ H-NMR (400MHz, CDCl ₃): 3.73-3.61(m,8H), 3.87(s, 3H), 6.65(d, 1H), 6.86(dd, 1H), 6.95(d, 1H)
46-46	 <chem>CC(C)NC(=O)CN1CCN(C1)C2=CC=C(N)C(OC)=C2</chem>	¹ H-NMR (400MHz, CDCl ₃): 1.17 (s, 3H), 1.19(s, 3H), 2.69(t, 4H), 3.04 (s, 2H), 3.08(t, 4H),4.15-4.07(m, 1H), 6.41 (dd, 1H), 6.51(d, 1H), 6.65(d, 1H), 7.01-6.94(m 1H)

46-47		¹ H-NMR (400MHz, CDCl ₃): 3.35-3.28 (m, 2H), 3.53-3.46(m, 2H), , 3.76 (s, 2H), 3.84(s, 3H), 5.92-5.83 (m, 1H), 6.40(dd, 1H), 6.48(d, 1H), 6.67 (d,1H)
46-48		¹ H-NMR (400MHz, CDCl ₃): 2.09-2.00 (m, 2H), 2.25-2.15 (m, 2H), 3.29-3.20 (m, 2H), 3.51-3.40(m, 4H), 3.85(s, 3H), 4.62-4.55(m, 1H), 6.08(d, 1H), 6.18(d, 1H), 6.67(d, 1H)
46-49		¹ H-NMR (400MHz, CDCl ₃): 1.52-1.40 (m, 2H), 1.90-1.84 (m, 2H), 2.68-2.59 (m, 2H), 3.51-3.45(m, 2H), 3.84(s, 3H), 6.44(dd, 1H), 6.54(d, 1H), 6.64(d, 1H)
46-50		¹ H-NMR (400MHz, CDCl ₃) : 2.14 (s, 3H), 2.66 (s, 6H), 6.44 (d, 1H), 6.54 (d, 1H), 6.98 (t, 1H).
46-51		¹ H-NMR (400MHz, CDCl ₃) :2.63 (s, 3H), 7.49-7.45 (m, 1H), 7.74-7.62 (m, 2H), 7.76 (dd, 1H), 8.24(d, 1H), 8.77-8.64 (m, 2H).
46-52		¹ H-NMR (400MHz, CDCl ₃) :3.84 (s, 3H), 3.88 (s, 3H), 6.78(d, 1H), 6.83 (d, 1H), 7.00-6.89 (m, 3H), 7.45 (d, 1H).

46-53		$^1\text{H-NMR}$ (400MHz, CDCl_3) : 1.40 (t, 3H), 3.03 (t, 4H), 3.84 (t, 4H), 4.00 (q, 2H), 6.27 (dd, 1H), 6.38 (d, 1H), 6.71 (dd, 1H),
46-54		$^1\text{H-NMR}$ (400MHz, CDCl_3) : 1.26 (t, 3H), 3.02 (t, 4H), 3.85 (t, 4H), 4.05 (q, 2H), 6.40 (dd, 1H), 6.49 (d, 1H), 6.66 (d, 1H),
46-55		$^1\text{H-NMR}$ (400MHz, CDCl_3) : 3.44 (s, 3H), 3.73 (s, 3H), 3.74-3.68 (m, 2H), 3.95-3.85 (m, 2H), 4.10-4.05 (m, 2H), 6.21 (dd, 1H), 6.32 (d, 1H), 6.75 (d, 1H).
46-56		$^1\text{H-NMR}$ (400MHz, CDCl_3) : 2.35-2.26 (m, 1H), 3.74 (s, 3H), 3.93-3.86 (m, 2H), 4.09-4.07 (m, 2H), 6.25 (dd, 1H), 6.34 (d, 1H), 6.76 (d, 1H).
46-57		$^1\text{H-NMR}$ (400MHz, CDCl_3) : 1.40 (t, 3H), 3.71 (s, 3H), 4.00 (q, 2H), 6.22 (dd, 1H), 6.33 (d, 1H), 6.69 (d, 1H).
46-58		$^1\text{H-NMR}$ (400MHz, CDCl_3) : 1.32 (d, 6H), 3.73 (s, 3H), 3.85-3.71 (m, 2H), 4.37 (sept, 1H), 6.22 (dd, 1H), 6.32 (d, 1H), 6.72 (d, 1H).

46-59		¹ H-NMR (400MHz, CDCl ₃) : 1.04 (t, 3H), 1.80 (ddd, 2H), 3.72 (s, 3H), 3.85-3.75 (m, 2H), 3.90 (t, 2H), 6.22 (dd, 1H), 6.33 (d, 1H), 6.69 (d, 1H).
46-60		¹ H-NMR (400MHz, CDCl ₃) : 2.94 (s, 6H), 3.89 (s, 3H), 6.16 (dd, 1H), 6.25 (d, 1H), 6.72 (d, 1H).
46-61		¹ H-NMR (400MHz, CDCl ₃) : 3.91(s, 3H), 6.87 (d, 1H), 7.02 (dd, 1H), 7.05 (d, 1H), 7.44 (dd, 2H), 8.59 (dd, 2H).
46-62		¹ H-NMR (400MHz, CDCl ₃) : 3.91 (s, 3H), 6.88 (d, 1H), 6.96-6.93(m, 1H), 7.31(ddd, 1H), 7.83-7.80 (m, 1H), 8.51(dd, 1H), 8.78(dd, 1H).
46-63		¹ H-NMR (400MHz, CDCl ₃) : 3.91 (s, 3H), 6.87 (dd, 1H), 7.16(ddd, 1H), 7.34(dd, 1H), 7.43 (d, 1H), 7.72-7.64 (m, 2H), 8.63-8.61 (m, 1H).
46-64		mp 148.6°C; ¹ H-NMR (500MHz, CDCl ₃) δ (ppm): 1.63 (m; 2H), 1.99 (m; 2H), 2.27 (m; 1H), 2.60 (m; 6H), 3.52 (m; 2H), 3.71 (m; 4H), 3.78 (s; 3H), 6.36 (dd; 1H); 6.52 (d; 1H), 6.73 (d; 1H).


46-65		
-------	--	--

47Preparation of 4-(3-amino-4-methylbenzoyl)-piperazine-1-carboxylic acid tert-butyl ester

To a solution of 4-methyl-3-nitro-benzoic acid (300 mg, 2.76 mmol), N-butoxycarbonyl-piperazine (340 mg, 1.83 mmol) in DMF (3.0 mL), triethylamine (300 μ L, 3.59 mmol), TBTU (800 mg, 2.49 mmol) and HOAt (270.5 mg, 1.99 mmol) are added and stirred at room temperature for 24 hours. The mixture is poured into water and extracted twice with ethyl acetate. The organic layer is washed with water and then brine, dried over magnesium sulfate, and evaporated in vacuo. The residue is purified with silica gel column chromatography (n-hexane : ethyl acetate = 5 : 1) to afford 4-(4-methyl-3-nitrobenzoyl)-piperazine-1-carboxylic acid tert-butyl ester as a colorless solid.

$^1\text{H-NMR}$ (δ , ppm) : 1.47 (s, 9H), 2.64 (s, 3H), 3.88-3.28 (m, 8H), 7.42 (d, 1H), 7.56 (dd, 1H), 8.03 (d, 1H). R_f (hexane : ethyl acetate = 10:1): 0.13.

The title compound is obtained by reduction with hydrogen over 10% palladium on charcoal in methanol solution.

48Preparation of 4-(3-amino-4-methylphenyl)-morpholine

To a suspension of 4-bromo-1-methyl-2-nitrobenzene (225 mg, 1.04 mmol), morpholine (125 μ L, 1.25 mmol), and cesium carbonate (474.4 mg, 1.46 mmol) in toluene, palladium diacetate (31.2 mg, 0.139 mmol) and 2-(di-*t*-butylphosphino)biphenyl (125 mg, 0.403 mmol) are added and stirred at 100°C for 5 hours. After cooling, the mixture is filtered to remove insoluble material. The filtrate is poured into water and extracted with ethyl acetate twice. The organic layer is washed with water and then brine, dried over magnesium sulfate, and evaporated in


vacuo. The residue is purified with silica gel column chromatography (n-hexane : ethyl acetate = 5 : 1) to afford 4-(4-methyl-3-nitrophenyl)-morpholine as a yellow solid.

$^1\text{H-NMR}$ (δ , ppm) : 2.50 (s, 3H), 3.19-3.17 (m, 4H), 3.88-3.86 (m, 4H), 7.04 (dd, 1H), 7.21 (d, 1H), 7.47 (d, 1H). R_f (hexane : ethyl acetate = 5:1): 0.20.

The title compound is obtained by reduction with hydrogen over 10% palladium on charcoal in methanol solution.

49

Preparation of 2-[5-chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-benzoic acid



To a solution of 1.0 g (3.37 mmol) of 2-(2,5-dichloro-pyrimidin-4-ylamino)-*N*-methyl-benzamide in 15 mL of acetic acid are added 2-methoxy-4-morpholinoaniline dihydrochloride (1.9 g, 6.73 mmol) and 6.0 mL of 1N ethanolic solution of hydrogen chloride (6.0 mmol). After the reaction mixture is stirred at 120°C for 16 hours and cooled to room temperature, aqueous NaHCO_3 solution is added to adjust the acidity between pH 5 and pH 6. The resulting precipitate is collected by a filtration and dried under reduced pressure to give 2-[5-chloro-2-(2-methoxy-4-morpholin-4-yl-phenyl-amino)-pyrimidin-4-ylamino]-benzoic acid (970 mg, 2.12 mmol, 63%) as ivory solid.


NMR (400MHz, DMSO-d_6 , δ): 3.10-3.20 (m, 4H), 3.78 (s, 3H), 3.70-3.80 (m, 4H), 6.52 (dd, 1H, $J = 8.56, 2.52$ Hz), 6.67 (d, 1H, $J = 2.52$ Hz), 7.08 (dd, 1H, $J = 8.04, 8.04$ Hz), 7.39 (d, 1H, $J = 8.56$ Hz), 7.35-7.45 (m, 1H), 7.99 (dd, 1H, $J = 8.04, 1.52$ Hz), 8.14 (s, 1H), 8.28 (s, 1H) 8.70-8.80 (m, 1H).

Example 50: Sulfonamide moieties are prepared as follows:

Preparation of 2-amino-4-chloro-5-methyl-benzenesulfonyl chloride

To a solution of 2-amino-5-chloro-4-methyl-benzenesulfonic acid (3.0 g, 1.35 mmol) in dichloroethane (10 mL) is added sulfonyl chloride (4.4 mL, 3.83 mmol) and stirred at 60°C. After one hour, thionyl chloride (1.3 mL) is added and the mixture is further stirred at 100°C for 7.0 hours. The mixture is poured into iced water and extracted with ether three times. The organic layer is washed with water and then brine, dried over sodium sulfate, and evaporated in vacuo. ¹H-NMR (δ, ppm) : 2.35 (s, 3H), 6.68 (s, 1H), 7.75 (s, 1H).


This substituted sulfonyl chloride is reacted with a suitable amine. On reaction e.g. with methylamine, 2-amino-5-chloro-4,N-dimethylbenzenesulfonamide is formed.

Example 51Preparation of 2-[5-bromo-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-N,N-dimethyl-benzensulfonamide

To a solution of 2-[5-Bromo-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzenesulfonamide (Ex3-19) (1.0g, 1.82mmol) in DMF (10mL), potassium carbonate (300mg, 2.17mmol) and iodomethane (116μl, 1.86mmol) are added. The resulting suspension is stirred at 50°C for 1h. To the reaction mixture, water is added and extracted with ethyl acetate three times. The organic layer is washed with water, dried over sodium sulfate, and concentrated in vacuo. The residue is purified by aluminum oxide column chromatography (AcOEt) to afford the title compound (728mg, 71% yield).

NMR (400MHz, CDCl₃, δ): 2.74 ((s, 6H), 3.05-3.18 (m, 4H), 3.84-3.93 (m, 4H), 3.88 (s, 3H), 6.43 (dd, 1H), 6.53 (d, 1H), 7.24 (m, 1H), 7.31 (s, 1H), 7.56 (m, 1H), 7.87 (dd, 1H), 8.05 (d, 1H), 8.21 (s, 1H), 8.49 (d, 1H), 8.49 (d, 1H), 9.27 (s, 1H). Rf: 0.23 (AcOEt:Hexane=1:1).

Example 52Preparation of 2-[5-Bromo-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-5-fluoro-N-methyl-benzenesulfonamide



Preparation of 7-Fluoro-1,1-dioxo-1,4-dihydro-2H-1 λ^6 -benzo[1,2,4]thiadiazin-3-one

To a solution of chlorosulfonylisocyanate (1.2mL, 13.5mmol) in nitroethane (10mL), 4-fluoroaniline (1.0g, 8.97mmol) is added dropwise at 0°C and the reaction mixture is stirred for 30min. To the solution, aluminum chloride (1.3g, 9.87mmol) is added at 0°C and the mixture is stirred at 100°C for 1 hour. After cooling to room temperature, water is added and the mixture is extracted with ethyl acetate twice. The organic layer is washed with brine, dried over sodium sulfate, and concentrated under reduced pressure. The resulting solids are collected by a filtration and washed with ether to give slightly gray solids (803.9mg, 41%).

NMR (400MHz, DMSO-d₆, δ): 7.22-7.28 (m, 1H), 7.45-7.57 (m, 1H), 7.60 (m, 1H), 11.15-11.30 (m, 1H). Rf: 0.43 (MeOH:AcOEt=1:5).

Preparation of 7-Fluoro-2-methyl-1,1-dioxo-1,4-dihydro-2H-1 λ^6 -benzo[1,2,4]thiadiazin-3-one

To a solution of 7-Fluoro-1,1-dioxo-1,4-dihydro-2H-1- λ^6 -benzo[1,2,4]thiadiazin-3-one (5.19g, 24.0mmol) in DMF (50mL), sodium hydride (1.04g, 26.0mmol) and iodomethane (1.5mL, 24.0mmol) are added successively and the mixture is stirred for 1 hour at 70°C. After cooling to room temperature, the mixture is poured into water and the precipitate is collected by a filtration and washed with water and hexane, successively, to give slightly gray solids (5.38g, 94%).

NMR (400MHz, DMSO-d₆, δ): 3.32 (s, 3H), 7.44 (dd, 1H), 7.75 (ddd, 1H), 7.94 (dd, 1H).

Rf (MeOH:AcOEt = 1:5): 0.21. Rf: 0.39 (Hexane:AcOEt=1:1).

Preparation of 2-Amino-5-fluoro-N-methyl-benzenesulfonamide

6.79g of 7-Fluoro-2-methyl-1,1-dioxo-1,4-dihydro-2H-1 λ^6 -benzo[1,2,4]thiadiazin-3-one (29.5mmol) is dissolved in 20% aq. sodium hydroxide and the resulting solution is stirred at 100°C for 13.5 hours. The mixture is cooled to room temperature and poured into water. 78mL of 5M HCl aq. is added and the precipitate is collected by a filtration and washed with water to afford slightly purple solids (3.96g, 65%).

NMR (400MHz, CDCl₃, δ): 2.60 (d, 3H), 4.55-4.82 (m, 3H), 6.74 (dd, 1H), 7.05-7.12 (m, 1H), 7.45 (dd, 1H). Rf: 0.41 (Hexane:AcOEt=1:1).

2-(5-Bromo-2-chloro-pyrimidin-4-ylamino)-5-fluoro-N-methyl-benzenesulfonamide

The reaction of pyrimidine with 2-Amino-5-fluoro-N-methyl-benzenesulfonamide is performed in the same manner described in example B.

NMR (400MHz, CDCl_3 , δ): 2.67 (d, 3H), 4.56 (m, 1H), 7.36-7.45 (m, 1H), 7.68 (dd, 1H), 8.39 (s, 1H), 8.42 (dd, 1H), 9.26 (s, 1H). Rf 0.59 (Hexane:AcOEt = 1:1).

2-[5-Bromo-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-5-fluoro-N-methyl-benzenesulfonamide

The introduction of substituted aniline is performed according to the manner described in Example A.

NMR (400MHz, CDCl_3 , δ): 2.65 (d, 3H), 3.09-3.16 (m, 4H), 3.87 (s, 3H), 4.50 (q, 1H), 6.41 (dd, 1H), 6.52 (d, 1H), 7.25-7.33 (m, 2H), 7.69 (dd, 1H), 7.95 (d, 1H), 8.20 (s, 1H), 8.37 (dd, 1H), 8.70 (s, 1H). Rf 0.30 (Hexane:AcOEt = 1:1)

Example 53:

Compound

Structure

Physical Data


d

^1H NMR 400 MHz ($\text{DMSO}-d_6$) and/or MS

Number


(m/z)

5



^1H NMR 400 MHz (CDCl_3) δ 12.5 (s, br, 1H), 10.3 (s, 1H), 8.39 (d, 1H, $J = 8.38$ Hz), 7.97 (s, 1H), 7.87 (dd, 1H, $J = 1.63$, 7.55 Hz), 7.49 (d, 1H, $J = 8.69$ Hz), 6.54 (d, 1H, $J = 2.49$ Hz), 6.43 (dd, 1H, $J = 2.51$, 8.73 Hz), 4.70 (s, 1H), 4.04 (q, 2H, $J = 6.98$ Hz), 3.51 (m, 2H), 3.29 (d, 1H, $J = 19.2$ Hz), 3.18 (m, 2H), 2.89 (d, 3H, $J = 3.26$ Hz), 2.77 (s, 6H), 2.39 (m, 2H), 2.22 (m, 2H), 1.40 (t, 3H, $J = 6.97$ Hz). MS m/z 561.4 ($M + 1$)


6




MS m/z 589.4 ($M + 1$)

- 161 -


7

MS m/z 633.4/635.4 ($M + 1$).


8

MS m/z 591.3/593.3 ($M + 1$).

9


MS m/z 547.4 ($M + 1$).

10




^1H NMR 400 MHz (acetone- d_6) δ 9.45 (br, 1H), 8.88 (br, 1H), 8.23 (m, 1H), 8.07 (s, 1H), 7.83 (m, 1H), 7.72 (dd, 1H, $J = 1.25$, 7.96 Hz), 7.45 (m, 1H), 7.14 (m, 2H), 6.86 (m, 1H), 6.79 (br, 1H), 6.62 (m, 1H), 6.13 (m, 1H), 3.97 (t, 2H, $J = 6.98$ Hz), 3.63 (m, 2H), 3.21 (m, 2H), 2.44 (m, 1H), 2.34 (s, 3H), 1.99 (m, 4H), 1.17 (t, 3H, $J = 6.98$ Hz). MS m/z 560.4 ($M + 1$).

11


MS m/z 561.4 ($M + 1$).

12


MS m/z 591.3/593.3 ($M + 1$).

- 162 -


13

MS m/z 547.4 ($M + 1$).


14

MS m/z 559.4 ($M + 1$).


15

MS m/z 564.5 ($M + 1$).

16


MS m/z 575.4 ($M + 1$).

17


MS m/z 605.3/607.3 ($M + 1$).

- 163 -


18

MS m/z 561.4 ($M + 1$).

19


MS m/z 532.4 ($M + 1$).

20




^1H NMR 400 MHz (acetone- d_6) δ 8.60 (d, 1H, $J = 8.27$ Hz), 8.40 (s, 1H), 8.00 (dd, $J = 1.55, 7.95$ Hz), 7.76 (m, 2H), 7.49 (m, 1H), 7.08 (d, $J = 8.94$ Hz, 1H), 6.84 (dd, $J = 3.00, 8.91$ Hz, 1H), 4.50 (t, $J = 4.99$ Hz, 2H), 4.08 (m, 2H), 4.00 (m, 2H), 3.90 (s, 3H), 3.78 (m, 4H), 3.37 (m, 4H), 1.76 (m, 2H), 0.99 (t, $J = 7.44$ Hz, 3H). MS m/z 562.4 ($M + 1$).

21


MS m/z 560.4 ($M + 1$).

22




^1H NMR 600 MHz (acetone- d_6) δ 11.20 (br, 1H), 9.75 (d, $J = 15.25$ Hz, 1H), 8.46 (m, 2H), 8.04 (dd, $J = 1.12, 7.93$ Hz, 1H), 7.71 (m, 1H), 7.69 (m, 1H), 7.47 (t, $J = 7.63$ Hz, 1H), 7.06 (m, 1H), 7.02 (m, 1H), 4.07 (m, 2H), 3.99 (s, 3H), 3.77 (m, 2H), 3.67 (m, 1H), 3.27 (m, 2H), 3.13 (t, $J = 12.2$ Hz, 2H), 2.85 (m, 2H), 2.46 (m, 2H), 2.26 (m, 3H), 2.08 (m, 3H), 1.95 (m, 1H), 1.75 (m, 1H), 1.66 (m, 1H), 0.89 (d, $J = 6.65$ Hz, 6H). MS m/z 672.4/674.4 ($M + 1$).


23

MS m/z 600.4 ($M + 1$).


24

MS m/z 586.4 ($M + 1$).


25

MS m/z 642.5 ($M + 1$).


26

MS m/z 640.5 ($M + 1$).


27

MS m/z 630.3/632.3 ($M + 1$).


28

MS m/z 633.4/635.3 ($M + 1$).


29

MS m/z 603.4 ($M + 1$).


30

MS m/z 601.4 ($M + 1$).


31

MS m/z 603.4 ($M + 1$).


32

MS m/z 601.4 ($M + 1$).


33

MS m/z 575.4 ($M + 1$).

34


MS m/z 561.4 ($M + 1$).

35




^1H NMR 600 MHz (acetone- d_6) δ 8.47(d, J = 7.13, 1H), 8.43 (s, 1H), 8.09 (d, J = 7.85 Hz, 1H), 7.83 (m, 1H), 7.73 (t, J = 7.67 Hz, 1H), 7.50 (m, 1H), 6.87 (m, 1H), 6.71 (m, 1H), 5.46 (d, J = 31.2 Hz, 1H), 4.22 (m, 3H), 3.55 (m, 5H), 2.51 (m, 1H), 2.49 (m, 1H), 2.21 (m, 1H), 1.50 (m, 6H), 1.10 (d, J = 6.56 Hz, 6H) .MS m/z 575.4 ($M + 1$).


36

MS m/z 574.4 ($M + 1$).


37

MS m/z 562.4 ($M + 1$).


38

MS m/z 560.4 ($M + 1$).


39

MS m/z 546.4 ($M + 1$).


40

MS m/z 560.4 ($M + 1$).


41

MS m/z 559.4 ($M + 1$).


42

MS m/z 547.4 ($M + 1$).


43

MS m/z 559.4 ($M + 1$).


44

MS m/z 600.4 ($M + 1$).


45

MS m/z 586.4 ($M + 1$).


46

MS m/z 600.4 ($M + 1$).


47

MS m/z 599.4 ($M + 1$).

48


MS m/z 587.4 ($M + 1$).

49




^1H NMR 600 MHz (acetone- d_6) δ 10.35 (m, 1H), 8.72 (s, br, 1H), 8.42 (s, 1H), 8.00 (d, $J = 7.93$ Hz, 1H), 7.73 (m, 1H), 7.64 (m, 1H), 7.50 (t, $J = 7.77$ Hz, 1H), 6.55 (s, 1H), 6.43 (d, $J = 8.51$ Hz, 1H), 4.31 (m, 1H), 4.22 (q, $J = 6.93$ Hz, 2H), 3.97 (m, 2H), 3.80 (t, $J = 8.45$ Hz, 1H), 3.58 (q, $J = 8.38$ Hz, 1H), 3.36 (m, 1H), 3.17 (s, 1H), 2.78 (m, 1H), 2.66 (m, 1H), 2.15 (m, 2H), 1.73 (d, $J = 12.9$ Hz, 2H), 1.59 (d, $J = 12.9$ Hz, 1H), 1.55 (m, 2H), 1.43 (m, 3H), 1.35 (m, 2H), 1.25 (m, 1H). MS m/z 599.4 ($M + 1$).

50




MS m/z 562.4 ($M + 1$).

51




MS m/z 614.5 ($M + 1$).

52




MS m/z 640.5 ($M + 1$).


53

MS m/z 640.5 ($M + 1$).


54

MS m/z 628.5 ($M + 1$).


55

MS m/z 600.4 ($M + 1$).


56

MS m/z 644.4/646.4 ($M + 1$).


57

MS m/z 613.5 ($M + 1$).


58

MS m/z 578.5 ($M + 1$).


59

MS m/z 560.4 ($M + 1$).


60

MS m/z 586.4 ($M + 1$).


61

MS m/z 586.4 ($M + 1$).


62

MS m/z 574.4 ($M + 1$).


63

MS m/z 546.4 ($M + 1$).


64

MS m/z 590.4/592.4 ($M + 1$).

65


MS m/z 559.4 ($M + 1$).

66




^1H NMR 600 MHz (CD_3OD) δ 8.65 (br, 1H), 8.03 (br, 1H), 7.78 (d, $J = 7.76$ Hz, 1H), 7.49 (br, 1H), 7.30 (s, 1H), 6.38 (m, 1H), 6.33 (dd, $J = 1.69, 8.64$ Hz, 1H), 4.12 (q, $J = 6.96$ Hz, 2H), 4.06 (m, 1H), 3.71 (dd, $J = 6.68, 10.74$ Hz, 1H), 3.66 (m, 1H), 3.56 (dd, $J = 4.71, 10.76$ Hz, 1H), 3.42 (m, 3H), 3.22 (q, $J = 7.28$ Hz, 2H), 2.57 (m, 1H), 2.26 (m, 1H), 1.39 (t, $J = 7.28$ Hz, 3H), 1.34 (t, $J = 6.96$ Hz, 3H). MS m/z 524.4 ($M + 1$).

67


MS m/z 574.4 ($M + 1$).

68




^1H NMR 600 MHz (CD_3OD) δ 8.23 (s, 1H), 8.18 (d, $J = 8.16$ Hz, 1H), 7.98 (d, $J = 8.44$ Hz, 2H), 7.68 (m, 1H), 7.43 (m, 1H), 7.22 (d, $J = 2.29$ Hz, 1H), 6.97 (dd, $J = 2.30$, 8.81 Hz, 1H), 4.20 (q, $J = 6.99$ Hz, 2H), 3.77 (m, 2H), 3.58 (m, 2H), 2.75 (d, $J = 7.03$ Hz, 2H), 2.72 (m, 1H), 2.18 (m, 4H), 1.47 (t, $J = 6.97$ Hz, 3H), 0.75 (m, 1H), 0.30 (m, 2H), 0.00 (m, 2H). MS m/z 600.4 ($M + 1$).


69

MS m/z 600.4 ($M + 1$).


70

MS m/z 588.4 ($M + 1$).


71

MS m/z 560.4 ($M + 1$).


72

MS m/z 604.3/606.3 ($M + 1$).


73

MS m/z 573.4 ($M + 1$).


74

MS m/z 538.4 ($M + 1$).


75

MS m/z 573.4 ($M + 1$).


76

MS m/z 578.5 ($M + 1$).


77

MS m/z 613.5 ($M + 1$).


78

MS m/z 511.4 ($M + 1$).


79

MS m/z 525.5 ($M + 1$).


80

MS m/z 524.5 ($M + 1$).


81

MS m/z 510.5 ($M + 1$).


82

MS m/z 525.4 ($M + 1$).

83


MS m/z 627.5 ($M + 1$).

84




^1H NMR 600 MHz (CD_3OD) δ 9.18 (m, 2H), 8.05 (d, $J = 7.93$ Hz, 1H), 7.93 (t, $J = 7.64$ Hz, 1H), 7.78 (m, 1H), 7.68 (m, 2H), 7.58 (t, $J = 7.54$ Hz, 1H), 4.72 (s, 2H), 4.07 (d, $J = 12.4$ Hz, 2H), 3.34 (br, 4H), 2.96 (t, $J = 12.3$ Hz, 2H), 2.87 (s, 6H), m, 2H), 2.28 (s, 1H), 2.00 (br, 7H), 1.81 (br, 2H).
MS m/z 627.5 ($M + 1$).


85

MS m/z 591.5 ($M + 1$).


86

MS m/z 601.5 ($M + 1$).


87

MS m/z 602.4 ($M + 1$).


88

MS m/z 566.5 ($M + 1$).

89


MS m/z 656.5 ($M + 1$).

90




^1H NMR 600 MHz (CD_3OD) δ 8.16 (d, J = 8.16 Hz, 1H), 8.23 (s, 1H), 7.94 (dd, J = 1.42, 7.98 Hz, 1H), 7.79 (d, J = 9.24 Hz, 1H), 7.72 (m, 1H), 7.48 (m, 1H), 7.15 (d, J = 2.53 Hz, 1H), 7.03 (dd, J = 2.58, 8.91 Hz, 1H), 4.00 (s, 1H), 3.82 (d, J = 12.63 Hz, 2H), 3.50 (s, br, 4H), 3.29 (m, 5H), 3.19 (m, 1H), 2.96 (s, 3H), 2.73 (s, 6H), 2.25 (d, J = 12.40 Hz, 2H), 2.03 (m, 2H), 1.40 (s, 9H). MS m/z 657.5 ($M + 1$).

91

MS m/z 621.6 ($M + 1$).


92




^1H NMR 600 MHz (CD_3OD) δ 8.31 (d, J = 7.65 Hz, 1H), 8.19 (s, 1H), 7.94 (dd, J = 1.33, 7.96 Hz, 1H), 7.67 (t, J = 7.70 Hz, 1H), 7.49 (t, J = 7.68 Hz, 1H), 7.44 (d, J = 8.56 Hz, 1H), 6.83 (d, J = 1.37 Hz, 1H), 6.65 (d, J = 8.66 Hz, 1H), 4.69 (m, 1H), 4.11 (br, 2H), 3.91 (d, J = 12.68 Hz, 2H), 3.83 (br, 2H), 3.58 (br, 2H), 3.46 (m, 1H), 3.23 (br, 2H), 3.00 (t, J = 12.19 Hz, 2H), 2.73 (s, 6H), 2.33 (d, J = 12.14 Hz, 2H), 1.97 (m, 2H), 1.30 (d, J = 6.04 Hz, 6H).

MS m/z 630.5 ($M + 1$).MS m/z 629.5 ($M + 1$).


93



94


MS m/z 594.5 ($M + 1$).

95

MS m/z 471.1 ($M + 1$).


^1H NMR 600 MHz ($\text{DMSO}-d_6$) δ 11.7 (bs, 1H), 11.14 (bs, 1H), 9.50 (b, 2H), 7.89 (d, 1H, $J = 9.1$ Hz), 7.72 (m, 2H), 7.55 (m, 2H), 7.18 (m, 1H), 6.62 (s, 1H), 6.46 (m, 1H), 4.06 (q, 2H, $J = 6.6$ Hz), 3.73 (t, 4H, $J = 4.8$ Hz), 3.11 (m, 4H), 2.40 (m, 3H), 1.24 (m, 3H); MS m/z 485.1 ($M + 1$).

97


MS m/z 577.1 ($M + 1$).

^1H NMR 600 MHz ($\text{DMSO}-d_6$) δ 9.97 (bs, 1H), 9.34 (s, 1H), 8.48 (bs, 1H), 8.34 (bs, 1H), 8.28 (s, 1H), 7.80 (q, 1H, $J = 4.8$ Hz), 7.77 (d, 1H, $J = 7.8$ Hz), 7.48 (m, 1H), 7.39 (d, 1H, $J = 8.4$ Hz), 7.25 (t, 1H, $J = 7.2$ Hz), 6.71 (s, 1H), 6.49 (d, 1H, $J = 8.4$ Hz), 3.93 (t, 2H, $J = 6.0$ Hz), 3.85 (d, 2H, $J = 12$ Hz), 3.54 (d, 2H, $J = 12$ Hz), 3.18 (q, 2H, $J = 9.0$ Hz), 2.94 (t, 2H, $J = 11$ Hz), 2.88 (s, 3H), 2.42 (d, 3H, $J = 4.8$ Hz), 1.64 (m, 2H), 0.85 (t, 3H, $J = 7.2$ Hz); MS m/z 590.1 ($M + 1$).


98




99

MS m/z 578.1 ($M + 1$).

100


MS m/z 593.1 ($M + 1$).

101




^1H NMR 600 MHz (DMSO- d_6) δ 10.0 (bs, 1H), 9.34 (s, 1H), 8.47 (bs, 1H), 8.33 (bs, 1H), 8.30 (s, 1H), 7.79 (s, 1H), 7.78 (d, 1H, $J = 7.2$ Hz), 7.50 (m, 2H), 7.27 (t, 1H, $J = 7.8$ Hz), 6.75 (s, 1H), 6.53 (d, 1H, $J = 7.8$ Hz), 4.12 (t, 2H, $J = 4.8$ Hz), 3.84 (d, 2H, $J = 12$ Hz), 3.59 (t, 2H, $J = 4.8$ Hz), 3.53 (d, 2H, $J = 12$ Hz), 3.23 (s, 3H), 3.17 (q, 2H, $J = 9.0$ Hz), 2.94 (t, 2H, $J = 11$ Hz), 2.88 (s, 3H), 2.43 (d, 3H, $J = 4.8$ Hz); MS m/z 606.1 ($M + 1$).

102




MS m/z 594.1 ($M + 1$).

103




MS m/z 547.1 ($M + 1$).

104




MS m/z 533.1 ($M + 1$).

105




MS m/z 496.4 ($M + 1$).

106




^1H NMR 400 MHz (DMSO- d_6) δ 9.35 (s, 1H), 9.02 (s, 1H), 8.52 (s, 1H), 8.17 (bs, 1H), 7.80 (m, 2H), 7.45 (bs, 1H), 7.29 (m, 2H), 6.66 (s, 1H), 6.47 (d, 1H, $J = 8.0$ Hz), 4.02 (q, 2H, $J = 7.2$ Hz), 3.76 (t, 4H, $J = 4.8$ Hz), 3.14 (t, 4H, $J = 4.8$ Hz), 2.41 (d, 3H, $J = 4.0$ Hz), 1.23 (t, 3H, $J = 7.2$ Hz); MS m/z 510.4 ($M + 1$).

107




MS m/z 524.5 ($M + 1$).

108




^1H NMR 600 MHz (DMSO- d_6) δ 9.61 (s, 1H), 9.10 (bs, 1H), 8.35 (bs, 1H), 8.18 (s, 1H), 7.77 (m, 2H), 7.41 (bs, 1H), 7.26 (t, 1H, $J = 7.2$ Hz), 7.15 (d, 1H, $J = 7.8$ Hz), 6.89 (s, 1H), 6.85 (d, 1H, $J = 9.6$ Hz), 3.77 (t, 4H, $J = 4.8$ Hz), 3.15 (t, 4H, $J = 4.8$ Hz), 2.52 (q, 2H, $J = 7.2$ Hz), 2.41 (d, 3H, $J = 4.8$ Hz), 1.06 (t, 3H, $J = 7.2$ Hz); MS m/z 503.2 ($M + 1$).

109




MS m/z 547.2 ($M + 1$).

110




MS m/z 494.2 ($M + 1$).

111




MS m/z 557.3 ($M + 1$).

112




MS m/z 559.3 ($M + 1$).

113




MS m/z 561.2 ($M + 1$).

114




MS m/z 508.2 ($M + 1$).

115




MS m/z 529.2 ($M + 1$).

116




^1H NMR 400 MHz (DMSO- d_6) δ 9.91 (bs, 1H), 9.50 (s, 1H), 8.94 (s, 1H), 8.49 (bs, 1H), 8.17 (s, 1H), 7.74 (d, 1H, $J = 7.6$ Hz), 7.41 (bs, 1H), 7.23 (t, 1H, $J = 7.2$ Hz), 7.18 (d, 1H, $J = 8.0$ Hz), 6.89 (m, 2H), 3.84 (d, 2H, $J = 11.4$ Hz), 3.54 (d, 2H, $J = 11.4$ Hz), 3.18 (q, 2H, $J = 8.0$ Hz), 2.95 (t, 2H, $J = 11.4$ Hz), 2.88 (s, 3H), 2.64 (s, 6H), 2.53 (m, 2H), 1.06 (t, 3H, $J = 8.0$ Hz); MS m/z 530.2 ($M + 1$).

117




MS m/z 556.2 ($M + 1$).

118




MS m/z 556.2 ($M + 1$).

119




MS m/z 548.2 ($M + 1$).

120




^1H NMR 400 MHz (DMSO- d_6) δ 10.24 (bs, 1H), 9.54 (s, 1H), 8.50 (bs, 1H), 8.41 (d, 1H, $J = 7.2$ Hz), 8.21 (s, 1H), 7.80 (m, 2H), 7.50 (t, 1H, $J = 7.2$ Hz), 7.32 (d, 1H, $J = 8.4$ Hz), 7.28 (t, 1H, $J = 8.0$ Hz), 6.32 (d, 1H, $J = 2.4$ Hz), 6.17 (dd, 1H, $J_1 = 2.4$ Hz, $J_2 = 8.4$ Hz), 4.61 (m, 2H), 4.01 (m, 1H), 3.60 (m, 1H), 3.48 (m, 2H), 3.25 (q, 1H, $J = 8.8$ Hz), 2.88 (s, 6H), 2.42 (d, 3H, $J = 4.8$ Hz), 2.19 (m, 1H), 1.20 (d, 6H, $J = 4.8$ Hz); MS m/z 560.2 ($M + 1$).

121













MS m/z 600.2 ($M + 1$).


122




MS m/z 562.2 ($M + 1$).

- 123  MS m/z 574.2 ($M + 1$).
- 124  MS m/z 575.2 ($M + 1$).
- 125  MS m/z 600.2 ($M + 1$).
- 126  MS m/z 574.2 ($M + 1$).
- 127  MS m/z 573.2 ($M + 1$).
- 128  MS m/z 604.1 ($M + 1$).
- 129  MS m/z 576.2 ($M + 1$).
- 130  MS m/z 588.2 ($M + 1$).
- 131  MS m/z 561.2 ($M + 1$).
- 132  MS m/z 561.1 ($M + 1$).


133

MS m/z 589.2 ($M + 1$).


134

MS m/z 575.2 ($M + 1$).


135

MS m/z 574.2 ($M + 1$).


136

MS m/z 601.2 ($M + 1$).


137

MS m/z 560.2 ($M + 1$).

138


MS m/z 575.2 ($M + 1$).

139




^1H NMR 600 MHz ($\text{DMSO}-d_6$) δ 10.35 (bs, 1H), 9.45 (s, 1H), 8.47 (d, 1H, $J = 7.2$ Hz), 8.33 (s, 1H), 8.23 (s, 1H), 7.79 (d, 1H, $J = 8.4$ Hz), 7.57 (t, 1H, $J = 7.2$ Hz), 7.52 (d, 1H, $J = 9.6$ Hz), 7.23 (t, 1H, $J = 7.2$ Hz), 6.68 (s, 0.4H), 6.64 (s, 0.6H), 6.50 (s, 0.6H), 6.49 (s, 0.4H), 5.19 (s, 0.6H), 5.15 (s, 0.4H), 4.58 (m, 1H), 3.99 (m, 0.4H), 3.73 (m, 1.6H), 3.36 (m, 0.6H), 3.26 (m, 0.8H), 3.15 (m, 0.6H), 2.96 (s, 1.2H), 2.89 (s, 1.8H), 2.64 (s, 6H), 2.60 (m, 1H), 2.24 (m, 0.6H), 2.07 (m, 0.4H), 1.20 (d, 6H, $J = 6.0$ Hz); MS m/z 561.2 ($M + 1$).


140

MS m/z 587.2 ($M + 1$).


141

MS m/z 587.2 ($M + 1$).


142

MS m/z 560.2 ($M + 1$).


143

MS m/z 575.2 ($M + 1$).


144

MS m/z 561.2 ($M + 1$).


145

MS m/z 587.2 ($M + 1$).

146


MS m/z 587.2 ($M + 1$).

147




^1H NMR 600 MHz ($\text{DMSO}-d_6$) δ 9.81 (bs, 0.4H), 9.74 (bs, 0.6H), 9.59 (bs, 1H), 8.48 (bs, 1H), 8.34 (s, 0.4H), 8.33 (s, 0.6H), 8.232 (s, 0.6H), 8.228 (s, 0.4H), 7.82 (d, 1H, $J = 7.8$ Hz), 7.59 (m, 1H), 7.49 (d, 0.6H, $J = 8.4$ Hz), 7.47 (d, 0.4H, $J = 8.4$ Hz), 7.34 (q, 1H, $J = 7.2$ Hz), 6.75 (s, 1H), 6.68 (s, 1H), 6.57 (dd, 0.6H, $J_1 = 2.4$ Hz, $J_2 = 8.4$ Hz), 6.54 (dd, 0.4H, $J_1 = 2.4$ Hz, $J_2 = 8.4$ Hz), 4.73 (m, 0.6H), 4.58 (m, 1H), 4.52 (m, 0.4H), 3.52 (m, 1H), 3.43 (m, 1H), 3.35 (m, 0.4H), 3.33 (m, 0.6H), 3.21 (m, 1H), 3.11 (m, 1H), 2.85 (d, 1.8H, $J = 4.8$ Hz), 2.82 (d, 1.2H, $J = 3.0$ Hz), 2.25 (m, 1H), 2.07 (m, 0.4H), 2.05 (m, 0.6H), 1.98 (m, 1H), 1.76 (m, 1H), 1.20 (d, 3.6H, $J = 6.0$ Hz), 1.18 (d, 2.4H, $J = 6.0$ Hz), 1.15 (d, 6H, $J = 7.2$ Hz); MS m/z 574.2 ($M + 1$).


148

MS m/z 589.3 ($M + 1$).


149

MS m/z 575.2 ($M + 1$).


150

MS m/z 601.3 ($M + 1$).


151

MS m/z 601.3 ($M + 1$).

152


MS m/z 559.1 ($M + 1$).

153




^1H NMR 400 MHz ($\text{DMSO}-d_6$) δ 10.07 (bs, 1H), 9.44 (s, 1H), 8.54 (s, 1H), 8.52 (bs, 1H), 8.29 (s, 1H), 7.91 (s, 1H), 7.79 (d, 1H, $J = 8.0$ Hz), 7.61 (t, 1H, $J = 7.6$ Hz), 7.32 (t, 1H, $J = 7.6$ Hz), 7.22 (d, 1H, $J = 8.8$ Hz), 7.14 (d, 1H, $J = 8.8$ Hz), 4.10 (m, 2H), 3.86 (s, 3H), 3.46 (m, 2H), 3.22 (m, 2H), 3.08 (m, 2H), 2.83 (s, 3H), 2.66 (s, 6H); MS m/z 560.1 ($M + 1$).


154

MS m/z 586.1 ($M + 1$).


155

MS m/z 586.1 ($M + 1$).

156


MS m/z 587.2 ($M + 1$).

157




^1H NMR 400 MHz (DMSO- d_6) δ 10.05 (bs, 1H), 9.42 (s, 1H), 8.43 (d, 1H, $J = 7.6$ Hz), 8.35 (s, 1H), 8.31 (s, 1H), 7.91 (s, 1H), 7.79 (dd, 1H, $J_1 = 1.6$ Hz, $J_2 = 8.0$ Hz), 7.58 (t, 1H, $J = 8.0$ Hz), 7.32 (t, 1H, $J = 8.4$ Hz), 7.15 (m, 2H), 4.70 (m, 1H), 4.07 (m, 2H), 3.45 (m, 2H), 3.21 (m, 2H), 3.06 (m, 2H), 2.83 (s, 3H), 2.65 (s, 6H), 1.27 (d, 6H, $J = 6.0$ Hz); MS m/z 588.2 ($M + 1$).

158




MS m/z 614.2 ($M + 1$).

159




MS m/z 573.2 ($M + 1$).

160




^1H NMR 600 MHz (DMSO- d_6) δ 9.45 (s, 1H), 8.44 (d, 1H, $J = 7.8$ Hz), 8.38 (s, 1H), 8.27 (s, 1H), 7.81 (d, 1H, $J = 7.8$ Hz), 7.66 (d, 1H, $J = 8.4$ Hz), 7.64 (t, 1H, $J = 7.2$ Hz), 7.35 (t, 1H, $J = 7.2$ Hz), 6.97 (s, 1H), 6.76 (d, 1H, $J = 8.4$ Hz), 4.06 (q, 2H, $J = 7.2$ Hz), 3.59 (m, 6H), 3.29 (m, 4H), 2.92 (m, 2H), 2.86 (s, 3H), 2.64 (s, 6H), 1.31 (t, 3H, $J = 6.6$ Hz); MS m/z 574.2 ($M + 1$).

161




MS m/z 600.2 ($M + 1$).

162




MS m/z 600.2 ($M + 1$).

163




MS m/z 627.5 ($M + 1$).

164




MS m/z 628.5 ($M + 1$).

165




MS m/z 654.5 ($M + 1$).

166




MS m/z 654.5 (M + 1).

167




MS m/z 592.3 ($M + 1$).

168




MS m/z 588.2 ($M + 1$).

169




MS m/z 614.2 (M + 1).

170




MS m/z 574.2 (M + 1).

171




¹H NMR 600 MHz (DMSO-*d*₆) δ 9.66 (s, 1H), 8.48 (bs, 1H), 8.40 (d, 1H, J = 8.8 Hz), 8.30 (s, 1H), 7.66 (dd, 1H, J = 1.6 Hz, J₂ = 7.6 Hz), 7.71 (t, 1H, J = 8.4 Hz), 7.46 (s, 1H), 7.41 (t, 1H, J = 8.0 Hz), 7.17 (bs, 1H), 6.96 (s, 1H), 6.93 (bs, 1H), 4.67 (m, 1H), 4.02 (q, 1H, J = 7.2 Hz), 3.65 (m, 2H), 3.46 (m, 1H), 3.34 (m, 1H), 2.54 (m, 1H), 1.94 (m, 4H), 1.26 (d, 6H, J = 6.0 Hz), 1.15 (d, 6H, J = 6.0 Hz); MS *m/z* 587.2 (M + 1).

172




^1H NMR 600 MHz ($\text{DMSO}-d_6$) δ 9.63 (bs, 1H), 8.50 (bs, 1H), 8.46 (bs, 1H), 8.29 (s, 1H), 7.82 (d, 1H, $J = 7.2$ Hz), 7.61 (t, 1H, $J = 7.2$ Hz), 7.43 (d, 1H, $J = 8.4$ Hz), 7.34 (t, 1H, $J = 7.2$ Hz), 6.75 (s, 1H), 6.58 (d, 1H, $J = 6.6$ Hz), 4.04 (q, 2H, $J = 7.2$ Hz), 3.82 (m, 2H), 3.76 (m, 5H), 3.44 (m, 1H), 3.22 (m, 4H), 2.85 (s, 3H), 2.82 (m, 2H), 2.09 (m, 2H), 1.71 (m, 2H), 1.25 (t, 3H, $J = 7.2$ Hz), 1.16 (d, 6H, $J = 7.2$ Hz); MS m/z 628.2 ($M + 1$).

173




MS m/z 615.2 ($M + 1$).

174




MS m/z 659.1 ($M + 1$).

175




^1H NMR 600 MHz ($\text{DMSO}-d_6$) δ 9.43 (s, 1H), 8.45 (bs, 1H), 8.38 (bs, 1H), 8.24 (s, 1H), 7.91 (d, 1H, $J = 8.4$ Hz), 7.84 (d, 1H, $J = 8.4$ Hz), 7.51 (t, 1H, $J = 7.2$ Hz), 7.48 (d, 1H, $J = 8.4$ Hz), 7.26 (t, 1H, $J = 7.2$ Hz), 6.76 (s, 1H), 6.55 (d, 1H, $J = 7.2$ Hz), 4.05 (q, 2H, $J = 6.6$ Hz), 3.82 (m, 2H), 3.59 (m, 5H), 3.30 (m, 1H), 3.25 (m, 4H), 2.85 (s, 3H), 2.82 (m, 2H), 2.09 (m, 2H), 1.71 (m, 2H), 1.26 (t, 3H, $J = 6.6$ Hz), 0.92 (d, 6H, $J = 7.2$ Hz); MS m/z 643.2 ($M + 1$).

176




MS m/z 642.2 ($M + 1$).

177




MS m/z 643.3 ($M + 1$).

178




MS m/z 669.3 ($M + 1$).


179

MS m/z 633.3 ($M + 1$).


180

MS m/z 629.2 ($M + 1$).


181

MS m/z 669.2 ($M + 1$).


182

MS m/z 642.2 ($M + 1$).

183


MS m/z 615.3 ($M + 1$).

184




^1H NMR 400 MHz ($\text{DMSO}-d_6$) δ 9.94 (bs, 1H), 9.53 (s, 1H), 8.49 (m, 2H), 8.23 (s, 1H), 7.79 (dd, 1H, $J_1 = 2.0$ Hz, $J_2 = 8.0$ Hz), 7.58 (t, 1H, $J = 7.6$ Hz), 7.38 (d, 1H, $J = 9.2$ Hz), 7.32 (t, 1H, $J = 8.0$ Hz), 6.68 (d, 1H, $J = 2.0$ Hz), 6.51 (dd, 1H, $J_1 = 2.0$ Hz, $J_2 = 8.4$ Hz), 4.21 (m, 2H), 4.04 (q, 2H, $J = 6.8$ Hz), 3.86 (d, 2H, $J = 12$ Hz), 3.68 (t, 2H, $J = 12$ Hz), 3.51 (d, 2H, $J = 11.2$ Hz), 3.38 (t, 1H, $J = 12$ Hz), 3.13 (m, 2H), 2.73 (t, 2H, $J = 11.2$ Hz); 2.64 (s, 6H), 2.16 (d, 2H, $J = 11.2$ Hz), 1.73 (m, 2H), 1.24 (t, 3H, $J = 7.2$ Hz); MS m/z 616.3 ($M + 1$).


185

MS m/z 642.3 ($M + 1$).


186

MS m/z 606.7 ($M + 1$).


187

MS m/z 599.3 ($M + 1$).


188

MS m/z 600.3 ($M + 1$).


189

MS m/z 626.3 ($M + 1$).


190

MS m/z 590.3 ($M + 1$).

191


MS m/z 599.3 ($M + 1$).

192




^1H NMR 400 MHz ($\text{DMSO}-d_6$) δ 9.86 (bs, 1H), 9.56 (s, 1H), 8.59 (bs, 1H), 8.47 (bs, 1H), 8.24 (s, 1H), 7.79 (dd, 1H, $J_1 = 1.6$ Hz, $J_2 = 8.0$ Hz), 7.58 (t, 1H, $J = 8.0$ Hz), 7.38 (d, 1H, $J = 8.4$ Hz), 7.33 (t, 1H, $J = 7.2$ Hz), 6.70 (d, 1H, $J = 2.4$ Hz), 6.51 (dd, 1H, $J_1 = 2.4$ Hz, $J_2 = 8.4$ Hz), 4.03 (q, 2H, $J = 6.8$ Hz), 3.81 (d, 2H, $J = 12.4$ Hz), 3.58 (m, 2H), 3.28 (m, 1H), 3.11 (m, 2H), 2.74 (t, 2H, $J = 12$ Hz), 2.64 (s, 6H), 2.15 (d, 2H, $J = 11.6$ Hz), 2.03 (m, 2H), 1.86 (m, 2H), 1.73 (m, 2H), 1.24 (t, 3H, $J = 6.8$ Hz); MS m/z 600.2 ($M + 1$).

193


MS m/z 626.2 ($M + 1$).

194




^1H NMR 400 MHz (CD_3OD) δ 8.29 (s, br, 1H), 8.07 (s, br, 1H), 7.96 (d, 1H, $J=7.8$ Hz), 7.60 (m, 1H), 7.39 (t, 1H, $J=7.2$ Hz), 7.22 (s, br, 1H), 6.31 (s, 1H), 6.22 (s, br, 1H), 4.08 (q, 1H, $J=6.6$ Hz), 3.95 (p, 1H, $J=6.0$ Hz), 3.61 (m, 2H), 3.53 (dd, 1H, $J=4.2, 11.4$ Hz), 3.36 (m, 2H), 2.78 (s, 3H), 2.52 (m, 1H), 2.22 (m, 1H), 1.31 (t, 3H, $J=6.6$ Hz), 0.98 (d, 6H, $J=6.6$ Hz); MS m/z 560.40 ($M + 1$).

195




^1H NMR 400 MHz (CD_3OD) δ 8.30 (s, br, 1H), 8.05 (s, br, 1H), 7.87 (d, 1H, $J=7.8$ Hz), 7.60 (m, 1H), 7.42 (m, 1H), 7.17 (s, br, 1H), 6.28 (s, 1H), 6.19 (s, br, 1H), 4.04 (q, 2H, $J=6.6$ Hz), 3.91 (p, 1H, $J=5.4$ Hz), 3.60 (m, 2H), 3.50 (dd, 1H, $J=3.6, 10.8$ Hz), 3.33 (m, 1H), 2.75 (s, 3H), 2.66 (s, 6H), 2.48 (m, 1H), 2.20 (m, 1H), 1.27 (t, 3H, $J=6.6$ Hz); MS m/z 546.40 ($M + 1$).

196




MS m/z 572.40 ($M + 1$).

197




MS m/z 545.40 ($M + 1$).

198




MS m/z 545.40 ($M + 1$).

199




^1H NMR 400 MHz (CD_3OD) δ 8.37 (s, br 1H), 8.11 (s, br, 1H), 8.00 (d, 1H, $J=7.8$ Hz), 7.65 (s, br, 1H), 7.43 (t, 1H, $J=7.2$ Hz), 7.25 (s, br, 1H), 6.34 (s, 1H), 6.25 (s, 1H), 4.11 (q, 2H, $J=7.2$ Hz), 3.99 (m, 1H), 3.65 (m, 2H), 3.57 (dd, 1H, $J=3.6, 10.8$ Hz), 3.41 (m, 2H), 2.82 (s, 3H), 2.56 (m, 1H), 2.28 (m, 1H), 1.35 (t, 3H, $J=6.6$ Hz), 1.01 (d, 6H, $J=6.6$ Hz); MS m/z 560.40 ($M + 1$).

200




^1H NMR 400 MHz (CD_3OD) δ 8.30 (s, br, 1H), 8.10 (s, br, 1H), 7.91 (d, 1H, $J=7.8$ Hz), 7.57 (s, br, 1H), 7.38 (t, 1H, $J=7.2$ Hz), 7.21 (s, 1H), 6.30 (s, 1H), 6.21 (s, 1H), 4.08 (q, 2H, $J=7.2$ Hz), 3.95 (m, 1H), 3.68 (m, 1H), 3.62 (m, 2H), 3.53 (dd, 1H, $J=4.2, 10.8$ Hz), 3.36 (m, 1H), 2.78 (s, 3H), 2.50 (m, 1H), 2.25 (m, 1H), 1.94 (m, 2H), 1.75 (m, 2H), 1.50 (m, 2H), 1.31 (t, 3H, $J=7.2$ Hz); MS m/z 572.40 ($M + 1$).

201




MS m/z 532.40 ($M + 1$).

202




MS m/z 546.40 ($M + 1$).

203




MS m/z 545.40 ($M + 1$).

204


MS m/z 532.40 ($M + 1$).

205




^1H NMR 400 MHz (CD_3OD) δ 8.27 (s, br, 1H), 8.07 (m, 1H), 7.94 (d, 1H, $J=7.8$ Hz), 7.61 (s, br, 1H), 7.40 (t, 1H, $J=7.2$ Hz), 7.20 (s, 1H), 6.30 (s, 1H), 6.20 (s, 1H), 4.08 (q, 2H, $J=7.2$ Hz), 3.95 (m, 1H), 3.60 (m, 2H), 3.53 (dd, 1H, $J=4.2, 10.8$ Hz), 3.35 (m, 1H), 2.78 (s, 3H), 2.73 (d, 1H, $J=6.6$ Hz), 2.51 (m, 1H), 2.23 (m, 1H), 1.31 (t, 3H, $J=7.2$ Hz), 0.742 (s, br 1H), 0.32 (s, 2H), 0.00 (s, 2H); MS m/z 572.40 ($M + 1$).

206




^1H NMR 400 MHz (CD_3OD) δ 8.24 (s, br, 1H), 8.07 (s, br, 1H), 7.93 (d, 1H, $J=8.4$ Hz), 7.57 (s, 1H), 7.38 (t, 1H, $J=7.2$ Hz), 7.20 (s, 1H), 6.29 (s, 1H), 6.19 (s, br, 1H), 4.06 (q, 2H, $J=7.2$ Hz), 3.93 (m, 1H), 3.60 (m, 2H), 3.50 (dd, 1H, $J=6.0, 10.8$ Hz), 3.35 (m, 1H), 2.76 (s, 3H), 2.72 (d, 1H, $J=6.6$ Hz), 2.50 (m, 1H), 2.22 (m, 1H), 1.30 (t, 3H, $J=7.2$ Hz), 0.73 (m, 1H), 0.31 (m, 2H), 0.01 (m, 2H); MS m/z 572.40 ($M + 1$).


207

MS m/z 545.40 ($M + 1$).


208

MS m/z 549.10 ($M + 1$).

209


MS m/z 532.40 ($M + 1$).

210




^1H NMR 400 MHz (CD_3OD) δ 8.22 (d, 1H, $J=7.8\text{Hz}$), 8.20 (s, 1H), 7.92 (dd, 1H, $J=1.2, 7.8\text{Hz}$), 7.58 (t, 1H, $J=6\text{Hz}$), 7.48(d, 1H, $J=9.1\text{Hz}$), 7.38 (t, 1H, $J=7.8\text{Hz}$), 6.65 (s, 1H), 6.46 (d, 1H, $J=7.8\text{Hz}$), 5.20 (s, 1H), 4.06 (q, 2H, $J=7.2\text{Hz}$), 3.86 (m, 1H), 3.38 (m, 1H), 3.05 (m, 3H), 2.64 (m, 1H), 2.50 (s, 3H), 2.30 (m, 1H), 1.34 (t, 3H, $J=7.2\text{Hz}$); MS m/z 577.30 ($M + 1$).

211


MS m/z 533.30 ($M + 1$).

212




^1H NMR 400 MHz (CD_3OD) δ 8.30 (d, 1H, $J=7.8\text{Hz}$), 8.11 (s, 1H), 7.91 (d, 1H, $J=7.8\text{Hz}$), 7.58 (t, 1H, $J=7.8\text{ Hz}$), 7.55 (d, 1H, $J=8.4\text{ Hz}$), 7.36 (t, 1H, $J=7.8\text{ Hz}$), 6.65 (s, 1H), 6.47 (d, 1H, $J=8.4\text{ Hz}$), 5.20 (s, 1H), 4.07 (q, 2H, $J=7.2\text{ Hz}$), 3.87 (m, 1H), 3.38 (m, 1H), 3.23 (m, 1H), 3.00 (m, 2H), 2.64 (m, 1H), 2.50 (s, 3H), 2.38 (m, 1H), 1.35 (t, 3H, $J=7.2\text{Hz}$); MS m/z 533.30 ($M + 1$).

213




MS m/z 533.30 ($M + 1$).

214




MS m/z 519.30 ($M + 1$).

215




MS m/z 563.30 ($M + 1$).


216

MS m/z 505.10 ($M + 1$).


217

MS m/z 563.30 ($M + 1$).


218






MS m/z 563.10 ($M + 1$).






219

MS m/z 561.40 ($M + 1$).


220

MS m/z 577.30 ($M + 1$).







231

MS m/z 519.35 ($M + 1$).

232


MS m/z 532.35 ($M + 1$).

233




^1H NMR 400 MHz (CD_3OD) δ 8.06 (d, 1H, $J=8.0$ Hz), 7.92 (s, 1H), 7.73 (dd, 1H, $J=1.6, 8.0$ Hz), 7.40 (t, 1H, $J=7.6$ Hz), 7.22 (m, 1H), 7.10 (d, 1H, $J=8.8$ Hz), 6.50 (d, 1H, $J=2.4$ Hz), 6.38 (dd, 1H, $J=2.0, 8.8$ Hz), 3.67 (m, 4H), 3.35 (m, 2H), 3.15 (m, 1H), 2.84 (m, 2H), 2.69 (m, 2H), 2.30 (s, 3H), 2.01 (m, 2H), 1.69 (m, 7H), 1.27 (m, 1H), 0.95 (m, 1H), 0.33 (m, 2H), 0.05 (m, 2H); MS m/z 626.5 ($M + 1$).


234

MS m/z 561.4 ($M + 1$).

235


MS m/z 547.25 ($M + 1$).

236




^1H NMR 400 MHz (CD_3OD) δ 8.23 (d, 1H, $J=7.6\text{Hz}$), 8.15 (s, 1H), 7.90 (dd, 1H, $J=1.6, 8.0\text{Hz}$), 7.56 (t, 1H, $J=7.2\text{ Hz}$), 7.36 (t, 1H, $J=7.6\text{ Hz}$), 7.31 (d, 1H, $J=8.8\text{ Hz}$), 6.63 (d, 1H, $J=2.4\text{ Hz}$), 6.51 (m, 1H), 4.10 (m, 2H), 3.83 (m, 6H), 3.34 (m, 2H), 2.82 (m, 2H), 2.48 (s, 3H), 2.23 (m, 2H), 1.78 (m, 2H), 1.14 (m, 1H), 0.51 (m, 2H), 0.23 (m, 2H); MS m/z 672.4 ($M + 1$).


237

MS m/z 532.4 ($M + 1$).


238

MS m/z 546.4 ($M + 1$).


239

MS m/z 576.3 ($M + 1$).


240

MS m/z 590.4 ($M + 1$).






241

MS m/z 533.3 ($M + 1$).


242




^1H NMR 400 MHz (CD_3OD) δ 8.26 (s, 1H), 8.10 (d, 1H, $J=8.0$ Hz), 7.90 (dd, 1H, $J=1.2, 8.0\text{Hz}$), 7.80 (d, 1H, $J=8.4\text{Hz}$), 7.62 (m, 1H), 7.38 (t, 1H, $J=8.0\text{Hz}$), 7.05 (d, 1H, $J=1.6\text{Hz}$), 6.83 (dd, 1H, $J=1.6, 8.4\text{Hz}$), 3.86 (s, 3H), 3.43 (m, 4H), 3.00 (m, 4H), 2.10 (m, 2H), 1.95 (m, 2H), 1.72 (m, 6H), 1.45 (m, 2H); MS m/z 658.4 ($M + 1$).




248

MS m/z .560.2 ($M + 1$).


249






MS m/z .604.3 ($M + 1$).

250

MS m/z .559.4 ($M + 1$).


251

MS m/z .573.4 ($M + 1$).




- 206 -


257

MS m/z .530.4 ($M + 1$).


258

MS m/z .590.4 ($M + 1$).

259


MS m/z .588.2 ($M + 1$).

260


MS m/z .546.4 ($M + 1$).

- 207 -


261

MS m/z .520.4 ($M + 1$).


262

MS m/z .534.4 ($M + 1$).

263


MS m/z .544.2 ($M + 1$).

264


MS m/z .516.4 ($M + 1$).

- 208 -


265

MS m/z .574.2 ($M + 1$).


266

MS m/z .577.1 ($M + 1$).


267

MS m/z .579.3 ($M + 1$)


268

MS m/z .614.5 ($M + 1$)


269

MS m/z .658.4 (M + 1)


270

MS m/z .590.3 (M + 1)


271

MS m/z .546.4 (M + 1)


272

MS m/z .524.4 (M + 1)

273


MS m/z 614.5 ($M + 1$)

274



^1H NMR 400 MHz (CD_3OD) δ 8.17 9s, 1H, 8.15 (d, 1H, $J=8.0\text{Hz}$), 7.89 9dd, 1H, $J=1.2, 8.0\text{ Hz}$), 7.86 (d, 1H, $J=8.4\text{Hz}$), 7.61 (m, 1H), 7.36 (m, 1H), 7.01 (d, 1H, $J=1.6\text{Hz}$), 6.81(dd, 1H, $J=1.6, 8.0\text{Hz}$), 4.10 (q, 2H, $J=7.2\text{Hz}$), 3.45 (m, 3H), 2.94 9m, 4H), 2.44 (s, 3H), 2.08 (m, 2H), 1.94 (m, 2H), 1.75 (m, 6H), 1.49 (m, 1H), 1.37 (t, 3H, $J=6.8\text{Hz}$); MS m/z 628.5 ($M + 1$)


275



^1H NMR 400 MHz (CD_3OD) δ 8.17 (d, 1H, $J=8.0\text{ Hz}$), 7.93 (s, 1H), 7.74 (dd, 1H, $J=1.6, 8.0\text{Hz}$), 7.47 (t, 1H, $J=8.0\text{ Hz}$), 7.25 (m, 1H), 7.13 (d, 1H, $J=8.8\text{Hz}$), 6.48 (d, 1H, $J=2.4\text{ Hz}$), 6.37 (dd, 1H, $J=2.4, 8.8\text{Hz}$), 3.66 (m, 4H), 3.34 (m, 2H), 3.14 (m, 4H), 2.81 (m, 2H), 2.65 (m, 2H), 2.98 (d, 2H, $J=12\text{Hz}$), 1.68 (m, 7H), 1.30 (m, 1H), 1.02 (d, 6H, $J=6.8\text{ Hz}$), 0.99 (m, 1H), 0.32 (m, 2H), 0.04 (m, 2H); MS m/z 639.5 ($M + 1$)


- 211 -

276




^1H NMR 400 MHz (CD_3OD) δ 8.32 (d, 1H, $J=8.0$ Hz), 8.11 (s, 1H), 7.92 (dd, 1H, $J=1.6, 8.0\text{Hz}$), 7.65 (t, 1H, $J=7.6\text{Hz}$), 7.43 (t, 1H, $J=7.6\text{Hz}$), 7.28 (d, 1H, $J=8.4\text{Hz}$), 6.67 (d, 1H, $J=2.4\text{Hz}$), 6.56 (dd, 1H, $J=2.4, 8.8\text{Hz}$), 4.03 (m, 2H), 3.86 (m, 4H), 3.73 (m, 2H), 3.36 (m, 2H), 3.35 (m, 2H), 3.23 (m, 2H), 2.85 (m, 2H), 2.23 (m, 2H), 1.83 (m, 2H), 1.19 (d, 6H), 1.13 (m, 1H), 0.48 (m, 2H), 0.21 (m, 2H); MS m/z 641.5 ($M + 1$)

277




^1H NMR 400 MHz (CD_3OD) δ 8.06 (d, 1H, $J=7.6\text{Hz}$), 7.91 (s, 1H), 7.72 (dd, 1H, $J=1.6, 8.0\text{Hz}$), 7.39 (t, 1H, $J=7.6$ Hz), 7.20 (m, 1H), 7.10 (d, 1H, $J=4.8$ Hz), 6.48 (d, 1H, $J=2.4\text{Hz}$), 6.36 (dd, 1H, $J=2.4, 8.8\text{Hz}$), 3.85 (m, 2H), 3.66 (m, 4H), 3.56 (m, 2H), 3.18 (m, 2H), 3.16 (m, 1H), 2.98 (m, 2H), 2.65 (t, 2H, $J=12\text{Hz}$), 2.29 (s, 3H), 2.05 (m, 2H), 1.64 (m, 2H), 0.95 (m, 1H), 0.32 (m, 2H), 0.03 (m, 2H); MS m/z 628.5 ($M + 1$)

278




^1H NMR 400 MHz (CD_3OD) δ 8.18 (d, 1H, $J=8.0\text{Hz}$), 8.15 (s, 1H), 7.89 (dd, 1H, $J=1.6, 8.0\text{Hz}$), 7.56 (t, 1H, $J=7.2\text{Hz}$), 7.38 (m, 1H), 7.25 (d, 1H, $J=8.8$ Hz), 6.65 (d, 1H, $J=2.4$ Hz), 6.53 (dd, 1H, $J=2.4, 8.8$ Hz), 3.82 (m, 4H), 3.49 (m, 2H), 3.31 (m, 1H), 2.95 (m, 2H), 2.83 (m, 2H), 2.47 (s, 3H), 2.17 (m, 2H), 1.86 (m, 7H), 1.52 (m, 1H), 1.15 (m, 1H), 0.49 (m, 2H), 0.21 (m, 2H); MS m/z 670.4 ($m + 1$)

279


MS m/z .641.5 ($M + 1$)

280




^1H NMR 400 MHz (CD_3OD) δ 8.13 (s, 1H), 8.10 (d, 1H, $J=8.0\text{Hz}$), 7.85 (t, 1H, $J=1.6\text{Hz}$), 7.83 (t, 1H, $J=1.6\text{Hz}$), 7.57 (m, 1H), 7.32 (m, 1H), 7.02 (d, 1H, $J=1.2\text{Hz}$), 6.81 (dd, 1H, $J=1.6, 8.4\text{Hz}$), 4.05 (q, 2H, $J=6.8\text{Hz}$), 3.36 (m, 4H), 3.08 (m, 2H), 2.84 (s, 3H), 2.40 (s, 3H), 1.33 (t, 3H, $J=6.8\text{Hz}$); MS m/z .560.4 ($M + 1$)

281

AMS m/z .600.5 ($M + 1$)


282




NMR 400 MHz (CD_3OD) δ 8.30 (d, 1H, $J=8.0\text{Hz}$), 8.21 (s, 1H), 8.00 (d, 1H, $J=8.4\text{Hz}$), 7.91 (dd, 1H, $J=1.63, 8.0\text{Hz}$), 7.71 (m, 1H), 7.42 (m, 1H), 7.08 (d, 1H, $J=2.0\text{Hz}$), 6.91 (dd, 1H, $J=1.6, 8.4\text{Hz}$), 4.30 (m, 2H), 4.13 (q, 2H, $J=7.2\text{Hz}$), 3.60 (m, 4H), 3.16 (m, 2H), 1.41 (t, 3H, $J=7.2\text{Hz}$), 1.90 (qd, 6H, $J=6.8\text{Hz}$); MS m/z .573.40 ($M + 1$)

- 213 -


283

MS m/z 681.5 ($M + 1$)


284

MS m/z 559.4 ($M + 1$)

285

MS m/z 577.3 ($M + 1$)


286




NMR 400 MHz (CD_3OD) δ 8.28 (s, br, 1H), 8.11 (s, br, 1H), 7.89 (d, 1H, $J=8.4$ Hz), 7.53 (s, br, 1H), 7.35 (t, 1H, $J=6.0$ Hz), 7.29 (m, 1H), 6.30 (s, 1H), 6.20 (d, 1H, $J=7.8$ Hz), 4.07 (q, 2H, $J=7.2$ Hz), 4.02 (m, 1H), 3.68 (m, 1H), 3.60 (m, 1H), 3.54 (m, 1H), 3.34 (m, 1H), 2.95 (s, 6H), 2.57 (m, 1H), 2.51 (s, 3H), 2.25 (m, 1H), 1.30 (t, 3H, $J=7.2$ Hz); MS m/z 590.3 ($M + 1$)

- 214 -


287

MS m/z .533.3 ($M + 1$)


288

MS m/z .565.3 ($M + 1$)

289


MS m/z .546.4 ($M + 1$)

290




NMR 400 MHz (CD_3OD) δ 8.29 (d, 1H, $J=8.4$ Hz), 8.24 (s, 1H), 7.91 (dd, 1H, $J=1.2, 7.8$ Hz), 7.57 (m, 2H), 7.35 (t, 1H, $J=7.2$ Hz), 6.98 (d, 1H, $J=9.0$ Hz), 6.77 (dd, 1H, $J=3.0, 9.0$ Hz), 4.18 (t, 2H, $J=4.8$ Hz), 3.82 (s, 3H), 3.56 (m, 2H), 3.17 (t, 2H, $J=4.8$ Hz), 3.3 (s, 1H), 3.01 (m, 2H), 2.53 (s, 3H), 1.92 (m, 2H), 1.79 (m, 3H), 1.53 (m, 1H); MS m/z .591.3 ($M + 1$)

291


MS m/z 521.4 ($M + 1$)

292




NMR 400 MHz (CD_3OD) δ 8.28 (d, 1H, $J=8.4$ Hz), 8.15 (s, 1H), 7.89 (dd, 1H, $J=2.4, 7.8$ Hz), 7.55 (t, 1H, $J=7.2$ Hz), 7.44 (d, 1H, $J=3.0$ Hz), 7.33 (t, 1H, $J=7.2$ Hz), 6.99 (dd, 1H, $J=3.6, 9.0$ Hz), 6.80 (dd, 1H, $J=3.0, 9.0$ Hz), 4.18 (t, 2H, $J=4.8$ Hz), 3.80 (s, 3H), 3.51 (t, 2H, $J=4.8$ Hz), 3.29 (m, 4H), 2.50 (s, 3H), 1.30 (m, 6H); MS m/z 535.4 ($M + 1$)

293




NMR 400 MHz (CD_3OD) δ 8.31 (d, 1H, $J=8.4$ Hz), 8.14 (s, 1H), 7.90 (dd, 1H, $J=1.2, 7.8$ Hz), 7.59 (m, 1H), 7.53 (d, 1H, $J=3.0$ Hz), 7.33 (t, 1H, $J=7.8$ Hz), 6.96 (d, 1H, $J=9.0$ Hz), 6.76 (dd, 1H, $J=3.0, 8.4$ Hz), 4.18 (t, 2H, $J=4.2$ Hz), 3.80 (s, 3H), 3.55 (m, 2H), 3.46 (t, 2H, $J=4.8$ Hz), 3.00 (t, 2H, $J=12.0$ Hz), 2.50 (s, 3H), 1.89 (m, 2H), 1.77 (m, 3H), 1.52 (m, 1H); MS m/z 547.4 ($M + 1$)

294

MS m/z 605.3 ($M + 1$)


- 216 -

295




NMR 400 MHz (CD₃OD) δ 8.33 (d, 1H, J=8.4 Hz), 8.14 (s, 1H), 7.90 (dd, 1H, J=1.2, 7.8 Hz), 7.57 (m, 1H), 7.53 (d, 1H, J=3.0 Hz), 7.33 (t, 1H, J=7.2 Hz), 6.95 (d, 1H, J=9.0 Hz), 6.74 (dd, 1H, J=3.0, 9.0 Hz), 4.17 (t, 1H, J=4.8 Hz), 3.80 (s, 3H), 3.51 (m, 4H), 3.27 (m, 2H), 2.50 (s, 3H), 1.85 (m, 4H), 1.70 (m, 4H); m/z 561.4 (M + 1)

296




MS m/z 587.4 (M + 1)

297







MS m/z 589.4 (M + 1)

298




MS m/z 590.4 (M + 1)

- 217 -




- 218 -


303

MS *m/z*.574.4 (*M* + 1)


304

MS *m/z*.618.2 (*M* + 1)


305

MS *m/z*.588.2 (*M* + 1)


306

MS *m/z*.562.4 (*M* + 1)


307

MS m/z .530.4 (M + 1)


308

MS m/z .562.4 (M + 1)


309

MS m/z .576.2 (M + 1)


310

MS m/z .546.5 (M + 1)


311

MS m/z 558.2 ($M + 1$)


312

MS m/z 604.5 ($M + 1$)

313


MS m/z 616.5 ($M + 1$)

314




NMR 400 MHz (CD_3OD) δ 8.17 (d, 1H, $J=8.0$ Hz), 7.88 (m, 2H), 7.56 (t, 1H, $J=8.0$ Hz), 7.37 (m, 1H), 7.17 (d, 1H, $J=8.8$ Hz), 6.61 (d, 1H, $J=2.4$ Hz), 6.47 (9dd, 1H, $J=2.4, 8.4$ Hz), 3.75 (m, 7H), 3.12 (m, 4H), 2.42 (s, 3H); MS m/z 489.1 ($M + 1$)

315


MS m/z : 546.4 ($M + 1$)

316




NMR 400 MHz (CD_3OD) δ 8.22 (d, 1H, $J=8.4$ Hz), 7.90 (d, 1H, $J=4.4$ Hz), 7.86 (dd, 1H, $J=1.6, 8.0$ Hz), 7.55 (m, 1H), 7.34 (m, 1H), 7.27 (d, 1H, $J=8.8$ Hz), 6.59 (d, 1H, $J=2.4$ Hz), 6.45 (dd, 1H, $J=1.6, 8.8$ Hz), 4.00 (q, 2H, $J=6.8$ Hz), 3.75 (t, 4H, $J=4.8$ Hz), 3.10 (m, 4H), 2.42 (s, 3H), 1.94 (m, 6H), 1.25 (t, 3H, $J=6.8$ Hz); MS m/z : 503.2 ($M + 1$)






317






MS m/z : 575.4 ($M + 1$)






318




NMR 400 MHz (CD_3OD) δ 8.24 (d, 1H, $J=7.6$ Hz), 8.17 (s, 1H), 7.94 (dd, 1H, $J=1.6, 8.0$ Hz), 7.60 (t, 1H, $J=7.6$ Hz), 7.44 (m, 1H), 7.39 (t, 1H, $J=7.23$ Hz), 6.68 (s, 1H), 6.50 (m, 1H), 4.10 (q, 2H, $J=7.2$ Hz), 3.84 (m, 4H), 3.16 (m, 4H), 2.83 (t, 2H, $J=7.2$ Hz), 1.37 (m, 6H), 0.75 (t, 3H, $J=7.6$ Hz); MS m/z : 591.1 ($M + 1$)









334

MS m/z 520.0 ($M + 1$)


335

MS m/z 478.0 ($M + 1$)


336

MS m/z 543.2 ($M + 1$)

337


MS m/z 563.2 ($M + 1$)

338




NMR 400 MHz (CD_3OD) δ 8.18 (d, 1H, $J=7.6$ Hz), 7.98 (d, 1H, $J=4.4$ Hz), 7.86 (dd, 1H, $J=1.6, 8.0$ Hz), 7.56 (m, 1H), 7.46 (s, 1H), 7.30 (m, 1H), 6.83 (m, 2H), 3.75 (s, 3H), 2.40 (s, 3H), 2.08 (s, 3H); MS m/z 418.1 ($M + 1$)

339




NMR 400 MHz (CD₃OD) δ 8.17 (d, 1H, J=8.0 Hz), 8.00 (d, 1H, J=4.4 Hz), 7.86 (dd, 1H, J=1.6, 8.0Hz), 7.53 (m, 1H), 7.43 (s, 1H), 7.32 (m, 1H), 6.79 (s, 1H), 4.00 (q, 2H, J=6.8Hz), 3.83 (m, 4H), 3.76 (m, 2H), 2.44 (s, 3H), 1.29 (t, 3H, J=7.2 Hz), 1.24 (t, 3H, J=7.2 Hz); MS *m/z* 547.2 (M + 1)

340




MS *m/z* 588.4 (M + 1)

341




MS *m/z* 600.4 (M + 1)

342




MS *m/z* 602.4 (M + 1)

343




MS m/z 593.00/ 595.00 (M + 1)

344




MS m/z 621.20 / 623.20 (M + 1)

345




¹H NMR 400 MHz (DMSO-d₆) δ 8.95 (s, br, 1H), 8.49 (s, br, 2H), 7.80 (dd, J = 7.54, 2.13 Hz, 1H), 7.34 (t, J = 8.36, 2H), 7.14 (m, 3 H), 6.93 (m, 3 H); MS m/z 540.10 / 542.10 (M + 1)

346




MS m/z 506.10 / 508.10 (M + 1)

347




MS m/z 522.10 / 524.10 (M + 1)

348




MS m/z 635.20 / 637.20 (M + 1)

349




MS m/z 456.10 / 458.10 (M + 1)

350




MS m/z 472.10 / 474.10 (M + 1)

351




MS m/z 585.20 / 587.20 (M + 1)

352




MS m/z 571.20 / 573.10 (M + 1)

353




MS m/z 448.10 / 450.10 (M + 1)

354




MS m/z 464.10 / 466.10 (M + 1)

355




¹H NMR 400 MHz (DMSO-*d*₆) δ 9.54 (s, br, 1H), 8.76 (s, br, 1H), 8.31 (s, 1), 8.28 (d, J = 8.14 Hz, 1H), 7.95 (t, J = 5.64 Hz, 7.85 (dd, J = 7.95, 1.48 Hz, 1H), 7.51 (t, J = 7.53 Hz, 1H), 7.33 (m, 2H), 6.73 (s, 1H), 4.05 (q, J = 6.92, 2H), 3.98 (m, 4H), 3.71 (m, 2H), 3.12 (m, 4H), 2.82 (p, J = 5.77
MS m/z 563.20 / 565.20 (M + 1)

356




357




MS m/z 412.10 / 414.10 (M + 1)

358









MS m/z 428.10 / 430.10 (M + 1)


359



MS m/z 541.20 / 543.20 (M + 1)




366




MS m/z 649.20 / 651.20 (M + 1)

367




MS m/z 591.20 / 593.20 (M + 1)

368




MS m/z 605.20 / 607.20 (M + 1)

369



MS m/z 448.10 / 450.10 (M + 1)


370




¹H NMR 600 MHz (DMSO-*d*₆) δ 9.46 (s, 1H), 8.46 (s, 1H), 8.37 (d, J = 8.12 Hz, 1H), 8.25 s, (1H), 7.75 (dd, 7.87, 1.40 Hz, 1H), 7.55 (t, J = 7.50 Hz, 1H), 7.37 (s, 1H), 7.30 (dt, J = 8.05, 0.8 Hz, 1H), 6.91 (d, J = 8.92 Hz, 1H), 6.58 (dd, J = 8.88, 3.04 Hz, 1H), 3.70 (s, 3H), 3.55 (s, 3H), 2.58 (s, 3H).

MS m/z 577.20 / 579.20 (M + 1)

371




372




MS m/z 563.20 / 565.20 (M + 1)

373




MS m/z 519.20 / 521.20 (M + 1)

374




MS m/z 533.20 / 535.20 (M + 1)

375




MS m/z 476.20 / 478.20 (M + 1)

376




¹H NMR 600 MHz (DMSO-*d*₆) δ 9.39 (s, 1H), 8.37 (s, 1H), 8.27 (s, 1H), 8.25 (s, 1H), 7.88 (t, *J* = 6.02 Hz, 1H), 7.76 (dd, *J* = 7.94, 1.35 Hz, 1H), 7.50 (dt, *J* = 8.24, 1.89 Hz, 1H), 7.40 (s, 1H), 7.25 (t, *J* = 7.39 Hz, 1H), 6.90 (d, *J* = 8.92 Hz, 1H) 5.65 (dd, *J* = 8.87, 3.03 Hz, 1H).. 3.70 (s, 3H), 3.54 (s, 3H)
MS m/z 605.30 / 607.20 (M + 1)

377




378




MS m/z 591.20 / 593.20 (M + 1)

379




MS m/z 547.20 / 549.20 (M + 1)

380




MS m/z 561.20 / 563.20 (M + 1)

381




MS m/z 548.20 / 550.20 / 552.20 (M + 1)

382




MS m/z 548.20 / 550.20 / 552.20 (M + 1)

383




MS m/z 563.30 / 565.30 (M + 1)

384




MS m/z 577.30 / 579.30 (M + 1)

385




MS m/z 591.40 / 593.30 (M + 1)

386




MS m/z 605.40 / 607.40 (M + 1)

387




MS m/z 617.40 / 619.40 (M + 1)

388




MS m/z 619.40 / 621.40 (M + 1)

389




MS m/z 576.30 / 578.30 (M + 1)

390




MS m/z 563.30 / 565.30 (M + 1)

391




MS m/z 560.40 / 562.40 (M + 1)

392




MS m/z 574.40 / 576.50 (M + 1)

393




¹H NMR 400 MHz (DMSO-*d*₆) δ 9.47 (s, 1H), 8.45 (d, J = 7.37 Hz, 1H), 8.36 (s, 1H), 8.20 (s, 1H), 7.83 (d, J = 7.93 Hz, 1H), 7.59 (t, J = 8.07 Hz, 1H), 7.47 (d, J = 8.69 Hz, 1H), 7.32 (t, J = 7.54 Hz, 1H), 6.63 (dd, J = 12.67, 2.42 Hz, 1H), 6.49 (m, 1H), 5.16 (m, 1H), 4.01-3.68 (m, 3H), 3.48-
MS m/z 587.40 / 589.40 (M + 1)

394




395




MS m/z 573.40 / 575.40 (M + 1)

396




MS m/z 587.40 / 589.40 (M + 1)

397




MS m/z 586.40 / 588.40 (M + 1)

398




MS m/z 586.40 / 588.40 (M + 1)

399




MS m/z 560.40 / 562.40 (M + 1)

400




¹H NMR 400 MHz (DMSO-*d*₆) δ 10.30 (s, br, 1H), 9.53 (s, 1H), 8.53 (s, 1H), 8.41 (d, J = 7.90 Hz, 1H), 8.26 (s, 1H), 7.90 (dd, J = 7.91, 1.25 Hz, 1H), 7.64 (t, J = 7.93 Hz, 1H), 7.46 (d, J = 8.70 Hz, 1H), 7.38 (t, J = 7.48 Hz, 1H), 6.67 (dd, J = 13.68, 2.49 Hz, 1H), 6.46 (m, 1H), 5.19 (m, 1H), 4.04 (m, 1H).
MS m/z 573.20 / 575.20 (M + 1)

401




402




MS m/z 573.20 / 575.20 (M + 1)

403




MS m/z 587.40 / 589.40 (M + 1)

404




MS m/z 573.40 / 575.40 (M + 1)

405




MS m/z 587.40 / 589.40 (M + 1)

406




MS m/z 586.40 / 588.40 (M + 1)

407




MS m/z 586.40 / 588.40 (M + 1)

408




^1H NMR 600 MHz (CD_3OD) δ 8.18 (d, J = 7.80 Hz, 1H), 8.03 (s, 1H), 7.87 (dd, J = 7.96, 1.21 Hz, 1H), 7.47 (m, 2H), 7.25 (t, J = 7.82 Hz, 1H), 6.59 (s, 1H), 6.40 (d, J = 8.56 Hz, 1H), 5.13 (s, 1H), 3.78 (m, 4H), 3.30 (m, 2H), 3.28 (s, 3H), 2.92 (m, 2H), 2.56 (m, 1H), 2.30 (m, 2H), 0.89 (d, J = MS m/z 547.40 / 549.40 ($M + 1$))

409




410




MS m/z 547.40 / 549.40 ($M + 1$)

411




MS m/z 587.50 / 589.40 ($M + 1$)

412




MS m/z 588.50 / 590.40 ($M + 1$)

413




MS m/z 637.50 / 639.50 ($M + 1$)

414




MS m/z 638.50 / 640.50 (M + 1)

415




MS m/z 602.50 / 604.50 (M + 1)

416




MS m/z 583.40 / 585.40 (M + 1)

417




MS m/z 584.40 / 586.40 (M + 1)

418




MS m/z 548.40 / 550.40 (M + 1)

419




MS m/z 540.40 / 542.40 (M + 1)

420




MS m/z 541.40 / 543.40 (M + 1)

421




MS m/z 505.40 / 507.40 (M + 1)

422




MS m/z 667.30 / 669.30 (M + 1)

423




MS m/z 632.30 / 634.30 (M + 1)

424




MS m/z 573.40 / 575.40 (M + 1)

425




^1H NMR 400 MHz (CD_3OD) δ 8.34 (d, $J = 8.4$ Hz, 1H), 8.13 (s, 1H), 7.96 (dd, $J = 1.2, 7.8$ Hz, 1H), 7.63 (t, $J = 7.8$ Hz, 1H), 7.43 (m, 2H), 6.72 (d, $J = 2.4$ Hz, 1H), 6.57 (m, 1H), 4.07 (m, 1H), 3.40 (m, 1H), 3.02 (t, $J=10.6$ Hz, 1H), 2.80 (t, $J=11.8$, 1H), 2.65 (m, 4H), 2.54, (s, 3H), 2.26 (s, 6H), 1.56 (m, 4H), 1.36 (d, $J = 7.2$ Hz, 6H). MS m/z 574.20 ($M + 1$).

426




MS m/z 603.20 ($M + 1$).

427




MS m/z 504.20 ($M + 1$).

428


MS m/z 585.30 ($M + 1$).

429


MS m/z 627.20 ($M + 1$).

- 243 -


430

MS m/z 616.20 ($M + 1$).


431

MS m/z 602.20 ($M + 1$).

432


MS m/z 562.10 ($M + 1$).

433




MS m/z 408.1 ($M + 1$).

434




MS m/z 421.1 ($M + 1$).

435




MS m/z 421.1 ($M + 1$).

436




MS m/z 480.10 ($M + 1$).


437

MS m/z 394.10 ($M + 1$).

438


MS m/z 408.10 ($M + 1$).

439




^1H NMR 400 MHz (CD_3OD) δ 8.34 (d, $J = 8.4$ Hz, 1H), 8.13 (s, 1H), 7.96 (dd, $J = 1.3, 7.9$ Hz, 1H), 7.64 (t, $J = 7.9$ Hz, 1H), 7.43 (m, 1H), 7.13 (d, $J = 2.8$ Hz, 1H), 6.24 (d, $J = 2.8$ Hz, 1H), 4.08 (m, 1H), 3.41 (m, 1H), 2.58 (s, 6H), 2.54 (s, 3H). MS m/z 574.20 ($M + 1$).


440

MS m/z 468.20 ($M + 1$).


441

MS m/z 552.30 ($M + 1$).

442


MS m/z 592.30 ($M + 1$)

443


MS m/z 504.20 ($M + 1$)

- 247 -


444

MS m/z 588.30 (M + 1)


445

MS m/z 628.30 (M + 1)


446

MS m/z 617.20 (M + 1)


447

MS m/z 503.20 ($M + 1$)


448

MS m/z 587.20 ($M + 1$)


449

MS m/z 627.30 ($M + 1$)


450

MS m/z 616.30 ($M + 1$)


451

MS m/z 549.30 ($M + 1$)


452

MS m/z 509.30 ($M + 1$)


453

MS m/z 495.20 ($M + 1$)


454

MS m/z 495.20 ($M + 1$)


455




458

MS *m/z* 531.20 (M + 1)


459

MS *m/z* 584.30 (M + 1)


460

MS *m/z* 544.20 (M + 1)


461

MS m/z 530.20 ($M + 1$)


462

MS m/z 530.20 ($M + 1$)

463

MS m/z 601.30 ($M + 1$)

464

MS m/z 626.30 ($M + 1$)

Example 54: Cell-free ZAP-70 Kinase assay

The ZAP-70 kinase assay is based on time-resolved fluorescence resonance energy transfer (FRET). 80 nM ZAP-70 are incubated with 80 nM Lck (lymphoid T-cell protein tyrosine kinase) and 4 μ M ATP in ZAP-70 kinase buffer (20 mM Tris, pH 7.5, 10 μ M Na_3VO_4 , 1 mM DTT, 1 mM MnCl_2 , 0.01 % BSA, 0.05 % Tween-20) for 1 hour at room temperature in a siliconized polypropylene tube. Then, the selective Lck inhibitor PP2 (1-tert-butyl-3-(4-chloro-phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-ylamine; Alexis Biochemicals) is added (final concentration 1.2 μ M) and incubated for further 10 min. 10 μ L of this solution is mixed with the 10 μ L biotinylated peptide LAT-11 (1 μ M) as substrate and 20 μ L of serial dilutions of inhibitors and incubated for 4 hours at room temperature. The kinase reaction is terminated with 10 μ L of a 10 mM EDTA solution in detection buffer (20 mM Tris, pH 7.5, 0.01 % BSA, 0.05 % Tween-20). 50 μ L europium-labelled anti-phosphotyrosine antibody (Eu-PT66; final concentration 0.125 nM); and 50 μ L streptavidin-allophycocyanine (SA-APC; final concentration 40 nM) in detection buffer are added. After 1 hour incubation at room temperature fluorescence is measured on the Victor2 Multilabel Counter (Wallac) at 665 nm. Background values (low control) are obtained in the absence of test samples and ATP and are subtracted from all values. Signals obtained in the absence of test samples are taken as 100% (high control). The inhibition obtained in the presence of test compounds is calculated as percent inhibition of the high control. The concentration of test compounds resulting in 50% inhibition (IC_{50}) is determined from the dose-response curves. In this assay, the agents of the invention have IC_{50} values in the range of 10 nM to 2 μ M, preferably from 10 nM to 100 nM.

Recombinant ZAP-70 kinase is obtained as follows: A nucleic acid encoding full-length human ZAP-70 (GenBank #L05148) is amplified from a Jurkat cDNA library by RT-PCR and cloned into the pBluescript KS vector (Stratagene, California, USA). The authenticity of the ZAP-70 cDNA insert is validated by complete sequence analysis. This donor plasmid is then used to construct a recombinant baculovirus transfer vector based on the plasmid pVL1392 (Pharmingen, California, USA) featuring in addition an N-terminal hexahistidine tag. Following co-transfection with AcNPV viral DNA, 10 independent viral isolates are derived via plaque-purification, amplified on small scale and subsequently analyzed for recombinant ZAP-70 expression by Western Blot using a commercially available anti-ZAP-70 antibody (Clone 2F3.1, Upstate Biotechnology, Lake Placid, NY, USA). Upon further amplification of one positive recombinant plaque, titrated virus stocks are prepared and used for infection of Sf9 cells grown in serum-free SF900 II medium (Life Technologies, Basel, Switzerland) under defined, optimized conditions. ZAP-70 protein is isolated from the lysate of infected Sf9 cells by affinity chromatography on a Ni-NTAcolumn (Qiagen, Basel, Switzerland).

Recombinant His-tagged ZAP-70 is also available from PanVera LLC, Madison, Wisconsin, USA.

LAT-11 (linker for activation of T cell): The biotinylated peptide LAT-11 (Biotin-EEGAPDYENLQELN) used as a substrate in the ZAP-70 kinase assay is prepared in analogy to known methods of peptide synthesis. The N- α Fmoc group of Fmoc-Asn(Trt)-oxymethyl-4-phenoxymethyl-co(polystyrene-1%-divinyl-benzene), content of Asn approx. 0.5 mmol/g, is cleaved using piperidine, 20% in DMF. Four equivalents per amino-group of Fmoc-amino acid protected in their side chains [Asp(OtBu), Glu(OtBu), Asn(Trt), Gln(Trt) and Tyr(tBu)] are coupled using DIPCDI and HOBt in DMF. After complete assembly of the peptide chain the terminal Fmoc-protecting group is removed with piperidine in DMF as before. L(+)-biotinyl-amino-hexanoic acid is then coupled to the terminal amino group using DIPCDI and HOBt in DMF using four equivalents of the reagents for four days at RT. The peptide is cleaved from the resin support and all side-chain protecting groups are simultaneously removed by using a reagent consisting of 5% dodecylmethylsulfide and 5% water in TFA for two hours at RT. Resin particles are filtered off, washed with TFA and the product is precipitated from the combined filtrates by the addition of 10 to 20 volumes of diethyl ether, washed with ether and dried. The product is purified by chromatography on a C-18 wide-pore silica column using a gradient of acetonitrile in 2% aqueous phosphoric acid. Fractions containing the pure compound are

collected, filtered through an anion-exchange resin (Biorad, AG4-X4 acetate form) and lyophilized to give the title compound. MS: 1958.0 (M-H)⁻¹

Example 56: Anchorage-independent tumor cell growth assay

Mouse mammary carcinoma 4T1 cells (5×10^3) are plated in 96-well Ultra low Attachment plates (#3474, Corning Inc.) in 100 μ L of Dulbecco's modified eagle medium containing 10% FBS. Cells are cultured for 2 h and inhibitors are added at various concentrations in a final concentration of 0.1% DMSO. After 48 h, cell growth is assayed with the cell counting kit-8 (Wako Pure Chemical), which uses a water soluble tetrazolium salt WST8. Twenty μ L of the reagent is added into each well and cells are further cultured for 2 h. The optical density is measured at 450 nm. The concentration of compound causing 50 % inhibition of growth is determined.

Example 59 *In vivo* activity in the nude mouse xenograft model:

female or male BALB/c nude mice (5–8 weeks old, Charles River Japan, Inc., Yokohama, Japan) are kept under sterile conditions with water and feed *ad libitum*. Tumours are induced by subcutaneous injection of tumour cells (human epithelial cell line MIA PaCa-2; European Collection of Cell Cultures (ECACC), Salisbury, Wiltshire, UK, Catalogue Number 85062806; cell line from a 65 year old Caucasian male; undifferentiated human pancreatic carcinoma cell line) into left or right flank of mice under Forene[®] anaesthesia (Abbott Japan Co., Ltd., Tokyo, Japan). Treatment with the test compound is started when the mean tumor volumes reached approximately 100 mm³. Tumour growth is measured two times per week and 1 day after the last treatment by determining the length of two perpendicular axis. The tumour volumes are calculated in accordance with published methods (see Evans et al., Brit. J. Cancer 45, 466-8, 1982). The anti-tumour efficacy is determined as the mean increase in tumour volume of the treated animals divided by the mean increase in tumour volume of the untreated animals (controls) and, after multiplication by 100, is expressed as delta T/C [%]. Tumour regression is reported as the mean changes of tumor volume of the treated animals divided by the mean tumor volume at start of treatment and, after multiplication by 100, is expressed as regression [%]. The test compound is orally administered daily with or without drug holidays.

As an alternative to cell line MIA PaCa-2, another cell line may also be used in the same manner, for example:

- the 4T1 breast carcinoma cell line (ATCC Number CRL-2539; see also Cancer. 88(12 Supple), 2979-2988, 2000) with female BALB/c mice (injection into mammary fat pad).

On the basis of these studies, a compound of formula I according to the invention shows therapeutic efficacy especially against proliferative diseases responsive to an inhibition of a tyrosine kinase.

Example 60: Tablets

Tablets comprising 50 mg of active ingredient, for example one of the compounds of formula I described in Examples 1 to 131, and having the following composition are prepared in customary manner:

Composition:

active ingredient	50 mg
wheat starch	150 mg
lactose	125 mg
colloidal silicic acid	12.5 mg
talc	22.5 mg
magnesium stearate	2.5 mg
Total:	362.5 mg

Preparation: The active ingredient is mixed with a portion of the wheat starch, with the lactose and the colloidal silicic acid and the mixture is forced through a sieve. A further portion of the wheat starch is made into a paste, on a water bath, with five times the amount of water and the powder mixture is kneaded with the paste until a slightly plastic mass is obtained.

The plastic mass is pressed through a sieve of about 3 mm mesh size and dried, and the resulting dry granules are again forced through a sieve. Then the remainder of the wheat starch, the talc and the magnesium stearate are mixed in and the mixture is compressed to form tablets weighing 145 mg and having a breaking notch.

Example 61: Soft Capsules

5000 soft gelatin capsules comprising each 50 mg of active ingredient, for example one of the compounds of formula I described in Examples 1 to 131, are prepared in customary manner:

Composition:

active ingredient	250 g
Lauroglykol	2 litres

Preparation: The pulverized active ingredient is suspended in Lauroglykol® (propylene glycol laurate, Gattefossé S.A., Saint Priest, France) and ground in a wet pulverizer to a particle size of approx. 1 to 3 μm . 0.419 g portions of the mixture are then dispensed into soft gelatin capsules using a capsule-filling machine.

Biological results:

Examp e	FA K IC50 (nM)	Pho s IC50 (μM)	Growt h IC50 (μM)	T Cell Migration IC50 (μM)	IGF- 1R IC50 (μM)
1.00	140	0.7	>10		
2.00	13	1.2			
3.01	44	0.34	>10		
3.02	36	0.85	4		
3.03	9.1	0.14	0.8		
3.04	32	0.53	2		
3.05	21	0.17	2		>10
3.06	13	0.11	2		
3.07	16	0.45	2		
3.08	74	0.3	6		
3.09	48	0.5	0.7		
3.10	52	0.95	>10		
3.11	9	0.04	0.3		0.2
3.12	5.4	0.01	1		
3.13	58	1.7	0.6		0.74
3.14	54	0.4	5		
3.15	7	0.02	0.8		0.94
3.16	48	1.1	3		
3.17	2.8	0.03	0.2		<0.0
3.18	130	1.5	9		
3.19	6.8	0.35	0.8		0.1
3.20	16	0.22	0.3		
3.22	120	0.9	2		
3.23	38	0.39	0.5		
3.24	64	3.5	5		
3.25	22	0.3	0.3		0.81
3.26	50	0.79	2		
3.28	43	0.71	0.7		
3.29	89	0.6	>10		
3.30	69	0.6	3		
3.31	13	1.1	5		
3.32	14	0.18	0.49	8	0.12
3.33	2.9	0.03	0.05	9	0.13
3.34	7	0.1	0.24	3	<0.0
3.35	13	0.02	0.17		8
3.36	43	1.8	2.8		3.55
3.37	39	1.1	2.6		
3.38	64	1.7	3.8		
3.39	2	0.02	0.03	1	0.09
3.40	9	>10	0.9		
3.41	22	>10	0.43		
3.42	29	0.35	0.3		
3.43	5.6	0.2	0.11		0.27
3.44	11	0.05	0.09		0.09
3.45	0.9	0.02	0.02		
3.46	4	0.1	0.18	0.3	

- 260 -

3.47	1	0.1	0.06		
3.48	7	0.07	0.3		0.21
3.49	39	10	0.39		
3.50	13	0.12	1		1.19
3.51	29	0.2	0.4		0.41
3.52	29	0.42	2		
3.53	6	0.07	0.21		
3.54	0.9	0.01	0.07		<0.0
3.55	34	>10	3		8
3.56	28	0.53	0.15		
3.57	28	0.61	3		
3.58	21	0.08	0.3		0.14
3.59	95	1.2	>10		
3.60	90	0.93	2		
3.61	12	10	>10		
3.62	63	>10	>10		
3.63	27	>10	>10		
3.64	5	0.13	0.7	1	0.2
3.65	8	0.08	0.1		0.15
3.66	1	0.08	0.07		0.25
3.67	6	0.38	0.39		
3.68	5.5	0.2	0.63		1
3.69	4	0.2	0.11	8	0.5
3.70	3.5	0.02	0.13		
3.71	11	0.05	0.08		
3.72	2.1	0.11	0.06		
3.73	11	0.03	0.29		1.63
3.74	15	0.1	0.15		
3.75	72	0.5	1.3		
3.76	15	0.29	1.3		0.7
3.77	65	>10	3		
3.78	10	>10	0.22		
3.79	5	1.3	0.12		
3.80	12	0.22	0.45		5
3.81	21	0.52	0.98		>10
3.82	4.8	0.2	0.07		
3.83	20	0.08	0.32		0.68
3.84	10	1	0.08		
6.00	110	0.35	5		
7.00	5.3	0.21	0.47		0.0
7.01	4.7	0.6	0.54	4	0.19
7.02	7.5	0.1	0.36		0.19
7.03	2.9	0.3	0.39		0.77
7.04	5.2	1	0.29		0.27
7.05	6.2	0.3	0.2		0.25
7.06	17	0.8	1.09		0.2
7.07	4.1	0.9	0.18	5	
7.08	8.7	0.8	1		
7.09	8.2	1	0.85		
7.10	6.6	1	0.98		
7.11	2.5	0.6	1.2		0.77
7.12	1.9	0.9	1		0.62

- 261 -

				1	
7.13	5.5	0.8	1.22		
7.14	7.6	0.3	0.36		0.33
7.15	4.5	0.06	0.19		0.26
7.16	6.4	0.2	0.42		
7.17	4.3	0.7	0.69		
7.18	6.2	0.5	0.7		
7.19	13		0.33		
7.20	2.5	>10	0.11		
7.21	3.3	>10	0.46		
7.22	25		0.48		
7.23	1.4		0.25		
7.24	5.1		0.09		
7.25	13	0.2	0.73		
7.25	2	>10	0.57		
7.26	4.1		0.15		
7.27	21	0.5	0.22		
7.28	34	1	0.15		
7.29	57	2	0.48		
7.30	2.1		0.3	1	
8.01	6.6	0.6	0.33		
8.02	2.4	0.5	0.99		
8.03	13	0.22	1		>10
8.04	8	>10	1.1		
9.01	22	0.36	1	0.6	
9.02	15	0.5	0.81		
9.03	18	0.1	0.37		
9.04	13	0.2	0.73		
9.05	22	0.36	1.6		0.6
9.06	23	3	0.4	0.3	
9.07	17	>10	0.26		
10.01	39	1	0.44		
10.02	26	0.9	1.06		
10.03	23	0.9	2.4		
11.01	9	0.7	0.85		
11.02	4.1	0.8	0.69		
11.03	26	0.41	0.1		
11.04	4.3	>10	3.2		
12.01	2.5	0.09	0.4	0.2	
12.02	1.6		0.05	2	
12.03	2.3		0.25		
12.04	1.1		0.14		
12.06	2.6				
13.01	65		0.81		
14.01	19	0.2	1.47	0.2	
14.02	190	2	1.1	8	
14.03	30	10	1.01	1	
14.04	18		0.54		
14.05	37	>10	1		
14.06	63	10	1.11		
14.07	7.5	0.2	1.4		
15.01	15	10	0.47		
15.02	21	>10	0.66		
15.03	44	2	1.67		
16.01	44	>10	4		

- 262 -

16.02	6	>10	0.6			
16.03	21	3	>10			
16.04	9.5	>10	0.92			
16B	11	3	7			
16.C	28	0.9	>10			
18.01	19	>10	1.29			
19.01	<1	0.2	0.3	9	0.2	1.41
19.02	1.6	0.13	0.38			0.91
19.03	<1	0.3	0.09			0.64
19.04	1.6	0.2	0.34			0.14
19.05	1.8	0.2	0.67	7	0.0	0.47
19.06	5	1	0.7			
19.07	2.1	0.3	0.11			
19.08	3.2	0.03	0.4	9	0.2	0.13
19.09	1.3	0.17	0.39		0.3	0.48
19.10	1.3	0.06	0.56			1.02
19.11	38	>10	2			
19.12	9	>10	0.7			0.63
19.13	2.5	0.3	1.1			
19.14	2.6	0.4	1.13			0.44
19.15	3.1	0.5	0.36			
19.16	2.3	0.7	1.1			
19.17	1	>10	0.17			
19.18	7	0.13	0.87			
19.19	5.7		0.4			
19.20	1.6	0.03	0.07			0.23
19.21	84	>10	1.71			
19.22	3.4	0.12	0.51			
19.23	6.4	0.7	0.71			
19.24	1.8	0.05	0.12			
19.25	7.2	1	0.49			0.24
19.26	6.1	0.1	0.3			
19.27	1.5	0.3	0.4			
19.28	4.8	0.1	0.12		0.3	0.46
19.29	1.9					
19.30	<1	0.06	0.1			
19.31	1.8	0.4	0.38			
19.32	1.4	0.2	0.31			
20.01	10	0.3	0.18	5	0.2	0.7
20.02	9	0.12	0.17	5	0.7	0.52
20.03	42	0.4	2.5			2.78
20.04	23	0.58	1.9			
20.05	6.8	0.87	1.46			
20.06	5	0.36	0.14		49	
20.07	3	0.1	0.05			0.38
20.08	6.8	0.17	0.05	9	0.2	
20.09	2	0.3	0.01			
20.10	2	0.1	0.02			
20.11	26	2	0.4			
20.12	9.5					
20.13	6.3		0.04			

- 263 -

20.14	33		0.32		
20.15	14	0.4	0.97	0.3	
20.16	7.5		0.06		
20.17	2		0.14		
20.18	15		0.81		
20.19	28		0.21		
20.20	3.1				0.1
20.21	2	26	3	0.68	
20.22		8	>10	0.19	
20.23		30	0.49	3	
20.24		19	0.48	2	
20.25		6.2	0.21	0.06	
20.26		5.3	0.76	0.27	
20.27		12	0.85	0.05	0.29
20.28		9.2	0.17	0.08	0.42
20.29		6.1	0.2	0.05	0.31
20.30		7.6	0.3	0.08	0.67
20.31		39		0.5	
20.32		13		0.11	
20.33		2.5		0.38	
20.34		13	1	0.12	
20.35		8.7	0.09	0.09	0.15
21.01		1	0.07	0.19	0.47
21.02		8.5	0.33	>10	
21.03		1.7	0.3	0.3	
21.04		1.8	0.05	0.3	
22.01		43	>10	>10	
22.02		26	1	3	
22.03		6.6	0.09	0.15	0.26
23.01		3.4	0.6	0.2	0.6
23.02		1.5	0.2	0.4	0.53
23.03		1.7	1	1.12	0.8
23.04		1.2	0.9	1.07	0.82
23.05		1.9	>10	0.59	0.6
23.06		16	1	0.57	
23.07		2.1	3	0.84	
23.08		6.7	0.3	0.49	
23.09		2.1	0.2	0.28	
24.01		3.6	0.11	0.44	0.05
24.02		2.1	0.5	0.11	0.39
24.03		1	0.3	1.08	
25.01		8.5	3	1	
25.02		3	0.4	0.13	0.64
26.01		4.4	0.05	0.35	0.29
26.02		1.9	0.03	0.12	0.0
26.03		1.4	0.1	0.13	0.39
26.04		4.9	0.05	0.43	0.2
26.05		2.1	0.09	0.23	1.16
26.06		4.4	0.1	0.35	1.5
26.07		11	0.5	0.95	
26.08		2.9	0.01	0.18	
26.09		2.3	0.04	0.22	
26.10		2	0.01	0.14	


26.11	4.4	0.4	0.78		0.5	
26.12	3.7	0.2	0.19			
26.13	1.6	0.2	0.44			
26.14	5		0.19			
26.15	6.9	1.2	0.08			0.07
26.16	9	0.32	2			
26.17	17	0.3	0.1	6	0.2	
26.18	1.3	6	1.17			
26.19	9.2	0.43	0.79			
26.20	10	0.14	0.22		0.6	0.49
26.21	1.1	0.1	0.49			
26.22	<1	0.1	0.28			
26.23	1.4	0.3	0.09		0.3	0.18
26.24	1	0.5	0.48		0.9	
26.25	<1	0.6	0.73		0.3	
26.26	1.9	0.2	0.07			0.34
26.27	4.8	0.6	1.49			
26.28	2.1	0.5	1.52			
26.29	<1	0.31	0.26			
26.30	4.4	1	0.76			
26.31	2	0.3	0.16			
26.32	1.6		0.05		0.6	
26.33	4		0.06	3	0.2	
26.34	7		0.1	5	0.2	
26.35	4.5		0.05		0.3	
26.36	1.9		0.07	9	0.0	
26.37	<1					
26.38	<1					
26.39	3.1					
27.01	14	0.06	0.47			
27.02	5.1	0.5	1.1			
27.03	6.3	>10	0.56			
27.04	11	0.1	0.27			
27.05	8.2	0.04	0.3			
27.06	1	0.08	0.31			
27.07	5.5	2	0.57			
27.08	9.3	0.6	0.75			
27.09	4.2	0.5	0.36			
28.01	12	0.3	0.46			0.3
28.02	1.9	0.08	0.44			3.71
28.03	7.4	0.07	0.29			
28.04	7.5	0.3	0.3			
28.05	6.7	0.1	0.12			1.39
28.06	17	0.6	0.56			
28.07	47	3	>10			
28.08	4.6	0.4	0.37			
28.09	3.1	0.5	0.36			
28.10	20	3	1.85			
28.11	4.2	0.5	0.63			
28.12	3.2	0.3	0.43			0.1
28.13	7.8	0.1	0.55	9	0.2	
28.14	3	0.1	1.44			

- 265 -

28.15	10	0.5	0.69		
28.16	11	0.11	1	0.6	
28.17	15	0.16	1.9		
28.18	9.1	>10	2.03		
28.19	3.7	0.5	0.14		
28.20	4.4	2	0.4		
28.21	1.3	0.1	0.23		
28.22	1.3	0.1	0.3		
28.23	5.9	0.5	0.28		
28.24	2.9	0.2	0.09		2.57
28.25	3.9	0.04	0.13		
28.26	6.6	0.2	0.57		
28.27	2.4	0.3	0.42	0.5	
28.28	5.2	0.4	0.52	1	
28.29	11	0.4	0.36		
28.30	2.3	0.9	0.11		
28.31	7.4	0.06	1.06		
29.01	13	0.7	2.2		0.09
29.02	3.3	0.7	1.1		
29.03	5.6	0.1	0.99		
30.01	22	0.2	0.89		
30.02	12	0.2	0.47		
30.03	19	0.5	0.68		
30.04	25	0.3	0.99		
30.05	8.5	2	0.29		
30.06	15	1	1.03		
30.07	8.8	0.6	0.47		
31.01	30	>10	1.6		
31.02	31	0.28	0.29		0.42
32.01	4.1	0.1	0.29		
32.02	5.9	0.05	0.37		0.12
33.01	2.5	0.08	0.25		
33.02	5.2	0.06	0.25		0.1
34.01	8	0.1	0.37		0.28
34.02	11	0.08	1.17		
34.03	33	0.19	2.25		
34.04	13	>10	1.22		
34.05	51	0.36	5.1		
34.06	14	>10	3		
34.07	27	>10	2.7		
34.08	8.7	>10	1.9		
35.01	6.8	>10	1.43		
35.02	6.1	0.7	0.23		
51.00	8.1	0.01	0.19		0.2
		3			
52.00	13	0.2	0.41		<0.0
				8	

Claims

1. A compound of formula I



wherein

R is selected from C₆₋₁₀aryl, C₅₋₁₀heteroaryl, C₃₋₁₂cycloalkyl and C₃₋₁₀heterocycloalkyl;
 each of R⁰, R¹, R², and R³ independently is hydrogen, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkinyl, C₃₋₈cycloalkyl, C₃₋₈cycloalkylC₁₋₈alkyl, C₅₋₁₀arylC₁₋₈alkyl, hydroxyC₁₋₈alkyl, C₁₋₈alkoxyC₁₋₈alkyl, aminoC₁₋₈alkyl, haloC₁₋₈alkyl, unsubstituted or substituted C₅₋₁₀aryl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1, 2 or 3 hetero atoms selected from N, O and S, hydroxy, C₁₋₈alkoxy, hydroxyC₁₋₈alkoxy, C₁₋₈alkoxyC₁₋₈alkoxy, haloC₁₋₈alkoxy, unsubstituted or substituted C₅₋₁₀arylC₁₋₈alkoxy, unsubstituted or substituted heterocyclloxy, or unsubstituted or substituted heterocyclylC₁₋₈alkoxy, unsubstituted or substituted amino, C₁₋₈alkylthio, C₁₋₈alkylsulfinyl, C₁₋₈alkylsulfonyl, C₅₋₁₀arylsulfonyl, halogen, carboxy, C₁₋₈alkoxycarbonyl, unsubstituted or substituted carbamoyl, unsubstituted or substituted sulfamoyl, cyano, nitro, -S(O)₀₋₂NR₁₂R₁₃, -S(O)₀₋₂R₁₃, -NR₁₂S(O)₀₋₂R₁₃, -C(O)NR₁₂R₁₃, -C(O)R₁₃ and -C(O)OR₁₃; wherein R₁₂ is selected from hydrogen and C₁₋₆alkyl; and R₁₃ is selected from hydrogen, C₁₋₆alkyl and C₃₋₁₂cycloalkyl;

or R⁰ and R¹, R¹ and R², and/or R² and R³ form, together with the carbon atoms to which they are attached, a 5 or 6 membered carbocyclic or heterocyclic ring comprising 0, 1, 2 or 3 heteroatoms selected from N, O and S;

R⁴ is hydrogen or C₁₋₈alkyl;

each of R⁵ and R⁶ independently is hydrogen, C₁₋₈alkyl, C₁₋₈alkoxyC₁₋₈alkyl, haloC₁₋₈alkyl, C₁₋₈alkoxy, halogen, carboxy, C₁₋₈alkoxycarbonyl, unsubstituted or substituted carbamoyl, cyano, or nitro;

R is unsubstituted or substituted by R₇, R₈, R₉, R₁₀, and R'₁₀;

R_7 , R_8 , R_9 , R_{10} , or R'_{10} is a substituent independently selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkinyl, C_3 - C_8 cycloalkyl, C_3 - C_8 cycloalkyl- C_1 - C_8 alkyl, C_5 - C_{10} aryl- C_1 - C_8 alkyl, hydroxy- C_1 - C_8 alkyl, C_1 - C_8 alkoxy- C_1 - C_8 alkyl, amino- C_1 - C_8 alkyl, halo- C_1 - C_8 alkyl, unsubstituted or substituted C_5 - C_{10} aryl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1, 2 or 3 hetero atoms selected from N, O and S, hydroxy, C_1 - C_8 alkoxy, hydroxy- C_1 - C_8 alkoxy, C_1 - C_8 alkoxy- C_1 - C_8 alkoxy, halo- C_1 - C_8 alkoxy, unsubstituted or substituted amino- C_1 - C_8 alkoxy, unsubstituted or substituted C_5 - C_{10} aryl- C_1 - C_8 alkoxy, unsubstituted or substituted heterocyclyloxy, or unsubstituted or substituted heterocyclyl- C_1 - C_8 alkyl, unsubstituted or substituted heterocyclyl- C_1 - C_8 alkoxy, unsubstituted or substituted amino, C_1 - C_8 alkylthio, C_1 - C_8 alkylsulfinyl, C_1 - C_8 alkylsulfonyl, C_5 - C_{10} arylsulfonyl, heterocyclosulfonyl, halogen, carboxy, C_1 - C_8 alkylcarbonyl, C_1 - C_8 alkoxycarbonyl, unsubstituted or substituted carbamoyl, unsubstituted or substituted sulfamoyl, cyano, nitro, $-S(O)_{0-2}NR_{12}R_{13}$, $-S(O)_{0-2}R_{12}$, $-C(O)R_{11}$, $-OXR_{11}$, $-NR_{12}XR_{11}$, $-NR_{12}XNR_{12}R_{13}$, $-OXNR_{12}R_{13}$, $-OXOR_{12}$ and $-XR_{11}$;

or two adjacent substituents on R may form together with the carbon atoms to which they are attached, a unsubstituted or substituted 5 or 6 membered carbocyclic or heterocyclic ring comprising 0, 1, 2 or 3 heteroatoms selected from N, O and S;

X is a bond or C_{1-6} alkylene; and

R_{11} is independently selected from C_{6-10} aryl, C_{5-10} heteroaryl, C_{3-12} cycloalkyl and C_{3-10} heterocycloalkyl;

and any aryl, heteroaryl, cycloalkyl or heterocycloalkyl of R_{11} is optionally substituted by 1 to 3 radicals independently selected from C_{1-6} alkyl, C_{3-10} heterocycloalkyl- C_{0-4} alkyl optionally substituted with C_{1-6} alkyl, $-C(O)R_{12}$, $-C(O)NR_{12}R_{13}$, $-XNR_{12}R_{13}$, $-NR_{12}XNR_{12}R_{13}$ and $-NR_{12}C(O)R_{13}$; wherein X is a bond or C_{1-6} alkylene; R_{12} and R_{13} are independently selected from hydrogen and C_{1-6} alkyl;

and salts thereof for the treatment of a disease associated to tyrosine kinase activity of anaplastic lymphoma kinase (ALK).

2. use of a compound of formula I according to claim 1 wherein

R^0 or R^2 independently is hydrogen, C_1 - C_8 alkyl, e.g. methyl, ethyl or isopropyl, hydroxy- C_1 - C_8 alkyl, e.g. hydroxyethyl or hydroxybutyl, halo- C_1 - C_8 alkyl, e.g. trifluoromethyl, unsubstituted or substituted C_5 - C_{10} aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C_1 - C_8 alkoxy, e.g. methoxy, ethoxy or isopropoxy, halo- C_1 - C_8 alkoxy, e.g. trifluoromethoxy, C_5 - C_{10} aryloxy, e.g. phenoxy,

unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclylC₁-C₈alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetylamino, C₁-C₈alkylsulfonyl, e.g. methylsulfonyl, halogen, e.g. fluoro or chloro, unsubstituted or substituted carbamoyl, e.g. cyclohexylcarbamoyl, piperidinocarbonyl, piperazinocarbonyl, N-methylpiperazinocarbonyl or morpholinocarbonyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl; preferably hydrogen, piperazino, N-methylpiperazino or 1-methyl-4-piperidyloxy, -S(O)₀₋₂NR₁₂R₁₃, -S(O)₀₋₂R₁₃, -NR₁₂S(O)₀₋₂R₁₃, -C(O)NR₁₂R₁₃, and -C(O)OR₁₃ in particular hydrogen;

R¹ is hydrogen, C₁-C₈alkyl, e.g. methyl, ethyl or isopropyl, hydroxyC₁-C₈alkyl, e.g. hydroxyethyl or hydroxybutyl, haloC₁-C₈alkyl, e.g. trifluoromethyl, unsubstituted or substituted C₅-C₁₀aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C₁-C₈alkoxy, e.g. methoxy, ethoxy or isopropoxy, haloC₁-C₈alkoxy, e.g. trifluoromethoxy, C₅-C₁₀aryloxy, e.g. phenoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclylC₁-C₈alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetylamino, C₁-C₈alkylsulfonyl, e.g. methylsulfonyl, halogen, e.g. fluoro or chloro, unsubstituted or substituted carbamoyl, e.g. cyclohexylcarbamoyl, piperidinocarbonyl, piperazinocarbonyl, N-methylpiperazinocarbonyl or morpholinocarbonyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl; preferably hydrogen, piperazino, N-methylpiperazino, morpholino, 1-methyl-4-piperidinyloxy, 3-morpholinopropoxy or 2-morpholinoethoxy, in particular hydrogen;

R³ is hydrogen, C₁-C₈alkyl, e.g. methyl or ethyl, hydroxyC₁-C₈alkyl, e.g. hydroxyethyl or hydroxybutyl, haloC₁-C₈alkyl, e.g. trifluoromethyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 heteroatoms selected from N, O and S, e.g. 2-pyrrolidinonyl or S,S-dioxoisothiazolidinyl, C₁-C₈alkoxy, e.g. methoxy, substituted amino, e.g. acetylamino, acetyl-methyl-amino, benzoylamino, methylsulfonylamino or phenylsulfonylamino, C₁-C₈alkylsulfonyl, e.g. methylsulfonyl, propyl-sulfonyl, cyclohexyl-sulfonyl, isopropyl-sulfonyl, C₅-C₁₀arylsulfonyl, e.g. phenylsulfonyl, halogen, e.g. fluoro or chloro, carboxy, substituted or unsubstituted carbamoyl, e.g. carbamoyl, methylcarbamoyl, ethyl-amino-carbonyl or dimethylcarbamoyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl,

propylsulfamoyl, isopropylsulfamoyl, isobutylsulfamoyl, cyclopropylmethyl-sulfamoyl, 2,2,2-trifluoroethylsulfamoyl, dimethylsulfamoyl or morpholinosulfonyl dimethyl-sulfamoyl, ethyl-sulfamoyl, 1-ethyl-propyl-sulfamoyl, cyclopentyl-sulfamoyl, cyclobutyl-sulfamoyl; preferably sulfamoyl, methylsulfamoyl or propylsulfamoyl;

each pair of adjacent substituents R^0 and R^1 , or R^1 and R^2 , or R^2 and R^3 are $-\text{CH}_2\text{-NH-CO-}$, $-\text{CH}_2\text{-CH}_2\text{-NH-CO-}$, $-\text{CH}_2\text{-CO-NH-}$, $-\text{CH}_2\text{-CH}_2\text{-CO-NH-}$, $-\text{CH}_2\text{-NH-SO}_2\text{-}$, $-\text{CH}_2\text{-CH}_2\text{-NH-SO}_2\text{-}$, $-\text{CH}_2\text{-SO}_2\text{-NH-}$, $-\text{CH}_2\text{-CH}_2\text{-SO}_2\text{-NH-}$, $-\text{CH}_2\text{-CH}_2\text{-SO}_2\text{-}$, $-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-SO}_2\text{-}$, $-\text{O-CH}_2\text{-O-}$, or $-\text{O-CF}_2\text{-O-}$, and such pairs wherein hydrogen in NH is replaced by $\text{C}_1\text{-C}_8$ alkyl; preferably the pair of adjacent substituents R^0 and R^1 , or R^1 and R^2 being $-\text{O-CH}_2\text{-O-}$, and the pair of adjacent substituents R^2 and R^3 being $-\text{CH}_2\text{-NH-CO-}$ or $-\text{CH}_2\text{-NH-SO}_2\text{-}$.

R^4 is hydrogen or $\text{C}_1\text{-C}_8$ alkyl, e.g. methyl; preferably hydrogen;

R^5 is hydrogen; $\text{C}_1\text{-C}_8$ alkyl, e.g. methyl or ethyl, halogen, e.g. chloro or bromo, halo $\text{C}_1\text{-C}_8$ alkyl, e.g. trifluoromethyl, cyano or nitro; preferably hydrogen, methyl, ethyl, chloro, bromo, trifluoromethyl or nitro; in particular chloro or bromo;

R^6 is hydrogen;

each of R^7 and R^9 independently is hydrogen, $\text{C}_1\text{-C}_8$ alkyl, e.g. methyl, ethyl or isopropyl, hydroxy $\text{C}_1\text{-C}_8$ alkyl, e.g. hydroxyethyl or hydroxybutyl, $\text{C}_1\text{-C}_8$ alkylcarbonyl, e.g. methyl carbonyl, aminoalkoxy, e.g. diethylaminoethoxy, halo $\text{C}_1\text{-C}_8$ alkyl, e.g. trifluoromethyl, unsubstituted or substituted $\text{C}_5\text{-C}_{10}$ aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, $\text{C}_1\text{-C}_8$ alkoxy, e.g. methoxy, ethoxy or isopropoxy, halo $\text{C}_1\text{-C}_8$ alkoxy, e.g. trifluoromethoxy, $\text{C}_5\text{-C}_{10}$ aryloxy, e.g. phenoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclyl $\text{C}_1\text{-C}_8$ alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetylamino, $\text{C}_1\text{-C}_8$ alkylsulfonyl, e.g. methylsulfonyl, heterocyclosulfonyl, e.g. piperazinylsulfonyl, heterocyclocarbonyl, e.g. methylpiperazinylcarbonyl, cyano, halogen, e.g. fluoro or chloro, unsubstituted or substituted carbamoyl, e.g. cyclohexylcarbamoyl, piperidinocarbonyl, piperazinocarbonyl, N-methylpiperazinocarbonyl or morpholinocarbonyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl; preferably hydrogen, methyl, isopropyl, trifluoromethyl, phenyl, methoxyphenyl, piperidino, piperazino, N-methylpiperazino, morpholino, methoxy, ethoxy, isopropoxy, phenoxy, 3-

morpholinopropoxy, 2-morpholinoethoxy, 2-(1-imidazolyl)ethoxy, dimethylamino, fluoro, morpholinocarbonyl, piperidinocarbonyl, piperazinocarbonyl or cyclohexylcarbonyl;

R^8 is hydrogen, C_1 - C_8 alkyl, e.g. methyl, ethyl or isopropyl, hydroxy C_1 - C_8 alkyl, e.g. hydroxyethyl or hydroxybutyl, halo C_1 - C_8 alkyl, e.g. trifluoromethyl, C_5 - C_{10} aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, heterocyclylalkyl, e.g. methylpiperazinoethyl, heterocyclylcarbonyl, e.g. piperazinocarbonyl, heterocyclyl C_1 - C_8 alkylamino, e.g. pyridylethyl(methyl)amino, C_1 - C_8 alkoxy, e.g. methoxy, ethoxy or isopropoxy, halo C_1 - C_8 alkoxy, e.g. trifluoromethoxy, C_5 - C_{10} aryloxy, e.g. phenoxy, unsubstituted or substituted heterocycliloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclyl C_1 - C_8 alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino or dimethylamino, C_1 - C_8 alkylamino- C_1 - C_8 alkylamino, e.g. dimethylamino-propylamino, C_1 - C_8 alkylsulfonyl, e.g. methylsulfonyl, halogen, e.g. fluoro or chloro, unsubstituted or substituted carbamoyl, e.g. cyclohexylcarbamoyl, piperidinocarbonyl, piperazinocarbonyl, N-methylpiperazinocarbonyl or morpholinocarbonyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl, cyano, or nitro; preferably hydrogen, methyl, piperidino, piperazino, N-methylpiperazino, morpholino, methoxy, ethoxy, trifluoromethoxy, phenoxy, 1-methyl-4-piperidyloxy, 3-morpholinopropoxy, 2-morpholinoethoxy, 3-(N-methylpiperazino)-propoxy, methylamino, fluoro, chloro, sulfamoyl or nitro;

R^{10} is hydrogen, C_1 - C_8 alkyl, e.g. methyl, ethyl or butyl, hydroxy, cyano, hydroxy C_1 - C_8 alkyl, e.g. hydroxyethyl or hydroxybutyl, halo C_1 - C_8 alkyl, e.g. trifluoromethyl, C_1 - C_8 alkoxy, e.g. methoxy or ethoxy, cycloalkylalkoxy, aryloxy, halo C_1 - C_8 alkoxy, unsubstituted or substituted heterocyclyl C_1 - C_8 alkoxy, e.g. 2-(1-imidazolyl)ethoxy, unsubstituted or substituted amino, e.g. methylamino or dimethylamino, halogen, e.g. fluoro or chloro; carboxy, carbamoyl, or unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl; preferably methyl, butyl, methoxy, ethoxy, 2-(1-imidazolyl)ethoxy, methylamino, dimethylamino or fluoro; and

each pair of adjacent substituents R^7 and R^8 , or R^8 and R^9 or R^9 and R^{10} , are $-NH-CH=CH-$, $-CH=CH-NH-$, $-NH-N=CH-$, $-CH=N-NH-$, $-CH_2-CH_2-CH_2-$, $-CH_2-CH_2-CH_2-CH_2-$, $-CH_2-CH_2-O-$, $-CH_2C(CH_3)_2O-$, $-CH=C(CH_3)O-$, $-OCH_2CH_2O-$, $-(Morpholinopropyl)N-CH=CH-$, $-CH=CH-O-$, $-O-CH_2-O-$, or $-O-CF_2-O-$; preferably the pair of adjacent substituents R^7 and R^8 or R^8

and R^9 being $-O-CH_2-O-$ or the pair of adjacent substituents R^9 and R^{10} being $-NH-CH=CH-$, $-CH=N-NH-$, $-CH_2-CH_2-CH_2-$, $-CH_2-CH_2-CH_2-CH_2-$ or $-O-CF_2-O-$.

3 use of a compound of formula I according to claim 1 or 2 wherein

R^7 , R^8 , R^9 , R^{10} and R^{10} are ethoxy, ethyl, propyl, methyl, t-butyl, trifluoromethyl, nitrile, cyclobutyloxy, 2,2,2-trifluoroethoxy, methoxy, isobutyloxy, t-butyloxy, isopropoxy, methyl-amino-carbonyl, cyclopropyl-methoxy, dimethylamino-propyl-amino, methoxy-ethoxy, $-XR_{11}$, $-C(O)R_{11}$ and $-OXR_{11}$; wherein X is a bond, methylene or ethylene; R_{11} is selected from piperaziny, piperidiny, pyrrolidiny, morpholino, azepanyl and 1,4-dioxo-8-aza-spiro[4.5]dec-8-yl; wherein R_{11} is optionally substituted by 1 to 3 radicals independently selected from methyl, isopropyl, acetyl, acetyl-methyl-amino, 3-dimethylamino-2,2-dimethyl-propylamino, ethyl-methyl-amino-ethoxy, diethyl-amino-ethoxy, amino-carbonyl, ethyl, 2-oxo-pyrrolidin-1-yl, pyrrolidiny, pyrrolidiny-methyl, piperidiny optionally substituted with methyl or ethyl, morpholino, dimethylamino, dimethylamino-propyl-amino, methyl-amino and ethyl-amino.

4. use of a compound of formula I according to claim wherein

R^0 or R^2 independently is hydrogen, C_1-C_8 alkyl, e.g. methyl, ethyl or isopropyl, halo C_1-C_8 alkyl, e.g. trifluoromethyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C_1-C_8 alkoxy, e.g. methoxy, ethoxy or isopropoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclyl C_1-C_8 alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetylamino, halogen, e.g. fluoro or chloro; preferably hydrogen, piperazino, N-methylpiperazino or 1-methyl-4-piperidyloxy, in particular hydrogen;

R^1 is hydrogen, C_1-C_8 alkyl, e.g. methyl, ethyl or isopropyl, halo C_1-C_8 alkyl, e.g. trifluoromethyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C_1-C_8 alkoxy, e.g. methoxy, ethoxy or isopropoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclyl C_1-C_8 alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetylamino, halogen, e.g. fluoro or chloro; preferably hydrogen, piperazino, N-

methylpiperazino, morpholino, 1-methyl-4-piperidinyloxy, 3-morpholinopropoxy or 2-morpholinoethoxy, in particular hydrogen;

R³ is hydrogen, C₁-C₈alkyl, e.g. methyl or ethyl, haloC₁-C₈alkyl, e.g. trifluoromethyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 heteroatoms selected from N, O and S, e.g. 2-pyrrolidonyl or S,S-dioxoisothiazolidinyl, C₁-C₈alkoxy, e.g. methoxy, substituted amino, e.g. acetamino, acetyl-methyl-amino, benzoylamino, methylsulfonylamino or phenylsulfonylamino, C₁-C₈alkylsulfonyl, e.g. methylsulfonyl, C₅-C₁₀arylsulfonyl, e.g. phenylsulfonyl, halogen, e.g. fluoro or chloro, carboxy, substituted or unsubstituted carbamoyl, e.g. carbamoyl, methylcarbamoyl or dimethylcarbamoyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl, propylsulfamoyl, isopropylsulfamoyl, isobutylsulfamoyl, cyclopropylmethyl-sulfamoyl, 2,2,2-trifluoroethylsulfamoyl, dimethylsulfamoyl or morpholinosulfonyl; preferably sulfamoyl, methylsulfamoyl or propylsulfamoyl;

each pair of adjacent substituents R⁰ and R¹, or R¹ and R², or R² and R³ are -CH₂-NH-CO-, -CH₂-NH-SO₂-, -CH₂-CH₂-SO₂-, -O-CH₂-O-, or -O-CF₂-O-, and such pairs wherein hydrogen in NH is replaced by C₁-C₈alkyl; preferably the pair of adjacent substituents R⁰ and R¹, or R¹ and R² being -O-CH₂-O-, and the pair of adjacent substituents R² and R³ being -CH₂-NH-CO- or -CH₂-NH-SO₂-.

R⁴ is hydrogen;

R⁵ is hydrogen, halogen, e.g. chloro or bromo, haloC₁-C₈alkyl, e.g. trifluoromethyl, or nitro; preferably hydrogen, chloro, bromo, trifluoromethyl or nitro; in particular chloro or bromo;

R⁶ is hydrogen;

each of R⁷ and R⁹ independently is hydrogen, C₁-C₈alkyl, e.g. methyl, ethyl or isopropyl, haloC₁-C₈alkyl, e.g. trifluoromethyl, unsubstituted or substituted C₅-C₁₀aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C₁-C₈alkoxy, e.g. methoxy, ethoxy or isopropoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclylC₁-C₈alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino, dimethylamino or acetamino, halogen, e.g. fluoro or chloro, unsubstituted or substituted carbamoyl, e.g. cyclohexylcarbamoyl, piperidinocarbonyl, piperazinocarbonyl, N-methylpiperazinocarbonyl or morpholinocarbonyl, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl; preferably hydrogen, methyl, isopropyl,


trifluoromethyl, phenyl, o-, m- or p-methoxyphenyl, piperidino, piperazino, N-methylpiperazino, morpholino, methoxy, ethoxy, isopropoxy, phenoxy, 3-morpholinopropoxy, 2-morpholinoethoxy, 2-(1-imidazolyl)ethoxy, dimethylamino, fluoro, morpholinocarbonyl, piperidinocarbonyl, piperazinocarbonyl or cyclohexylcarbonyl; R^8 is hydrogen, C_1 - C_8 alkyl, e.g. methyl, ethyl or isopropyl, halo C_1 - C_8 alkyl, e.g. trifluoromethyl, C_5 - C_{10} aryl, e.g. phenyl or methoxyphenyl, unsubstituted or substituted 5 or 6 membered heterocyclyl comprising 1 or 2 hetero atoms selected from N, O and S, e.g. morpholino, piperidino, piperazino or N-methylpiperazino, C_1 - C_8 alkoxy, e.g. methoxy, ethoxy or isopropoxy, halo C_1 - C_8 alkoxy, e.g. trifluoromethoxy, C_5 - C_{10} aryloxy, e.g. phenoxy, unsubstituted or substituted heterocyclyloxy, e.g. 1-methyl-4-piperidyloxy, unsubstituted or substituted heterocyclyl C_1 - C_8 alkoxy, e.g. 2-(1-imidazolyl)ethoxy, 3-morpholinopropoxy or 2-morpholinoethoxy, unsubstituted or substituted amino, e.g. methylamino or dimethylamino, halogen, e.g. fluoro or chloro, unsubstituted or substituted sulfamoyl, e.g. sulfamoyl, methylsulfamoyl or dimethylsulfamoyl, or nitro; preferably hydrogen, methyl, piperidino, piperazino, N-methylpiperazino, morpholino, methoxy, ethoxy, trifluoromethoxy, phenoxy, 1-methyl-4-piperidyloxy, 3-morpholinopropoxy, 2-morpholinoethoxy, 3-(N-methylpiperazino)-propoxy, methylamino, fluoro, chloro, sulfamoyl or nitro;

R^{10} is C_1 - C_8 alkyl, e.g. methyl, ethyl or butyl, halo C_1 - C_8 alkyl, e.g. trifluoromethyl, C_1 - C_8 alkoxy, e.g. methoxy or ethoxy, unsubstituted or substituted heterocyclyl C_1 - C_8 alkoxy, e.g. 2-(1-imidazolyl)ethoxy, unsubstituted or substituted amino, e.g. methylamino or dimethylamino, halogen, e.g. fluoro or chloro; preferably methyl, butyl, methoxy, ethoxy, 2-(1-imidazolyl)ethoxy, methylamino, dimethylamino or fluoro; and

each pair of adjacent substituents R^7 and R^8 , or R^8 and R^9 or R^9 and R^{10} , are $-NH-CH=CH-$, $-CH=CH-NH-$, $-NH-N=CH-$, $-CH=N-NH-$, $-CH_2-CH_2-CH_2-$, $-CH_2-CH_2-CH_2-CH_2-$, $-O-CH_2-O-$, or $-O-CF_2-O-$; preferably the pair of adjacent substituents R^7 and R^8 or R^8 and R^9 being $-O-CH_2-O-$ or the pair of adjacent substituents R^9 and R^{10} being $-NH-CH=CH-$, $-CH=N-NH-$, $-CH_2-CH_2-CH_2-$, $-CH_2-CH_2-CH_2-CH_2-$ or $-O-CF_2-O-$.

5. Use of a compound of formula I wherein the compound is selected from a compound of examples 1 to 53.

6. a compound of formula I' with the proviso that this does not include any of the compounds of examples 1 to 52 inclusive.



in which:

n' is selected from 1, 2 and 3;

R'₁ is selected from C₆₋₁₀aryl, C₅₋₁₀heteroaryl, C₃₋₁₂cycloalkyl and C₃₋₁₀heterocycloalkyl;

wherein any aryl, heteroaryl, cycloalkyl or heterocycloalkyl of R'₁ is optionally substituted by 1 to 3 radicals independently selected from C₁₋₆alkyl, C₁₋₆alkoxy, alkoxy-substituted-C₁₋₆alkyl, halo-substituted-C₁₋₆alkyl, halo-substituted-C₁₋₆alkoxy, -C(O)NR'₅R'₆, -S(O)₀₋₂NR'₅R'₆, -S(O)₀₋₂R'₅, -C(O)R'₄, -OXR'₄, -NR'₅XNR'₅R', -OXNR'₅R'₆, -OXOR'₅ and -XR'₄;

wherein X' is a bond or C₁₋₆alkylene; R'₅ is selected from hydrogen and C₁₋₆alkyl; R'₆ is selected from hydrogen, C₁₋₆alkyl and C₃₋₁₂cycloalkyl-C₁₋₄alkyl; and R'₄ is independently selected from C₆₋₁₀aryl, C₅₋₁₀heteroaryl, C₃₋₁₂cycloalkyl and C₃₋₁₀heterocycloalkyl;

and any aryl, heteroaryl, cycloalkyl or heterocycloalkyl of R'₄ is optionally substituted by 1 to 3 radicals independently selected from C₁₋₆alkyl, C₃₋₁₀heterocycloalkyl-C₀₋₄alkyl optionally substituted with C₁₋₆alkyl, -C(O)NR'₅R'₆, -XNR'₅R'₆, -NR'₅XNR'₅R'₆ and -NR'₅C(O)R'₆; wherein X is a bond or C₁₋₆alkylene; R'₅ and R'₆ are independently selected from hydrogen and C₁₋₆alkyl;

R'₂ is selected from hydrogen and halo, cyano, C₁₋₆alkyl, halo-substituted-C₁₋₆alkyl;

R'₃ is selected from halo, -S(O)₀₋₂NR'₅R'₆, -S(O)₀₋₂R'₆, -NR'₅S(O)₀₋₂R'₆, -C(O)NR'₅R'₆, -C(O)R'₆ and -C(O)OR'₆; wherein R'₅ is selected from hydrogen and C₁₋₆alkyl; and R'₆ is selected from hydrogen, C₁₋₆alkyl and C₃₋₁₂cycloalkyl;

and the pharmaceutically acceptable salts, hydrates, solvates, isomers and prodrugs thereof.

7. A compound of formula I' according to claim 6 in which:

n' is selected from 1 and 2;

R'_1 is selected from C_{6-10} aryl and C_{5-10} heteroaryl; wherein any aryl or heteroaryl of R'_1 is optionally substituted by 1 to 3 radicals independently selected from C_{1-6} alkyl, C_{1-6} alkoxy, $-C(O)NR'_5R'_6$, $-OX'R'_4$, $-C(O)R'_4$, $-NR'_5X'NR'_5R'_6$, $-OX'NR'_5R'_6$, $-OX'OR'_5$ and $-X'R'_4$; wherein X' is a bond or C_{1-6} alkylene; R'_5 is selected from hydrogen and C_{1-6} alkyl; R'_6 is selected from hydrogen, C_{1-6} alkyl and C_{3-12} cycloalkyl- C_{1-4} alkyl; and R'_4 is C_{3-10} heterocycloalkyl optionally substituted by 1 to 3 radicals independently selected from C_{1-6} alkyl, halo-substituted- C_{1-6} alkyl, C_{3-10} heterocycloalkyl- C_{0-4} alkyl optionally substituted with C_{1-6} alkyl, $-C(O)NR'_5R'_6$, $-X'NR'_5R'_6$, $-NR'_5X'NR'_5R'_6$ and $-NR'_5C(O)R'_6$; wherein X' is a bond or C_{1-6} alkylene; R'_5 and R'_6 are independently selected from hydrogen and C_{1-6} alkyl;

R'_2 is selected from hydrogen and halo;

R'_3 is selected from halo, $-S(O)_{0-2}NR'_5R'_6$, $-S(O)_{0-2}R'_6$, $-NR'_5S(O)_{0-2}R'_6$, $-C(O)NR'_5R'_6$ and $-C(O)OR'_6$; wherein R'_5 is selected from hydrogen and C_{1-6} alkyl; and R'_6 is selected from hydrogen, C_{1-6} alkyl and C_{3-12} cycloalkyl.

8. A compound of formula I' according to claim 6 or 7 in which R'_1 is selected from phenyl, pyridinyl, pyrazolyl and pyrimidinyl; wherein any aryl or heteroaryl of R'_1 is optionally substituted by 1 to 3 radicals independently selected from ethoxy, ethyl, propyl, methyl, t-butyl, trifluoromethyl, nitrile, cyclobutyloxy, 2,2,2-trifluoroethoxy, methoxy, isobutyloxy, t-butyloxy, isopropoxy, methyl-amino-carbonyl; cyclopropyl-methoxy, dimethylamino-propyl-amino, methoxy-ethoxy, $-X'R'_4$, $-C(O)R'_4$ and $-OX'R'_4$; wherein X' is a bond, methylene or ethylene; R'_4 is selected from piperazinyl, piperidinyl, pyrrolidinyl, morpholino, azepanyl and 1,4-dioxo-8-aza-spiro[4.5]dec-8-yl; wherein R'_4 is optionally substituted by 1 to 3 radicals independently selected from methyl, isopropyl, acetyl, acetyl-methyl-amino, 3-dimethylamino-2,2-dimethyl-propylamino, ethyl-methyl-amino-ethoxy, diethyl-amino-ethoxy, amino-carbonyl, ethyl, 2-oxo-pyrrolidin-1-yl, pyrrolidinyl, pyrrolidinyl-methyl, piperidinyl optionally substituted with methyl or ethyl, morpholino, dimethylamino, dimethylamino-propyl-amino, methyl-amino and ethyl-amino.

9. A compound of formula I' according to claim 6, 7 or 8 in which R'_2 is selected from hydrogen and halo; and R'_3 is selected from halo, dimethyl-sulfamoyl, isobutyl-sulfamoyl, methyl-sulfamoyl, ethyl-sulfamoyl, propyl-sulfonyl, ethyl-amino-carbonyl, 1-ethyl-propyl-

sulfamoyl, cyclopentyl-sulfamoyl, isopropyl-sulfamoyl, cyclohexyl-sulfonyl, cyclopropyl-methyl-sulfamoyl, cyclobutyl-sulfamoyl, isopropyl-sulfonyl,

10. A compound of formula I according to any one of claim 6 to 9 wherein the compound is a compound of example 53.

11. A pharmaceutical composition comprising a compound according to any one of claims 1 to 9, as active ingredient together with one or more pharmaceutically acceptable diluents or carriers.

12. The use of a compound according to any one of claims 1 to 9 for the manufacture of a medicament for the treatment or prevention of neoplastic diseases and immune system disorders.

13. A combination comprising a therapeutically effective amount a compound according to any one of claims 1 to 9 and one or more further drug substances, said further drug substance being useful in the treatment of neoplastic diseases or immune system disorders.

14. A method for the treatment of neoplastic diseases and immune system disorders in a subject in need thereof which comprises administering an effective amount of a compound according to any one of claims 1 to 9 or a pharmaceutical composition comprising same.

15. Use of a compound according to any one of claims 1 to 9 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment or prevention of a disease which responds to inhibition of FAK and/or ALK and/or ZAP-70 and/or IGF-IR.

16. The use according to claim 15, wherein the disease to be treated is selected from proliferative disease .

17. The use according to claim 16, wherein the proliferative disease to be treated is selected from a tumor of, breast, renal , prostate, colorectal, thyroid, ovarian, pancreas, neuronal, lung, uterine and gastro-intestinal tumours as well as osteosarcomas and melanomas.

18. The use according to claim 15, wherein the disease to be treated is an immune disease.

19. Use of a compound according to any one of claims 1 to 9 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment or prevention of inflammatory and/or an immune disorder.

20. Use according to claim 19 wherein the inflammatory and/or immune disorder is selected from transplant rejection, allergy and autoimmune disorders mediated by immune cells including T lymphocytes, B lymphocytes, macrophages, dendritic cells, mast cells and eosinophils.

21. The use according to any one of claims 14 to 19, wherein the compound is 2-[5-Bromo-2-(2-methoxy-5-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzenesulfonamide or 5-Chloro-N²*-[2-methoxy-4-[4-(4-methyl-piperazin-1-yl)-piperidin-1-yl]-phenyl]-N⁴*-[2-(propane-2-sulfonyl)-phenyl]-pyrimidine-2,4-diamine or a pharmaceutically acceptable salt thereof.

22. The use according to any one of claims 14 to 19, wherein the compound is selected from 2-[5-chloro-2-(2-methoxy-4-morpholin-4-yl-phenylamino)-pyrimidin-4-ylamino]-N-methyl-benzamide, N²-(4-[1,4']Bipiperidiny-1'-yl-2-methoxy-phenyl)-5-chloro-N⁴-[2-(propane-1-sulfonyl)-phenyl]-pyrimidine-2,4-diamine and 2-[5-Chloro-2-[2-methoxy-4-(4-methyl-piperazin-1-yl)-phenylamino]-pyrimidin-4-ylamino]-N-isopropyl-benzenesulfonamide, or 5-Chloro-N²*-[2-methoxy-4-[4-(4-methyl-piperazin-1-yl)-piperidin-1-yl]-phenyl]-N⁴*-[2-(propane-2-sulfonyl)-phenyl]-pyrimidine-2,4-diamine a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/009099

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/48 C07D405/12 C07D403/12 C07D401/12 C07D401/14
A61K31/506 A61K35/00 A61P37/00 A61P29/00 C07D401/14
C07D405/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BEILSTEIN Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 03/018021 A1 (AMGEN INC., USA) 6 March 2003 (2003-03-06) page 186, line 15 - page 189, line 8; examples 66,67,70,72,74,76,79,81,83,97,105,107,109, 116	11-20
X	EP 1 054 004 A (YAMANOUCHI PHARMACEUTICAL CO., LTD., JAPAN) 22 November 2000 (2000-11-22) table 5, Ex.2,8,34,35,37,39,40. tables 7 and 9 page 12, line 45, paragraph 56; claims 1,7 ----- -/--	11-14, 19,20



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

11 November 2004

Date of mailing of the international search report

22/11/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Schuemacher, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2004/009099

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GHOSH D: "2,4-BIS(ARYLAMINO)-6-METHYL PYRIMIDINES AS ANTIMICROBIAL AGENTS" JOURNAL OF THE INDIAN CHEMICAL SOCIETY, vol. 58, no. 5, May 1981 (1981-05), pages 512-513, XP000918018 ISSN: 0019-4522 table I; compound V -----	6,11
X	WO 01/60816 A1 (AMGEN INC., USA) 23 August 2001 (2001-08-23) page 18, line 22 - line 30; claims 1,19-24; examples 37,42-44,54,56-60 -----	1-22
X	WO 00/39101 A (BREAU LT GLORIA ANNE ; PEASE JANET ELIZABETH (GB); ASTRAZENECA UK LTD () 6 July 2000 (2000-07-06) page 2, line 1 - line 9; claims 1,12-14; examples 191,192,196,199-203,213,218,219 -----	6-20
X	WO 97/19065 A (CELLTECH THERAPEUTICS LTD ; DAVIS PETER DAVID (GB); MOFFAT DAVID FESTU) 29 May 1997 (1997-05-29) page 1, line 35 - page 2, line 4; claim 1; example 125 -----	6-20
A	WO 01/65655 A (HYDE DOUGLAS) 7 September 2001 (2001-09-07) page 2, line 1 - line 9; claim 1 -----	6-22
X	WO 01/64656 A1 (ASTRAZENECA AB, SWED.; ASTRAZENECA UK LTD.) 7 September 2001 (2001-09-07) claims 1,14; example 29 -----	6-20
X	WO 00/12485 A1 (ZENECA LIMITED, UK) 9 March 2000 (2000-03-09) page 1, line 30 - page 2, line 7; claims 1,10; examples 6,9-11,106 -----	11-20
X	WO 03/030909 A1 (BAYER CORPORATION, USA) 17 April 2003 (2003-04-17) examples -----	11-14, 19,20
X	WO 03/063794 A2 (RIGEL PHARMACEUTICALS, INC., USA) 7 August 2003 (2003-08-07) examples 7.3; claims 1,47,52,57-62 -----	11-14, 19,20
X	WO 99/50250 A1 (JANSSEN PHARMACEUTICA N.V., BELG.; ET AL.) 7 October 1999 (1999-10-07) claim 1; tables 5,6 -----	11-14, 19,20
	----- -/--	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/009099

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 03/066601 A1 (SMITHKLINE BEECHAM CORPORATION, USA) 14 August 2003 (2003-08-14) claims 1,23,27,28; example 48 -----	11-14, 19,20
X	EP 1 184 376 A (YAMANOUCHI PHARMACEUTICAL CO., LTD., JAPAN) 6 March 2002 (2002-03-06) page 9, lines 50-58; claim 1; table 10 -----	11-14, 19,20
P,X	WO 03/078404 A1 (NOVARTIS A.-G., SWITZ.; NOVARTIS PHARMA G.M.B.H.) 25 September 2003 (2003-09-25) claims 1,4-10; examples 2,7,10-12,16,30,51,236 -----	6-20
P,X	WO 2004/002964 A1 (YAMANOUCHI PHARMACEUTICAL CO., LTD., JAPAN) 8 January 2004 (2004-01-08) examples 79-86,167,168; tables 4,17-20 -----	11-14, 19,20
P,X	WO 03/095448 A (CHEN JIANQING ; LIU DONGLEI (US); WOOD JILL E (US); CHEN YUANWEI (US);) 20 November 2003 (2003-11-20) claim 6; tables 1,2 -----	11-14, 19,20
E	WO 2004/074244 A2 (SMITHKLINE BEECHAM CORPORATION, USA) 2 September 2004 (2004-09-02) the whole document -----	6-14,19, 22
E	WO 2004/080980 A1 (NOVARTIS A.-G., SWITZ.; NOVARTIS PHARMA G.M.B.H.) 23 September 2004 (2004-09-23) the whole document -----	6-22

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2004/009099

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 14 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/009099

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03018021	A1	06-03-2003	US 2004063705 A1 CA 2457838 A1 EP 1427421 A1	01-04-2004 06-03-2003 16-06-2004
EP 1054004	A	22-11-2000	AU 1507199 A EP 1054004 A1 US 6432963 B1 WO 9931073 A1	05-07-1999 22-11-2000 13-08-2002 24-06-1999
WO 0160816	A1	23-08-2001	AU 3704101 A CA 2400447 A1 CN 1429222 T EP 1257546 A1 HU 0301117 A2 JP 2003532635 T US 2003199534 A1 US 2002052386 A1 ZA 200206386 A	27-08-2001 23-08-2001 09-07-2003 20-11-2002 29-12-2003 05-11-2003 23-10-2003 02-05-2002 26-11-2003
WO 0039101	A	06-07-2000	AT 277020 T AU 763091 B2 AU 1874300 A BR 9916590 A CA 2352896 A1 CN 1335838 T DE 69920509 D1 EP 1140860 A1 WO 0039101 A1 JP 2002533446 T NO 20013038 A NZ 512118 A US 6593326 B1 ZA 200104413 A	15-10-2004 10-07-2003 31-07-2000 23-10-2001 06-07-2000 13-02-2002 28-10-2004 10-10-2001 06-07-2000 08-10-2002 22-08-2001 29-08-2003 15-07-2003 29-08-2002
WO 9719065	A	29-05-1997	AU 7631496 A DE 69627179 D1 DE 69627179 T2 EP 0862560 A1 ES 2195020 T3 WO 9719065 A1 US 6235746 B1 US 5958935 A	11-06-1997 08-05-2003 29-01-2004 09-09-1998 01-12-2003 29-05-1997 22-05-2001 28-09-1999
WO 0165655	A	07-09-2001	US 6376770 B1 AU 3354901 A WO 0165655 A2 CA 2401436 A1 EP 1260006 A2 EP 1376800 A1 US 2003124905 A1	23-04-2002 12-09-2001 07-09-2001 07-09-2001 27-11-2002 02-01-2004 03-07-2003
WO 0164656	A1	07-09-2001	AU 3397901 A BR 0108879 A CA 2398887 A1 CN 1416423 T EP 1278735 A1 JP 2003525279 T MX PA02007581 A	12-09-2001 29-04-2003 07-09-2001 07-05-2003 29-01-2003 26-08-2003 13-12-2002

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/009099

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0164656	A1	NO 20024126 A NZ 520502 A US 2003181474 A1 ZA 200206192 A	29-08-2002 28-05-2004 25-09-2003 26-11-2003
WO 0012485	A1	09-03-2000 AU 5438299 A EP 1107957 A1 JP 2002523497 T	21-03-2000 20-06-2001 30-07-2002
WO 03030909	A1	17-04-2003 NONE	
WO 03063794	A2	07-08-2003 CA 2474277 A1 EP 1471915 A2 US 2004029902 A1 EP 1468510 A2 EP 1468511 A2 EP 1467646 A1 WO 03063333 A1 WO 03061449 A1 WO 03063390 A2 WO 03063391 A2 US 2003137264 A1 US 2003137263 A1 US 2003142982 A1 US 2003170033 A1 WO 2004014382 A1	07-08-2003 03-11-2004 12-02-2004 20-10-2004 20-10-2004 20-10-2004 31-07-2003 31-07-2003 31-07-2003 31-07-2003 24-07-2003 24-07-2003 31-07-2003 11-09-2003 19-02-2004
WO 9950250	A1	07-10-1999 AT 232521 T AU 751573 B2 AU 3599699 A BG 104738 A BR 9909191 A CA 2324919 A1 CN 1295564 T DE 69905306 D1 DE 69905306 T2 DK 945443 T3 EA 2973 B1 EE 200000532 A EP 1245567 A1 EP 0945443 A1 ES 2193660 T3 HR 20000620 A1 HU 0101204 A2 ID 26291 A JP 3507917 B2 JP 2002509920 T NO 20004810 A NZ 506679 A OA 11492 A PL 343196 A1 PT 945443 T SI 945443 T1 SK 14062000 A3 TR 200002760 T2 TW 531534 B US 2003083317 A1 US 6197779 B1	15-02-2003 22-08-2002 18-10-1999 30-04-2001 05-12-2000 07-10-1999 16-05-2001 20-03-2003 27-11-2003 02-06-2003 26-12-2002 15-02-2002 02-10-2002 29-09-1999 01-11-2003 30-06-2001 28-10-2001 14-12-2000 15-03-2004 02-04-2002 26-09-2000 26-11-2002 06-05-2004 30-07-2001 30-06-2003 31-08-2003 11-06-2001 21-12-2000 11-05-2003 01-05-2003 06-03-2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP2004/009099

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9950250	A1	US 2001011094 A1 EP 0945442 A1 ZA 200006044 A	02-08-2001 29-09-1999 26-10-2001
WO 03066601	A1	14-08-2003 CA 2476281 A1 EP 1472233 A1	14-08-2003 03-11-2004
EP 1184376	A	06-03-2002 AU 5107900 A EP 1184376 A1 US 6797706 B1 WO 0075113 A1 JP 2001055378 A	28-12-2000 06-03-2002 28-09-2004 14-12-2000 27-02-2001
WO 03078404	A1	25-09-2003 NONE	
WO 2004002964	A1	08-01-2004 NONE	
WO 03095448	A	20-11-2003 WO 03095448 A1 WO 2004046118 A2	20-11-2003 03-06-2004
WO 2004074244	A2	02-09-2004 NONE	
WO 2004080980	A1	23-09-2004 NONE	